

β -openness and β -compactness in graded ditopological texture spaces

RAMAZAN EKMEKÇİ

Received 07 April 2025; Revised 26 April 2025; Accepted 14 June 2025

ABSTRACT. In this work, β -openness and β -compactness in graded ditopological texture spaces are defined and the properties of these concepts are examined. Furthermore, several relationships between the structures ditopological texture spaces and graded ditopological texture spaces in the context of β -openness are studied.

2020 AMS Classification: 54A05, 54D30, 54C08

Keywords: β -continuity, β -compactness, Graded ditopology, Texture.

Corresponding Author: Ramazan Ekmekçi (rekmekci@klu.edu.tr)

1. INTRODUCTION

Abd El Monsef *et al.* introduced β -open sets in [1] and several studies have been developed this area such as [2, 3, 4].

In [5] Brown and Šostak introduced the theory of graded ditopology as a more comprehensive structure than ditopology presented in [6, 7] and fuzzy topology given independently by Šostak in [8] and Kubiak in [9]. In the structure of graded ditopology, openness and closedness are defined by independent grading functions instead of elements of a texture as in ditopological case. The theory of graded ditopological texture spaces continues to be developed by various recent studies such as [10, 11].

The purpose of this study is to generalize the concepts of β -openness and β -compactness in ditopological texture spaces defined in [12] to the structure of graded ditopological texture spaces and investigate the properties of these concepts in this structure. For this generalization, the spectral approach as in [13, 14, 15] is used and several relationships between the structures ditopological texture spaces and graded ditopological texture spaces in the context of β -openness are examined.

Our basic motivation is to define β -openness and β -compactness in g.d.t.s. and so develop the theory of graded ditopologies by investigating the properties of β -openness and β -compactness in g.d.t.s.

2. PRELIMINARIES

Ditopological Texture Spaces ([6, 16, 17]) Let S be a set and $\mathcal{S} \subseteq \mathcal{P}(S)$ with $S, \emptyset \in \mathcal{S}$. If \mathcal{S} is a point separating, complete, completely distributive lattice with respect to inclusion and for which meet \bigwedge coincides with intersection \bigcap and finite joins \bigvee coincide with unions \bigcup then the pair (S, \mathcal{S}) is called a *texture* or a *texture space*.

In general, a texturing of S may not be closed under set complementation. However, if there is a mapping $\sigma : \mathcal{S} \rightarrow \mathcal{S}$ satisfying $\sigma(\sigma(A)) = A$ and $A \subseteq B \Rightarrow \sigma(B) \subseteq \sigma(A)$ for all $A, B \in \mathcal{S}$ then σ is called a *complementation* on (S, \mathcal{S}) and (S, \mathcal{S}, σ) is called a *complemented texture*.

The p -sets given by $P_s = \bigcap \{A \in \mathcal{S} \mid s \in A\}$ and the q -sets given by $Q_s = \bigvee \{A \in \mathcal{S} \mid s \notin A\} = \bigvee \{P_u \mid u \in S, s \notin P_u\}$ are essential to define several concepts in a texture space (S, \mathcal{S}) .

Product of textures ([7, 16, 17]) Let (S_j, \mathcal{S}_j) , $j \in J$ be textures, $S = \prod_{j \in J} S_j$ and $A_k \in \mathcal{S}_k$ for some $k \in J$. If we write

$$E(k, A_k) = \prod_{j \in J} Y_j \text{ where } Y_j = \begin{cases} A_j & \text{if } j = k \\ S_j & \text{otherwise} \end{cases}$$

then the product texturing $\mathcal{S} = \bigotimes_{j \in J} \mathcal{S}_j$ of S consists of arbitrary intersections of elements of the set

$$\varepsilon = \left\{ \bigcup_{j \in J_1} E(j, A_j) \mid J_1 \subseteq J, A_j \in \mathcal{S}_j \text{ for } j \in J_1 \right\}.$$

Consider two textures (S, \mathcal{S}) and (V, \mathcal{V}) . The p -sets and q -sets of the product texture $(S \times V, \mathcal{P}(S) \otimes \mathcal{V})$ will be denoted by $\bar{P}_{(s,v)}$, $\bar{Q}_{(s,v)}$ respectively.

If $\mathcal{P}(X)$ is the power set of a set X , then $(X, \mathcal{P}(X))$ is the discrete texture on X . For $x \in X$, $P_x = \{x\}$ and $Q_x = X \setminus \{x\}$. The mapping $\pi_X : \mathcal{P}(X) \rightarrow \mathcal{P}(X)$, $\pi_X(Y) = X \setminus Y$ for $Y \subseteq X$ is a complementation on the texture $(X, \mathcal{P}(X))$.

Definition 2.1 ([16]). Let (S, \mathcal{S}) and (V, \mathcal{V}) be textures. Then

- (i) $r \in \mathcal{P}(S) \otimes \mathcal{V}$ is called a *relation* on (S, \mathcal{S}) to (V, \mathcal{V}) , if it satisfies
 - R1 $r \not\subseteq \bar{Q}(s, v), P_{s'} \not\subseteq Q_s \Rightarrow r \not\subseteq \bar{Q}(s', v)$,
 - R2 $r \not\subseteq \bar{Q}(s, v) \Rightarrow \exists s' \in S$ such that $P_{s'} \not\subseteq Q_s$ and $r \not\subseteq \bar{Q}(s', v)$,
- (ii) $R \in \mathcal{P}(S) \otimes \mathcal{V}$ is called a *co-relation* on (S, \mathcal{S}) to (V, \mathcal{V}) , if it satisfies
 - CR1 $\bar{P}(s, v) \not\subseteq R, P_s \not\subseteq Q_{s'} \Rightarrow \bar{P}(s', v) \not\subseteq R$,
 - CR2 $\bar{P}(s, v) \not\subseteq R \Rightarrow \exists s' \in S$ such that $P_{s'} \not\subseteq Q_s$ and $\bar{P}(s', v) \not\subseteq R$,
- (iii) A pair (r, R) , where r is a relation and R a co-relation on (S, \mathcal{S}) to (V, \mathcal{V}) is called a *direlation* on (S, \mathcal{S}) to (V, \mathcal{V}) .

For a texture (S, \mathcal{S}) the identity direlation $(i_{(S,S)}, I_{(S,S)})$ is defined by

$$i_{(S,S)} = \bigvee \{\bar{P}(s, s) \mid s \in S\} \text{ and } I_{(S,S)} = \bigcap \{\bar{Q}(s, s) \mid s \in S^b\}.$$

For $A \subseteq S$, $r \rightarrow A = \bigcap \{Q_v \mid \forall s, r \not\subseteq \bar{Q}_{(s,v)} \Rightarrow A \subseteq Q_s\}$ is called the *A-section* of r and $R \rightarrow A = \bigvee \{P_v \mid \forall s, \bar{P}_{(s,v)} \not\subseteq R \Rightarrow P_s \subseteq A\}$ is called the *A-section* of R .

For $B \subseteq V$, $r \leftarrow B = \bigvee \{P_s \mid \forall v, r \not\subseteq \bar{Q}_{(s,v)} \Rightarrow P_v \subseteq B\}$ is called the *B-presection* of r and $R \leftarrow B = \bigcap \{Q_s \mid \forall v, \bar{P}_{(s,v)} \not\subseteq R \Rightarrow B \subseteq Q_v\}$ is called the *B-presection* of R .

Proposition 2.2 ([16]). If (r, R) is a direlation on (S, \mathcal{S}) to (V, \mathcal{V}) , then $r^\rightarrow(\bigvee_{i \in I} A_i) = \bigvee_{i \in I} r^\rightarrow A_i$, $R^\rightarrow(\bigcap_{i \in I} A_i) = \bigcap_{i \in I} R^\rightarrow A_i$, $r^\leftarrow(\bigcap_{j \in J} B_j) = \bigcap_{j \in J} r^\leftarrow B_j$ and $R^\leftarrow(\bigvee_{j \in J} B_j) = \bigvee_{j \in J} R^\leftarrow B_j$ for any $A_i \in \mathcal{S}$, $B_j \in \mathcal{V}$, $i \in I$, $j \in J$.

Definition 2.3 ([16]). A direlation (f, F) from (S, \mathcal{S}) to (V, \mathcal{V}) is called a *difunction* from (S, \mathcal{S}) to (V, \mathcal{V}) , if it satisfies the following two conditions:

- (DF1) For $s, s' \in S$, $P_s \not\subseteq Q_{s'} \Rightarrow \exists v \in V$ with $f \not\subseteq \overline{Q}_{(s,v)}$ and $\overline{P}_{(s',v)} \not\subseteq F$.
 - (DF2) For $v, v' \in V$ and $s \in S$, $f \not\subseteq \overline{Q}_{(s,v)}$ and $\overline{P}_{(s,v')} \not\subseteq F \Rightarrow P_{v'} \not\subseteq Q_v$.
- (f, F) is called *surjective*, if $\forall v, v' \in V$ $P_v \not\subseteq Q_{v'} \Rightarrow \exists s \in S$ with $f \not\subseteq \overline{Q}_{(s,v')}$ and $\overline{P}_{(s,v)} \not\subseteq F$.

In particular, the identity direlation (i_S, I_S) is a difunction on (S, \mathcal{S}) .

- Proposition 2.4** ([16]).
- (1) $f^\leftarrow B = F^\leftarrow B$ for each $B \in \mathcal{V}$.
 - (2) $f^\leftarrow \emptyset = F^\leftarrow \emptyset = \emptyset$ and $f^\leftarrow V = F^\leftarrow V = S$.
 - (3) $f^\leftarrow(F^\rightarrow A) \subseteq A \subseteq F^\leftarrow(f^\rightarrow A)$ and $f^\rightarrow(F^\leftarrow B) \subseteq B \subseteq F^\rightarrow(f^\leftarrow B)$ for all $A \in \mathcal{S}$, $B \in \mathcal{V}$.
 - (4) If (f, F) is surjective, then $F^\rightarrow(f^\leftarrow B) = B = f^\rightarrow(F^\leftarrow B)$ for all $B \in \mathcal{V}$.

A ditopology on a texture (S, \mathcal{S}) is a pair (τ, κ) , where $\tau, \kappa \subseteq \mathcal{S}$ and the set of open sets τ satisfies

- (T₁) $S, \emptyset \in \tau$,
- (T₂) $G_1, G_2 \in \tau \Rightarrow G_1 \cap G_2 \in \tau$,
- (T₃) $G_i \in \tau, i \in I \Rightarrow \bigvee_i G_i \in \tau$

and the set of closed sets κ satisfies

- (CT₁) $S, \emptyset \in \kappa$,
- (CT₂) $K_1, K_2 \in \kappa \Rightarrow K_1 \cup K_2 \in \kappa$,
- (CT₃) $K_i \in \kappa, i \in I \Rightarrow \bigcap_i K_i \in \kappa$.

In this case, $(S, \mathcal{S}, \tau, \kappa)$ is called a *ditopological texture space* (or "d.t.s." for short). So a ditopology can be considered as a "topology" in which there is no need to exist a relation between the open and closed sets [6].

Let $(S, \mathcal{S}, \tau, \kappa)$ be a d.t.s. For a subset $A \in \mathcal{S}$, the closure (interior) of A is defined by $[A] = \bigcap \{B \in \kappa \mid A \subseteq B\}$ ($]A[= \bigvee \{B \in \tau \mid B \subseteq A\}$) respectively [17]. $A \in \mathcal{S}$ is called *semi open* (*semi closed*), if $A \subseteq []A[$ ($]A[\subseteq A$) respectively [18]. $A \in \mathcal{S}$ is called β -open (β -closed), if $A \subseteq []A[$ ($]A[\subseteq A$) respectively [12].

Definition 2.5 ([19]). Let $(S, \mathcal{S}, \tau, \kappa)$ be a d.t.s. and $A \in \mathcal{S}$.

- (i) A is called *compact*, if whenever $\{G_i \mid i \in I\}$ is an open cover of A (i.e. $\forall i \in I$ $G_i \in \tau$ and $A \subseteq \bigvee_{i \in I} G_i$) then there is a finite subset J of I with $A \subseteq \bigvee_{i \in J} G_i$. In particular, $(S, \mathcal{S}, \tau, \kappa)$ is called *compact*, if S is compact.
- (ii) A is called *cocompact*, if whenever $\{K_i \mid i \in I\}$ is a closed cocover of A (i.e. $\forall i \in I$ $K_i \in \kappa$ and $\bigcap_{i \in I} K_i \subseteq A$) then there is a finite subset J of I with $\bigcap_{i \in J} K_i \subseteq A$. In particular, $(S, \mathcal{S}, \tau, \kappa)$ is called *cocompact*, if \emptyset is compact.
- (iii) $(S, \mathcal{S}, \tau, \kappa)$ is called *stable*, if every $K \in \kappa$ with $K \neq S$ is compact.
- (iv) $(S, \mathcal{S}, \tau, \kappa)$ is called *costable*, if every $G \in \tau$ with $G \neq \emptyset$ is cocompact.
- (v) $(S, \mathcal{S}, \tau, \kappa)$ is called *dicompact*, if it is compact, cocompact, stable and costable.

$A \in \mathcal{S}$ is called β -compact (β -cocompact), if every cover (cocover) of A by β -open (β -closed) sets has a finite subcover (subcocover) respectively and $(S, \mathcal{S}, \tau, \kappa)$ is called β -compact (β -cocompact), if S is β -compact (if \emptyset is β -cocompact) respectively [12].

Let $(S_k, \mathcal{S}_k, \tau_k, \kappa_k)$, $k = 1, 2$ be d.t.s. and $(f, F) : (S_1, \mathcal{S}_1) \rightarrow (S_2, \mathcal{S}_2)$ be a difunction. (f, F) is called β -continuous ($M\beta$ -continuous), if $F^{\leftarrow}(G)$ is β -open in $(S_1, \mathcal{S}_1, \tau_1, \kappa_1)$ for every open (β -open) set G in $(S_2, \mathcal{S}_2, \tau_2, \kappa_2)$. (f, F) is called β -cocontinuous ($M\beta$ -cocontinuous), if $f^{\leftarrow}(K)$ is β -closed in $(S_1, \mathcal{S}_1, \tau_1, \kappa_1)$ for every closed (β -closed) set K in $(S_2, \mathcal{S}_2, \tau_2, \kappa_2)$. (f, F) is called β -bicontinuous ($M\beta$ -bicontinuous), if it is both β -continuous and β -cocontinuous (both $M\beta$ -continuous and $M\beta$ -cocontinuous) respectively [12].

Graded Ditopological Texture Spaces [5] Consider two textures (S, \mathcal{S}) and (V, \mathcal{V}) . A graded ditopological texture space (or "g.d.t.s." for short) is a tuple $(S, \mathcal{S}, \mathcal{T}, \mathcal{K}, V, \mathcal{V})$ where the mappings $\mathcal{T}, \mathcal{K} : \mathcal{S} \rightarrow \mathcal{V}$ satisfy following conditions:

- (GT₁) $\mathcal{T}(S) = \mathcal{T}(\emptyset) = V$,
- (GT₂) $\mathcal{T}(A_1) \cap \mathcal{T}(A_2) \subseteq \mathcal{T}(A_1 \cap A_2) \quad \forall A_1, A_2 \in \mathcal{S}$,
- (GT₃) $\bigcap_{j \in J} \mathcal{T}(A_j) \subseteq \mathcal{T}(\bigcup_{j \in J} A_j) \quad \forall A_j \in \mathcal{S}, j \in J$,
- (GCT₁) $\mathcal{K}(S) = \mathcal{K}(\emptyset) = V$,
- (GCT₂) $\mathcal{K}(A_1) \cap \mathcal{K}(A_2) \subseteq \mathcal{K}(A_1 \cup A_2) \quad \forall A_1, A_2 \in \mathcal{S}$,
- (GCT₃) $\bigcap_{j \in J} \mathcal{K}(A_j) \subseteq \mathcal{K}(\bigcap_{j \in J} A_j) \quad \forall A_j \in \mathcal{S}, j \in J$.

In this case, \mathcal{T} is called a (V, \mathcal{V}) -graded topology and \mathcal{K} a (V, \mathcal{V}) -graded cotopology on (S, \mathcal{S}) . For $v \in V$ it is defined that

$$\mathcal{T}^v = \{A \in \mathcal{S} \mid P_v \subseteq \mathcal{T}(A)\}, \quad \mathcal{K}^v = \{A \in \mathcal{S} \mid P_v \subseteq \mathcal{K}(A)\}.$$

So $(\mathcal{T}^v, \mathcal{K}^v)$ is a ditopology on (S, \mathcal{S}) for each $v \in V$. Namely, if $(S, \mathcal{S}, \mathcal{T}, \mathcal{K}, V, \mathcal{V})$ is a g.d.t.s., then there exists a d.t.s. $(S, \mathcal{S}, \mathcal{T}^v, \mathcal{K}^v)$ for each $v \in V$.

$[A]^v$ and $]A[^v$ stand for the closure and the interior of a set $A \in \mathcal{S}$ in the d.t.s. $(S, \mathcal{S}, \mathcal{T}^v, \mathcal{K}^v)$ respectively, so we have $[A]^v = \bigcap \{B \in \mathcal{S} \mid A \subseteq B, B \in \mathcal{K}^v\}$, $]A[^v = \bigcup \{B \in \mathcal{S} \mid B \subseteq A, B \in \mathcal{T}^v\}$.

Let (S, \mathcal{S}, σ) be a complemented texture. If $(S, \mathcal{S}, \mathcal{T}, \mathcal{K}, V, \mathcal{V})$ is a g.d.t.s. then $(S, \mathcal{S}, \mathcal{K} \circ \sigma, \mathcal{T} \circ \sigma, V, \mathcal{V})$ is also a g.d.t.s. Additionally, $(\mathcal{T}, \mathcal{K})$ is called *complemented*, if $(\mathcal{T}, \mathcal{K}) = (\mathcal{K} \circ \sigma, \mathcal{T} \circ \sigma)$ and in this case, we say that $(S, \mathcal{S}, \mathcal{T}, \mathcal{K}, \sigma, V, \mathcal{V})$ is a *complemented g.d.t.s.*

Let $(S_k, \mathcal{S}_k, \mathcal{T}_k, \mathcal{K}_k, V_k, \mathcal{V}_k)$, $k = 1, 2$ be g.d.t.s., $(f, F) : (S_1, \mathcal{S}_1) \rightarrow (S_2, \mathcal{S}_2)$, $(h, H) : (V_1, \mathcal{V}_1) \rightarrow (V_2, \mathcal{V}_2)$ difunctions. For the pair $((f, F), (h, H))$, (f, F) is called *continuous with respect to (h, H)* , if $H^{\leftarrow} \mathcal{T}_2(A) \subseteq \mathcal{T}_1(F^{\leftarrow} A) \quad \forall A \in \mathcal{S}_2$, and *cocontinuous with respect to (h, H)* , if $h^{\leftarrow} \mathcal{K}_2(A) \subseteq \mathcal{K}_1(f^{\leftarrow} A) \quad \forall A \in \mathcal{S}_2$. If (f, F) is continuous and cocontinuous with respect to (h, H) , then it is said to be a *bicontinuous difunction with respect to (h, H)* .

Example 2.6 ([5]). Consider the discrete texture $(V, \mathcal{V}) = (1, \mathcal{P}(1))$ (The notation 1 denotes the set $\{0\}$) and take a d.t.s. $(S, \mathcal{S}, \tau, \kappa)$. Then the mappings $\tau^g, \kappa^g : \mathcal{S} \rightarrow \mathcal{P}(1)$ defined by $\tau^g(A) = 1 \Leftrightarrow A \in \tau$ and $\kappa^g(A) = 1 \Leftrightarrow A \in \kappa$ form a g.d.t.s. $(S, \mathcal{S}, \tau^g, \kappa^g, V, \mathcal{V})$. In this case, (τ^g, κ^g) is called a *graded ditopology* on (S, \mathcal{S}) corresponding to ditopology (τ, κ) . Thus g.d.t.s. are more general than d.t.s.

Definition 2.7 ([13]). Let $(S, \mathcal{S}, \mathcal{T}, \mathcal{K}, V, \mathcal{V})$ be a g.d.t.s. and $A \in \mathcal{S}$. The families defined by

$$\mathcal{C}(A) = \{P_v \in \mathcal{V} \mid [\mathcal{U} \subseteq \mathcal{T}^v, A \subseteq \bigvee \mathcal{U}] \Rightarrow \exists \mathcal{U}_0 \subseteq \mathcal{U} : A \subseteq \bigvee \mathcal{U}_0\}$$

$$\mathcal{C}^*(A) = \{P_v \in \mathcal{V} \mid [\mathcal{U} \subseteq \mathcal{K}^v, \bigwedge \mathcal{U} \subseteq A] \Rightarrow \exists \mathcal{U}_0 \subseteq \mathcal{U} : \bigwedge \mathcal{U}_0 \subseteq A\}$$

where \mathcal{U}_0 denotes a finite subfamily of \mathcal{U} , are called *compactness* and *co-compactness spectrums* of $A \in \mathcal{S}$ respectively. In particular, the compactness spectrum and the co-compactness spectrum of $(S, \mathcal{S}, \mathcal{T}, \mathcal{K}, V, \mathcal{V})$ are $\mathcal{C}(S)$ and $\mathcal{C}^*(\emptyset)$ respectively.

Definition 2.8 ([20]). Let $(S, \mathcal{S}, \mathcal{T}, \mathcal{K}, V, \mathcal{V})$ be a graded ditopological texture space and the mappings $p\mathcal{T}, p\mathcal{K}, s\mathcal{T}, s\mathcal{K} : \mathcal{S} \rightarrow \mathcal{V}$ be defined by

$$(2.1) \quad p\mathcal{T}(A) = \bigvee \{P_v \mid A \subseteq]A]^v[v^v\}, \quad p\mathcal{K}(A) = \bigvee \{P_v \mid [A]^v[v^v \subseteq A\}$$

$$(2.2) \quad s\mathcal{T}(A) = \bigvee \{P_v \mid A \subseteq [A]^v[v^v\}, \quad s\mathcal{K}(A) = \bigvee \{P_v \mid [A]^v[v^v \subseteq A\}$$

for all $A \in \mathcal{S}$. Then $p\mathcal{T}$ ($p\mathcal{K}$) is called pre-openness (pre-closedness) function; $s\mathcal{T}$ ($s\mathcal{K}$) is called *semi-openness* (*semi-closedness*) function of $(S, \mathcal{S}, \mathcal{T}, \mathcal{K}, V, \mathcal{V})$ and $p\mathcal{T}(A)$ ($p\mathcal{K}(A)$) is called pre-openness (pre-closedness) grade; $s\mathcal{T}(A)$ ($s\mathcal{K}(A)$) is called *semi-openness* (*semi-closedness*) grade of A respectively.

3. MAIN RESULTS

Definition 3.1. Let $(S, \mathcal{S}, \mathcal{T}, \mathcal{K}, V, \mathcal{V})$ be a graded ditopological texture space and the mappings $\beta\mathcal{T}, \beta\mathcal{K} : \mathcal{S} \rightarrow \mathcal{V}$ be defined by

$$(3.1) \quad \beta\mathcal{T}(A) = \bigvee \{P_v \mid A \subseteq [A]^v[v^v\}, \quad \beta\mathcal{K}(A) = \bigvee \{P_v \mid [A]^v[v^v \subseteq A\}$$

for all $A \in \mathcal{S}$. Then $\beta\mathcal{T}$ ($\beta\mathcal{K}$) is called β -openness (β -closedness) function of $(S, \mathcal{S}, \mathcal{T}, \mathcal{K}, V, \mathcal{V})$ and $\beta\mathcal{T}(A)$ ($\beta\mathcal{K}(A)$) is called β -openness (β -closedness) grade of A respectively.

Proposition 3.2. For a graded ditopological texture space $(S, \mathcal{S}, \mathcal{T}, \mathcal{K}, V, \mathcal{V})$ following statements are hold:

- (1) (a) $\mathcal{T} \subseteq p\mathcal{T} \subseteq \beta\mathcal{T}$ and $\mathcal{K} \subseteq p\mathcal{K} \subseteq \beta\mathcal{K}$,
 (b) $\mathcal{T} \subseteq s\mathcal{T} \subseteq \beta\mathcal{T}$ and $\mathcal{K} \subseteq s\mathcal{K} \subseteq \beta\mathcal{K}$,
 (2) The functions $\beta\mathcal{T}$ and $\beta\mathcal{K}$ satisfy the property (GT_3) and (GCT_3) respectively where the texture (V, \mathcal{V}) is discrete.

Proof. (1) (a) For any $A \in \mathcal{S}$ and $v \in V$, we have

$$P_v \subseteq \mathcal{T}(A) \Rightarrow]A]^v = A \Rightarrow A =]A]^v \subseteq [A]^v[v^v \Rightarrow P_v \subseteq p\mathcal{T}(A)$$

and

$$A \subseteq [A]^v[v^v \Rightarrow A \subseteq [A]^v[v^v \subseteq [A]^v[v^v.$$

Then this implies

$$p\mathcal{T}(A) = \bigvee \{P_v \mid A \subseteq [A]^v[v^v\} \subseteq \bigvee \{P_v \mid A \subseteq [A]^v[v^v \subseteq [A]^v[v^v\} = \beta\mathcal{T}(A).$$

Thus we get $\mathcal{T} \subseteq p\mathcal{T} \subseteq \beta\mathcal{T}$. Similarly, it can be shown that $\mathcal{K} \subseteq p\mathcal{K} \subseteq \beta\mathcal{K}$.

(b) For any $A \in \mathcal{S}$ and $v \in V$, we have

$$P_v \subseteq \mathcal{K}(A) \Rightarrow [A]^v = A \Rightarrow [A]^v[v] = A[v] \subseteq A \Rightarrow P_v \subseteq s\mathcal{K}(A)$$

and

$$][A]^v[v] \subseteq A \Rightarrow][A]^v[v] \subseteq][A]^v[v] \subseteq A.$$

Then this implies

$$s\mathcal{K}(A) = \bigvee \{P_v \mid][A]^v[v] \subseteq A\} \subseteq \bigvee \{P_v \mid][A]^v[v] \subseteq A\} = \beta\mathcal{K}(A).$$

Thus we get $\mathcal{K} \subseteq s\mathcal{K} \subseteq \beta\mathcal{K}$. Similarly, it can be shown that $\mathcal{T} \subseteq s\mathcal{T} \subseteq \beta\mathcal{T}$.

(2) Let $A_j \in \mathcal{S}$, $j \in J$ where J is an index set.

(a) Assume that $\bigcap_{j \in J} \beta\mathcal{T}(A_j) \not\subseteq \beta\mathcal{T}(\bigvee_{j \in J} A_j)$. Then $\bigcap_{j \in J} \beta\mathcal{T}(A_j) \not\subseteq Q_v$ and $P_v \not\subseteq \beta\mathcal{T}(\bigvee_{j \in J} A_j)$ for an element $v \in V$. Thus we get $P_v \subseteq \bigcap_{j \in J} \beta\mathcal{T}(A_j)$ by $\bigcap_{j \in J} \beta\mathcal{T}(A_j) \not\subseteq Q_v$. So considering (V, \mathcal{V}) is discrete, we have

$$\begin{aligned} \forall j \in J \ P_v = \{v\} \subseteq \beta\mathcal{T}(A_j) &\Rightarrow \forall j \in J \ A_j \subseteq [][A_j]^v[v] \subseteq [] [\bigvee_{j \in J} A_j]^v [v] \\ &\Rightarrow \bigvee_{j \in J} A_j \subseteq [] [\bigvee_{j \in J} A_j]^v [v] \Rightarrow P_v \subseteq \beta\mathcal{T}(\bigvee_{j \in J} A_j) \end{aligned}$$

and this contradicts with $P_v \not\subseteq \beta\mathcal{T}(\bigvee_{j \in J} A_j)$. Hence $\bigcap_{j \in J} \beta\mathcal{T}(A_j) \subseteq \beta\mathcal{T}(\bigvee_{j \in J} A_j)$.

(b) Assume that $\bigcap_{j \in J} \beta\mathcal{K}(A_j) \not\subseteq \beta\mathcal{K}(\bigcap_{j \in J} A_j)$. Then $\bigcap_{j \in J} \beta\mathcal{K}(A_j) \not\subseteq Q_v$ and $P_v \not\subseteq \beta\mathcal{K}(\bigcap_{j \in J} A_j)$ for an element $v \in V$. Thus we get $P_v \subseteq \bigcap_{j \in J} \beta\mathcal{K}(A_j)$ by $\bigcap_{j \in J} \beta\mathcal{K}(A_j) \not\subseteq Q_v$. So considering (V, \mathcal{V}) is discrete, we have

$$\begin{aligned} \forall j \in J \ P_v = \{v\} \subseteq \beta\mathcal{K}(A_j) &\Rightarrow \forall j \in J \ [] \bigcap_{j \in J} A_j [v] [v] \subseteq [] A_j [v] [v] \subseteq A_j \\ &\Rightarrow [] \bigcap_{j \in J} A_j [v] [v] \subseteq \bigcap_{j \in J} A_j \Rightarrow P_v \subseteq \beta\mathcal{K}(\bigcap_{j \in J} A_j) \end{aligned}$$

and this contradicts with $P_v \not\subseteq \beta\mathcal{K}(\bigcap_{j \in J} A_j)$. Hence $\bigcap_{j \in J} \beta\mathcal{K}(A_j) \subseteq \beta\mathcal{K}(\bigcap_{j \in J} A_j)$. \square

Example 3.3. Let $(S, \mathcal{S}, \tau, \kappa)$ be a ditopological texture space. Then a set $D \in \mathcal{S}$ is β -open (not β -open) in $(S, \mathcal{S}, \tau, \kappa)$ if and only if $\beta\tau^g(D) = 1$ ($\beta\tau^g(D) = 0$) in the graded ditopological texture space $(S, \mathcal{S}, \tau^g, \kappa^g, 1, \mathcal{P}(1))$ respectively. Similarly, a set $D \in \mathcal{S}$ is β -closed (not β -closed) in $(S, \mathcal{S}, \tau, \kappa)$ if and only if $\beta\kappa^g(D) = 1$ ($\beta\kappa^g(D) = 0$) in $(S, \mathcal{S}, \tau^g, \kappa^g, 1, \mathcal{P}(1))$ respectively.

Proposition 3.4. If $(S, \mathcal{S}, \mathcal{T}, \mathcal{K}, \sigma, V, \mathcal{V})$ is a complemented g.d.t.s., then $\beta\mathcal{K} \circ \sigma = \beta\mathcal{T}$ and $\beta\mathcal{T} \circ \sigma = \beta\mathcal{K}$.

Proof. Since $(S, \mathcal{S}, \mathcal{T}, \mathcal{K}, \sigma, V, \mathcal{V})$ is complemented, we get $\sigma([A]^v) =]\sigma(A)[^v$ and $\sigma([A]^v) = [\sigma(A)]^v$ for all $A \in \mathcal{S}$ and $v \in V$ by [20]. Then we have

$$\begin{aligned} (\beta\mathcal{K} \circ \sigma)(A) &= \beta\mathcal{K}(\sigma(A)) = \bigvee \{P_v \mid [] \sigma(A)[^v]^v [^v \subseteq \sigma(A)\} \\ &= \bigvee \{P_v \mid A \subseteq \sigma([] \sigma(A)[^v]^v [^v)\} = \bigvee \{P_v \mid A \subseteq [\sigma([] \sigma(A)[^v]^v)]^v\} \\ &= \bigvee \{P_v \mid A \subseteq [] \sigma([\sigma(A)]^v)[^v]^v\} = \bigvee \{P_v \mid A \subseteq [] [\sigma(\sigma(A))]^v [^v]^v\} \\ &= \bigvee \{P_v \mid A \subseteq [] [A]^v [^v]^v\} = \beta\mathcal{T}(A) \end{aligned}$$

and

$$\begin{aligned} (\beta\mathcal{T} \circ \sigma)(A) &= \beta\mathcal{T}(\sigma(A)) = \bigvee \{P_v \mid \sigma(A) \subseteq [] [\sigma(A)]^v [^v]^v\} \\ &= \bigvee \{P_v \mid \sigma([] [\sigma(A)]^v [^v]^v) \subseteq A\} = \bigvee \{P_v \mid]\sigma([] [\sigma(A)]^v [^v]^v) \subseteq A\} \\ &= \bigvee \{P_v \mid [] [\sigma([\sigma(A)]^v)]^v [^v \subseteq A\} = \bigvee \{P_v \mid [] [\sigma(\sigma(A))]^v [^v]^v \subseteq A\} \\ &= \bigvee \{P_v \mid [] [A]^v [^v]^v \subseteq A\} = \beta\mathcal{K}(A) \end{aligned}$$

for all $A \in \mathcal{S}$. \square

Definition 3.5. Let $(S_k, \mathcal{S}_k, \mathcal{T}_k, \mathcal{K}_k, V_k, \mathcal{V}_k)$, $k = 1, 2$ be g.d.t.s. and $(f, F) : (S_1, \mathcal{S}_1) \rightarrow (S_2, \mathcal{S}_2)$, $(h, H) : (V_1, \mathcal{V}_1) \rightarrow (V_2, \mathcal{V}_2)$ be difunctions. (f, F) is said to be:

- (i) β -continuous w.r.t. (h, H) , if $H^\leftarrow \mathcal{T}_2(A) \subseteq \beta\mathcal{T}_1(F^\leftarrow A)$ for all $A \in \mathcal{S}_2$,
- (ii) $M\beta$ -continuous w.r.t. (h, H) , if $H^\leftarrow \beta\mathcal{T}_2(A) \subseteq \beta\mathcal{T}_1(F^\leftarrow A)$ for all $A \in \mathcal{S}_2$,
- (iii) β -cocontinuous w.r.t. (h, H) , if $h^\leftarrow \mathcal{K}_2(A) \subseteq \beta\mathcal{K}_1(f^\leftarrow A)$ for all $A \in \mathcal{S}_2$,
- (iv) $M\beta$ -cocontinuous w.r.t. (h, H) , if $h^\leftarrow \beta\mathcal{K}_2(A) \subseteq \beta\mathcal{K}_1(f^\leftarrow A)$ for all $A \in \mathcal{S}_2$,
- (v) β -bicontinuous w.r.t. (h, H) , if it is both β -continuous and β -cocontinuous w.r.t. (h, H) ,
- (vi) $M\beta$ -bicontinuous w.r.t. (h, H) , if it is both $M\beta$ -continuous and $M\beta$ -cocontinuous w.r.t. (h, H) .

We say that $((f, F), (h, H))$ is a *relatively $(M)\beta$ -bicontinuous difunction pair*, if (f, F) is $(M)\beta$ -bicontinuous w.r.t. (h, H) .

Example 3.6. Let $(S_k, \mathcal{S}_k, \tau_k, \kappa_k)$, $k = 1, 2$ be d.t.s. and $(f, F) : (S_1, \mathcal{S}_1) \rightarrow (S_2, \mathcal{S}_2)$ be a difunction. Consider graded ditopological texture spaces $(S_k, \mathcal{S}_k, \tau_k^g, \kappa_k^g, V, \mathcal{V})$, $k = 1, 2$ corresponding to ditopological texture spaces $(S_k, \mathcal{S}_k, \tau_k, \kappa_k)$, $k = 1, 2$.

(1) If (f, F) is $M\beta$ -continuous, then we have $I_V^\leftarrow \beta\tau_2^g(A) = \beta\tau_2^g(A) \subseteq \beta\tau_1^g(F^\leftarrow A)$ for each $A \in \mathcal{S}_2$ and so (f, F) is $M\beta$ -continuous w.r.t. the identity difunction (i_V, I_V) . If (f, F) is $M\beta$ -cocontinuous, then we have $i_V^\leftarrow \beta\kappa_2^g(A) = \beta\kappa_2^g(A) \subseteq \beta\kappa_1^g(f^\leftarrow A)$ for each $A \in \mathcal{S}_2$ and so (f, F) is $M\beta$ -cocontinuous w.r.t. the identity difunction (i_V, I_V) .

(2) If (f, F) is β -continuous, then we have $I_V^\leftarrow \tau_2^g(A) = \tau_2^g(A) \subseteq \beta\tau_1^g(F^\leftarrow A)$ for each $A \in \mathcal{S}_2$ and so (f, F) is β -continuous w.r.t. the identity difunction (i_V, I_V) . If (f, F) is β -cocontinuous, then we have $i_V^\leftarrow \kappa_2^g(A) = \kappa_2^g(A) \subseteq \beta\kappa_1^g(f^\leftarrow A)$ for each $A \in \mathcal{S}_2$ and so (f, F) is β -cocontinuous w.r.t. the identity difunction (i_V, I_V) .

In this sense, every β -bicontinuous $(M\beta$ -bicontinuous) difunction between two d.t.s. is considered as β -bicontinuous $(M\beta$ -bicontinuous) difunction w.r.t. identity difunction on the discrete texture on a singleton.

(3) Clearly, each relatively bicontinuous difunction pair is relatively β -bicontinuous by Proposition 3.2 (1).

Definition 3.7. Let $(S, \mathcal{S}, \mathcal{T}, \mathcal{K}, V, \mathcal{V})$ be a g.d.t.s. and $A \in \mathcal{S}$. Then we say that the families

$$\beta\mathcal{C}(A) = \{P_v \mid [\mathcal{U} \subseteq \beta\mathcal{T}^v, A \subseteq \bigvee \mathcal{U}] \Rightarrow \exists \mathcal{U}_0 \subseteq \mathcal{U} : A \subseteq \bigvee \mathcal{U}_0\}$$

$$\beta\mathcal{C}^*(A) = \{P_v \mid [\mathcal{U} \subseteq \beta\mathcal{K}^v, \bigwedge \mathcal{U} \subseteq A] \Rightarrow \exists \mathcal{U}_0 \subseteq \mathcal{U} : \bigwedge \mathcal{U}_0 \subseteq A\}$$

where \mathcal{U}_0 denotes a finite subfamily of \mathcal{U} , are β -compactness and β -cocompactness spectrums of $A \in \mathcal{S}$ respectively. We also say that $\beta\mathcal{C}(S)$ and $\beta\mathcal{C}^*(\emptyset)$ are β -compactness spectrum and β -cocompactness spectrum of the g.d.t.s. $(S, \mathcal{S}, \mathcal{T}, \mathcal{K}, V, \mathcal{V})$ respectively.

Corollary 3.8. Let $(S, \mathcal{S}, \mathcal{T}, \mathcal{K}, V, \mathcal{V})$ be a g.d.t.s. and $A \in \mathcal{S}$. Then $\beta\mathcal{C}(A) \subseteq \mathcal{C}(A)$ and $\beta\mathcal{C}^*(A) \subseteq \mathcal{C}^*(A)$.

Proof. It is clear by Proposition 3.2. \square

Proposition 3.9. If $(S, \mathcal{S}, \mathcal{T}, \mathcal{K}, \sigma, V, \mathcal{V})$ is a complemented g.d.t.s. and $A \in \mathcal{S}$ then $\beta\mathcal{C}(A) = \beta\mathcal{C}^*(\sigma(A))$. Namely, β -compactness and β -cocompactness spectrums of a complemented g.d.t.s. are equal.

Proof. Let $P_v \in \beta\mathcal{C}^*(\sigma(A))$, $\mathcal{U} \subseteq \beta\mathcal{T}^v$ and $A \subseteq \bigvee \mathcal{U}$. Since $(S, \mathcal{S}, \mathcal{T}, \mathcal{K}, \sigma, V, \mathcal{V})$ is complemented, we have $P_v \subseteq \beta\mathcal{T}(U) = (\beta\mathcal{K} \circ \sigma)(U) = \beta\mathcal{K}(\sigma(U))$ for all $U \in \mathcal{U}$ by Proposition 3.4. This implies $\sigma(\mathcal{U}) = \{\sigma(U) \mid U \in \mathcal{U}\} \subseteq \beta\mathcal{K}^v$. On the other hand, we have also $A \subseteq \bigvee \mathcal{U} \Rightarrow \sigma(\bigvee \mathcal{U}) \subseteq \sigma(A) \Rightarrow \bigwedge \sigma(\mathcal{U}) \subseteq \sigma(A)$. Then we get $\bigwedge \sigma(\mathcal{U}_0) \subseteq \sigma(A)$ for a finite subfamily $\mathcal{U}_0 \subseteq \mathcal{U}$ by $P_v \in \beta\mathcal{C}^*(\sigma(A))$. Also $\bigwedge \sigma(\mathcal{U}_0) \subseteq \sigma(A)$ implies $\sigma(\sigma(A)) \subseteq \sigma(\bigwedge \sigma(\mathcal{U}_0))$ and thus $A \subseteq \bigvee \mathcal{U}_0$. So we obtain that $P_v \in \beta\mathcal{C}(A)$, i.e. $\beta\mathcal{C}^*(\sigma(A)) \subseteq \beta\mathcal{C}(A)$. The other inclusion $\beta\mathcal{C}(A) \subseteq \beta\mathcal{C}^*(\sigma(A))$ can be shown similarly.

Since $\sigma(S) = \emptyset$, we get $\beta\mathcal{C}(S) = \beta\mathcal{C}^*(\sigma(S)) = \beta\mathcal{C}^*(\emptyset)$. Then β -compactness and β -cocompactness spectrums of $(S, \mathcal{S}, \mathcal{T}, \mathcal{K}, \sigma, V, \mathcal{V})$ are equal. \square

Theorem 3.10. Let $(S_k, \mathcal{S}_k, \mathcal{T}_k, \mathcal{K}_k, V_k, \mathcal{V}_k)$, $k = 1, 2$ be g.d.t.s. and $(f, F) : (S_1, \mathcal{S}_1) \rightarrow (S_2, \mathcal{S}_2)$, $(h, H) : (V_1, \mathcal{V}_1) \rightarrow (V_2, \mathcal{V}_2)$ be difunctions.

(1) If (f, F) is $M\beta$ -continuous w.r.t. (h, H) and $A \in \mathcal{S}_1$, then

$$P_v \in \beta\mathcal{C}_1(A) \Rightarrow P_t \in \beta\mathcal{C}_2(f^\rightarrow A),$$

(2) If (f, F) is $M\beta$ -cocontinuous w.r.t. (h, H) and $A \in \mathcal{S}_1$, then

$$P_v \in \beta\mathcal{C}_1^*(A) \Rightarrow P_t \in \beta\mathcal{C}_2^*(F^\rightarrow A)$$

where $P_v \in \mathcal{V}_1$, $P_t \in \mathcal{V}_2$ with $P_v \subseteq h^\leftarrow P_t$.

Proof. We omit the proof of (1) and prove (2) since the proofs of (1) and (2) are similar. Let $P_v \in \beta\mathcal{C}_1^*(A)$, $P_v \in \mathcal{V}_1$, $P_t \in \mathcal{V}_2$ and $P_v \subseteq h^\leftarrow P_t$. Considering the definition of $\beta\mathcal{C}_2^*(F^\rightarrow A)$, if we suppose that $(\mathcal{U} \subseteq \beta\mathcal{K}_2^t, \bigwedge \mathcal{U} \subseteq F^\rightarrow A)$ then we have

$$\bigwedge_{U \in \mathcal{U}} f^\leftarrow U = \bigwedge f^\leftarrow \mathcal{U} = f^\leftarrow (\bigwedge \mathcal{U}) \subseteq f^\leftarrow (F^\rightarrow A) \subseteq A$$

by Proposition 2.2 and Proposition 2.4 (3). Besides, since $\mathcal{U} \subseteq \beta\mathcal{K}_2^t \Rightarrow (\forall U \in \mathcal{U} P_t \subseteq \beta\mathcal{K}_2(U)) \Rightarrow P_t \subseteq \beta\mathcal{K}_2(\mathcal{U})$ and (f, F) is $M\beta$ -cocontinuous w.r.t. (h, H) we get

$$P_v \subseteq h^\leftarrow P_t \subseteq h^\leftarrow(\beta\mathcal{K}_2(\mathcal{U})) \subseteq \beta\mathcal{K}_1(f^\leftarrow \mathcal{U}).$$

Since $P_v \in \beta\mathcal{C}_1^*(A)$, we have $\bigwedge_{U \in \mathcal{U}_0} f^\leftarrow U \subseteq A$ for a finite subfamily $\mathcal{U}_0 \subseteq \mathcal{U}$. Then we get

$$F^\rightarrow A \supseteq F^\rightarrow \left(\bigwedge_{U \in \mathcal{U}_0} f^\leftarrow U \right) = \bigwedge_{U \in \mathcal{U}_0} F^\rightarrow(f^\leftarrow U) \supseteq \bigwedge_{U \in \mathcal{U}_0} U = \bigwedge \mathcal{U}_0$$

by Proposition 2.2 and Proposition 2.4 (3). Thus we obtain that $P_t \in \beta\mathcal{C}_2^*(F^\rightarrow A)$. \square

Corollary 3.11. *In addition to the conditions in the above theorem, if (f, F) is surjective, then the following results hold:*

- (1) *if (f, F) is $M\beta$ -continuous w.r.t. (h, H) , then*

$$P_v \in \beta\mathcal{C}_1(S_1) \Rightarrow P_t \in \beta\mathcal{C}_2(S_2),$$

- (2) *if (f, F) is $M\beta$ -cocontinuous w.r.t. (h, H) , then*

$$P_v \in \beta\mathcal{C}_1^*(\emptyset) \Rightarrow P_t \in \beta\mathcal{C}_2^*(\emptyset)$$

where $P_v \in \mathcal{V}_1, P_t \in \mathcal{V}_2$ with $P_v \subseteq h^\leftarrow P_t$.

Proof. We get $f^\leftarrow \emptyset = F^\leftarrow \emptyset = \emptyset$ and $f^\leftarrow S_2 = F^\leftarrow S_2 = S_1$ by using Proposition 2.4 (2). Besides, surjectivity of (f, F) implies $F^\rightarrow(f^\leftarrow S_2) = S_2 = f^\rightarrow(F^\leftarrow S_2)$ and $F^\rightarrow(f^\leftarrow \emptyset) = \emptyset = f^\rightarrow(F^\leftarrow \emptyset)$ by using Proposition 2.4 (4). Then we have $f^\rightarrow S_1 = S_2$ and $F^\rightarrow \emptyset = \emptyset$. Thus the result is obtained by Theorem 3.10. \square

Definition 3.12. Let $(S, \mathcal{S}, \mathcal{T}, \mathcal{K}, V, \mathcal{V})$ be a g.d.t.s. Then the families

$$\beta\Omega = \{P_v \mid [A \in \mathcal{S}, A \neq S] \Rightarrow [P_v \subseteq \beta\mathcal{K}(A) \Rightarrow P_v \in \beta\mathcal{C}(A)]\}$$

$$\beta\Omega^* = \{P_v \mid [A \in \mathcal{S}, A \neq \emptyset] \Rightarrow [P_v \subseteq \beta\mathcal{T}(A) \Rightarrow P_v \in \beta\mathcal{C}^*(A)]\}$$

are called β -stableness spectrum and β -costableness spectrum of the g.d.t.s. $(S, \mathcal{S}, \mathcal{T}, \mathcal{K}, V, \mathcal{V})$ respectively.

Proposition 3.13. *If $(S, \mathcal{S}, \mathcal{T}, \mathcal{K}, \sigma, V, \mathcal{V})$ is a complemented g.d.t.s. then the β -stableness spectrum and the β -costableness spectrum are equal, i.e., $\beta\Omega = \beta\Omega^*$.*

Proof. Let $P_v \in \beta\Omega, A \in \mathcal{S}, A \neq \emptyset$ and $P_v \subseteq \beta\mathcal{T}(A)$. Since $(S, \mathcal{S}, \mathcal{T}, \mathcal{K}, \sigma, V, \mathcal{V})$ is complemented, we have $P_v \subseteq \beta\mathcal{T}(A) = (\beta\mathcal{K} \circ \sigma)(A) = \beta\mathcal{K}(\sigma(A))$. Moreover, $A \in \mathcal{S}, A \neq \emptyset$ implies $\sigma(A) \in \mathcal{S}, \sigma(A) \neq S$. Then using $P_v \in \beta\Omega$ and Proposition 3.9, we get $P_v \in \beta\mathcal{C}(\sigma(A)) = \beta\mathcal{C}^*(A)$. Thus we obtain that $\beta\Omega \subseteq \beta\Omega^*$. Similarly, it can be shown that $\beta\Omega^* \subseteq \beta\Omega$. \square

Theorem 3.14. *Let $(S_k, \mathcal{S}_k, \mathcal{T}_k, \mathcal{K}_k, V_k, \mathcal{V}_k), k = 1, 2$ be g.d.t.s. and $(f, F) : (S_1, \mathcal{S}_1) \rightarrow (S_2, \mathcal{S}_2), (h, H) : (V_1, \mathcal{V}_1) \rightarrow (V_2, \mathcal{V}_2)$ be difunctions. If (f, F) is surjective and $M\beta$ -bicontinuous w.r.t. (h, H) , then*

$$(1) P_v \in \beta\Omega_1 \Rightarrow P_t \in \beta\Omega_2,$$

$$(2) P_v \in \beta\Omega_1^* \Rightarrow P_t \in \beta\Omega_2^*$$

where $P_v \in \mathcal{V}_1, P_t \in \mathcal{V}_2$ with $P_v \subseteq h^\leftarrow P_t$.

Proof. (1) Let $P_v \in \beta\Omega_1$. Let $P_t \subseteq \beta\mathcal{K}_2(B)$ is given where $B \in \mathcal{S}_2$, $B \neq S_2$. Then we get $P_v \subseteq h^\leftarrow P_t \subseteq h^\leftarrow(\beta\mathcal{K}_2(B)) \subseteq \beta\mathcal{K}_1(f^\leftarrow B)$ by using that $P_v \subseteq h^\leftarrow P_t$ and (f, F) is $M\beta$ -bicontinuous w.r.t. (h, H) . Thus we have $P_v \subseteq \beta\mathcal{K}_1(f^\leftarrow B)$. On the other hand, since (f, F) is surjective and $B \neq S_2$, we get $f^\leftarrow B \neq S_1$. So we obtain that $P_v \in \beta\mathcal{C}_1(f^\leftarrow B)$ by using $P_v \in \beta\Omega_1$. Hence we have $P_t \in \beta\mathcal{C}_2(f^\rightarrow(f^\leftarrow B)) = \beta\mathcal{C}_2(B)$ by Theorem 3.10 and Proposition 2.4 (1) and (4). Therefore we get $P_t \in \beta\Omega_2$.

(2) Let $P_v \in \beta\Omega_1^*$. Let $P_t \subseteq \beta\mathcal{T}_2(B)$ is given where $B \in \mathcal{S}_2$, $B \neq \emptyset$. Then we get $P_v \subseteq h^\leftarrow P_t \subseteq h^\leftarrow(\beta\mathcal{T}_2(B)) = H^\leftarrow(\beta\mathcal{T}_2(B)) \subseteq \beta\mathcal{T}_1(F^\leftarrow B)$ by using that $P_v \subseteq h^\leftarrow P_t$ and (f, F) is $M\beta$ -bicontinuous w.r.t. (h, H) . Thus we have $P_v \subseteq \beta\mathcal{T}_1(F^\leftarrow B)$. On the other hand, since (f, F) is surjective and $B \neq \emptyset$, we get $F^\leftarrow B \neq \emptyset$. So we obtain that $P_v \in \beta\mathcal{C}_1^*(F^\leftarrow B)$ by using $P_v \in \beta\Omega_1^*$. Hence we have $P_t \in \beta\mathcal{C}_2^*(F^\rightarrow(F^\leftarrow B)) = \beta\mathcal{C}_2^*(B)$ by Theorem 3.10 and Proposition 2.4 (1) and (4). Therefore we get $P_t \in \beta\Omega_2^*$. \square

Definition 3.15. Let $(S, \mathcal{S}, \mathcal{T}, \mathcal{K}, V, \mathcal{V})$ be a g.d.t.s. Then the family defined by

$$\beta\mathcal{DC} = \beta\mathcal{C}(S) \cap \beta\mathcal{C}^*(\emptyset) \cap \beta\Omega \cap \beta\Omega^*$$

is called β -dcompactness spectrum of the g.d.t.s. $(S, \mathcal{S}, \mathcal{T}, \mathcal{K}, V, \mathcal{V})$.

Corollary 3.16. Let $(S_k, \mathcal{S}_k, \mathcal{T}_k, \mathcal{K}_k, V_k, \mathcal{V}_k)$, $k = 1, 2$ be g.d.t.s. and $(f, F) : (S_1, \mathcal{S}_1) \rightarrow (S_2, \mathcal{S}_2)$, $(h, H) : (V_1, \mathcal{V}_1) \rightarrow (V_2, \mathcal{V}_2)$ be difunctions. If (f, F) is surjective and $M\beta$ -bicontinuous w.r.t. (h, H) , then

$$P_v \in \beta\mathcal{DC}_1 \Rightarrow P_t \in \beta\mathcal{DC}_2$$

where $P_v \in \mathcal{V}_1$, $P_t \in \mathcal{V}_2$ with $P_v \subseteq h^\leftarrow P_t$.

Proof. It is an immediate result of Corollary 3.11 and Theorem 3.14. \square

Example 3.17. Let $(S, \mathcal{S}, \tau, \kappa)$ be a d.t.s. If $(S, \mathcal{S}, \tau, \kappa)$ is β -compact (β -cocompact, β -dcompact), then for the g.d.t.s. $(S, \mathcal{S}, \tau^g, \kappa^g, 1, \mathcal{P}(1))$, $P_v \in \beta\mathcal{C}(S)$ ($P_v \in \beta\mathcal{C}^*(\emptyset)$, $P_v \in \beta\mathcal{DC}$) respectively for all $v \in 1 = \{0\}$, i.e. $v = 0$.

Following proposition shows β -compactness relationships between ditopological texture spaces and graded ditopological texture spaces:

Proposition 3.18. Let $(S, \mathcal{S}, \mathcal{T}, \mathcal{K}, V, \mathcal{V})$ be a g.d.t.s. Then the following hold for each $v \in V$:

- (1) $P_v \in \beta\mathcal{C}(S) \Leftrightarrow$ The d.t.s. $(S, \mathcal{S}, \mathcal{T}^v, \mathcal{K}^v)$ is β -compact,
- (2) $P_v \in \beta\mathcal{C}^*(\emptyset) \Leftrightarrow$ The d.t.s. $(S, \mathcal{S}, \mathcal{T}^v, \mathcal{K}^v)$ is β -cocompact,
- (3) $P_v \in \beta\Omega \Leftrightarrow$ The d.t.s. $(S, \mathcal{S}, \mathcal{T}^v, \mathcal{K}^v)$ is β -stable,
- (4) $P_v \in \beta\Omega^* \Leftrightarrow$ The d.t.s. $(S, \mathcal{S}, \mathcal{T}^v, \mathcal{K}^v)$ is β -costable,
- (5) $P_v \in \beta\mathcal{DC} \Leftrightarrow$ The d.t.s. $(S, \mathcal{S}, \mathcal{T}^v, \mathcal{K}^v)$ is β -dcompact.

Proof. It is straightforward. \square

Example 3.19. Take discrete textures $(S, \mathcal{S} = \mathcal{P}(S))$ and $(V, \mathcal{V} = \mathcal{P}(V))$ where $S \neq \emptyset$ and $V = \{x, y, z, r\}$. If we define the mappings $\mathcal{T}, \mathcal{K} : \mathcal{S} \rightarrow \mathcal{V}$ as

$$\mathcal{T}(A) = \begin{cases} V, & A = \emptyset \text{ or } A = S \\ P_z = \{z\}, & \text{otherwise} \end{cases}$$

$$\mathcal{K}(A) = \begin{cases} V, & A = \emptyset \text{ or } A = S \\ P_r = \{r\}, & \text{otherwise} \end{cases}$$

for all $A \in \mathcal{S}$, then we get a g.d.t.s. $(S, \mathcal{S}, \mathcal{T}, \mathcal{K}, V, \mathcal{V})$. Thus we obtain $\mathcal{T}^z = \mathcal{S} = \mathcal{P}(S)$, $\mathcal{T}^x = \mathcal{T}^y = \mathcal{T}^r = \{S, \emptyset\}$, $\mathcal{K}^r = \mathcal{S} = \mathcal{P}(S)$, $\mathcal{K}^x = \mathcal{K}^y = \mathcal{K}^z = \{S, \emptyset\}$. So we have $]A[^z = A$ and $]A[^r = A$ for all $A \in \mathcal{S}$. Also $A \neq S$, $A \in \mathcal{S}$ implies $]A[^x =]A[^y =]A[^r = \emptyset$ and $A \neq \emptyset$, $A \in \mathcal{S}$ implies $]A[^x =]A[^y =]A[^z = S$. Hence we get $\beta\mathcal{T}^x = \beta\mathcal{T}^y = \beta\mathcal{T}^z = \mathcal{S} = \mathcal{P}(S)$, $\beta\mathcal{T}^r = \{\emptyset, S\}$, $\beta\mathcal{K}^x = \beta\mathcal{K}^y = \beta\mathcal{K}^r = \mathcal{S} = \mathcal{P}(S)$, $\beta\mathcal{K}^z = \{\emptyset, S\}$. Therefore we obtain that

$$\beta\mathcal{C}(S) = \beta\mathcal{C}^*(\emptyset) = \beta\Omega = \beta\Omega^* = \beta\mathcal{DC} = \{\{x\}, \{y\}, \{z\}, \{r\}\}$$

if S is finite.

If a subset $A \in \mathcal{S}$ is infinite, then $A \subseteq \bigvee \mathcal{U} = \bigvee_{s \in A} \{s\}$ where $\mathcal{U} = \{P_s \mid s \in A\}$. Yet there is no finite subfamily \mathcal{U}_0 of \mathcal{U} such that $A \subseteq \bigvee \mathcal{U}_0$. Thus we get that $\beta\mathcal{C}(S) = \beta\Omega = \{P_r\} = \{r\}$ if S is infinite.

On the other hand, for a subset $A \in \mathcal{S}$, if $S \setminus A$ is infinite, then $\bigwedge \mathcal{U} = \bigwedge_{x \in (S \setminus A)} ((S \setminus A) \setminus P_x) = \emptyset \subseteq A$ where $\mathcal{U} = \{(S \setminus A) \setminus P_x \mid x \in (S \setminus A)\}$. Yet there is no finite subfamily \mathcal{U}_0 of \mathcal{U} such that $\bigwedge \mathcal{U}_0 \subseteq A$. So we have $\beta\mathcal{C}^*(\emptyset) = \beta\Omega^* = \{P_z\} = \{z\}$ and $\beta\mathcal{DC} = \emptyset$ if S is infinite.

Moreover, we get

$$p\mathcal{T}(A) = \begin{cases} V, & A = \emptyset \text{ or } A = S \\ \{x, y, z\}, & \text{otherwise} \end{cases}$$

$$p\mathcal{K}(A) = \begin{cases} V, & A = \emptyset \text{ or } A = S \\ \{x, y, r\}, & \text{otherwise} \end{cases}$$

$$s\mathcal{T}(A) = \begin{cases} V, & A = \emptyset \text{ or } A = S \\ P_z = \{z\}, & \text{otherwise} \end{cases}$$

$$s\mathcal{K}(A) = \begin{cases} V, & A = \emptyset \text{ or } A = S \\ P_r = \{r\}, & \text{otherwise} \end{cases}$$

$$\beta\mathcal{T}(A) = \begin{cases} V, & A = \emptyset \text{ or } A = S \\ \{x, y, z\}, & \text{otherwise} \end{cases}$$

$$\beta\mathcal{K}(A) = \begin{cases} V, & A = \emptyset \text{ or } A = S \\ \{x, y, r\}, & \text{otherwise} \end{cases}$$

for all $A \in \mathcal{S}$. Hence we have $\mathcal{T} \subseteq p\mathcal{T} = \beta\mathcal{T}$, $\mathcal{K} \subseteq p\mathcal{K} = \beta\mathcal{K}$, $\mathcal{T} = s\mathcal{T} \subseteq \beta\mathcal{T}$, $\mathcal{K} = s\mathcal{K} \subseteq \beta\mathcal{K}$ and this result is also an example for Proposition 3.2 (1).

4. CONCLUSION

This study focuses on generalizing the concepts of β -openness and β -compactness in d.t.s. to the g.d.t.s. To enhance the theory of g.d.t.s., it is important to examine the properties of these new concepts. The interrelations of openness, semi-openness, pre-openness and β -openness (closedness, semi-closedness, pre-closedness and β -closedness) grade of an element of a texture are investigated in a g.d.t.s.: $\mathcal{T} \subseteq p\mathcal{T} \subseteq \beta\mathcal{T}$, $\mathcal{K} \subseteq p\mathcal{K} \subseteq \beta\mathcal{K}$, $\mathcal{T} \subseteq s\mathcal{T} \subseteq \beta\mathcal{T}$, $\mathcal{K} \subseteq s\mathcal{K} \subseteq \beta\mathcal{K}$ (Proposition 3.2). In a complemented g.d.t.s. with complementation σ ; $\beta\mathcal{K} \circ \sigma = \beta\mathcal{T}$ and $\beta\mathcal{T} \circ \sigma = \beta\mathcal{K}$

(Proposition 3.4). We also show that β -compactness and β -cocompactness spectrums of a complemented g.d.t.s. are equal in Proposition 3.9.

The relationships between the concepts of $M\beta$ -continuity (β -continuity) in d.t.s. and relatively $M\beta$ -continuity (relatively β -continuity) in g.d.t.s. are studied respectively. Besides, the properties of β -(di)compactness spectrum in g.d.t.s., the relationship between β -(di)compactness in d.t.s. and β -(di)compactness spectrum in g.d.t.s. are examined.

Thus, this work develops the theory of graded ditopology by these findings and it allows a more comprehensive approach to the theory.

ACKNOWLEDGEMENTS

The author would like to thank the referees for their helpful suggestions and comments.

REFERENCES

- [1] M. E. Abd El-Monsef, S. N. El-Deep and R. A. Mahmoud, β -open sets and β -continuous mappings, Bull. Fac. Sci. Assiut Univ. 12 (1983) 77–90.
- [2] D. Andrijević, Semi-preopen sets, Mat. Vesnik 38 (1986) 24–32.
- [3] M. Caldas and S. Jafari, A new decomposition of β -open functions, Chaos Solitons & Fractals 40 (2009) 10–12.
- [4] I. M. Hanafy, Fuzzy β -compactness and fuzzy β -closed spaces, Turk. J. Math. 28 (2004) 281–294.
- [5] L. M. Brown, A. P. Šostak, Categories of fuzzy topology in the context of graded ditopologies on textures, Iran. J. Fuzzy Syst. 11(6) (2014) 1–20.
- [6] L. M. Brown, M. Diker, Ditopological texture spaces and intuitionistic sets, Fuzzy Sets and Systems 98 (1998) 217–224.
- [7] L. M. Brown, R. Ertürk, Fuzzy sets as texture spaces, I. Representation theorems, Fuzzy Sets and Systems 110 (2000) 227–236.
- [8] A. P. Šostak, On a fuzzy topological structure, Rend. Circ. Mat. Palermo II 11 (1985) 89–103.
- [9] T. Kubiak, On fuzzy topologies, PhD Thesis, A. Mickiewicz University Poznan, Poland, 1985.
- [10] R. Ekmekçi and R. Ertürk, Convergence in graded ditopological texture spaces, Applied General Topology 17(1) (2016) 17–35.
- [11] R. Ekmekçi and R. Ertürk, Q-Convergence in graded ditopological texture spaces, Mathematica Moravica 21(1) (2017) 27–36.
- [12] Ş. Dost, L. M. Brown and R. Ertürk, β -open and β -closed sets in ditopological texture spaces, Filomat 24(2) (2010) 11–26.
- [13] R. Ekmekçi, A Tychonoff theorem for graded ditopological texture spaces, Commun. Fac. Sci. Univ. Ank. Ser. A1. Math. Stat. 69(1) (2020) 193–212.
- [14] A. P. Šostak, On compactness and connectedness degrees of fuzzy topological spaces, General Topology and its Relations to Modern Analysis and Algebra, Heldermann Verlag, Berlin (1988) 519–532.
- [15] A. P. Šostak, Two decades of fuzzy topology: basic ideas, notions and results, Russian Math. Surveys 44(6) (1989), 125–186.
- [16] L. M. Brown, R. Ertürk, Ş. Dost, Ditopological texture spaces and fuzzy topology, I. Basic concepts, Fuzzy Sets and Systems 147(2) (2004) 171–199.
- [17] L. M. Brown, R. Ertürk, Ş. Dost, Ditopological texture spaces and fuzzy topology, II. Topological considerations, Fuzzy Sets and Systems 147(2) (2004) 201–231.
- [18] Ş. Dost, Semi-open and semi-closed sets in ditopological texture spaces, J. Adv. Math. Stud. 5(1) (2012) 97–110.
- [19] L. M. Brown, M. M. Gohar, Compactness in ditopological texture spaces, Hacet. J. Math. Stat. 38(1) (2009), 21–43.

- [20] R. Ekmekçi, Semi-compactness spectrum in graded ditopological texture spaces, Annals of Oradea Univ. Math. Fasc. 28(2) (2021) 51–60.

RAMAZAN EKMEKÇİ (rekmekci@klu.edu.tr) – Department of Mathematics, Faculty of Art and Science, Kırklareli University, postal code 39100, Kırklareli, Turkey