

Cosine similarity measures of neutrosophic soft set

I. R. SUMATHI, I. AROCKIARANI

Received 8 January 2016; Revised 12 February 2016; Accepted 15 May 2016

ABSTRACT. In this paper we have introduced the concept of cosine similarity measures for neutrosophic soft set and interval valued neutrosophic soft set. An application is given to show its practicality and effectiveness.

2010 AMS Classification: 03B99, 03E99

Keywords: Neutrosophic soft set, Cosine similarity of Neutrosophic soft set, Interval valued Neutrosophic soft set.

Corresponding Author: I. R. Sumathi (sumathi_raman2005@yahoo.co.in)

1. INTRODUCTION

Neutrosophy is a branch of philosophy, which emphasizes the origin and nature of neutralities, along with their interaction with different conceptive domains. Fuzzy logic extends classical logic by assigning a membership function ranging in degree between 0 and 1 to the variables. As a generalization of fuzzy logic, neutrosophic logic introduces a new component called indeterminacy and carries more information than fuzzy logic. The application of neutrosophic logic would lead to better performance than fuzzy logic. Neutrosophic logic is so new that its use in many fields merit exploration. The words "neutrosophy" and "neutrosophic" were introduced by F. Smarandache in his 1998 book [23]. Etymologically, "neutro-sophy" (noun) [French *neutre* < Latin *neuter*, neutral, and Greek *sophia*, skill/wisdom] means knowledge of neutral thought. Neutrosophic set [24],[25] is a mathematical tool for handling problems involving imprecise, indeterminacy and inconsistent data. In neutrosophic logic a proposition has a degree of truth (T), a degree of indeterminacy (I), and a degree of falsity (F), where T, I, F are standard or non-standard subsets of $]-0, 1+[$. A. A. Salama [7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22] studied the various notions of neutrosophic sets.

In several application it is often needed to compare two sets and we are interested to know whether two patterns or images are identical or approximately identical of atleast to what degree they are identical. Several researchers like Hung

[3], Jun Ye [4, 5], P.Majumdar [6], C.Wang[26] and many authors have studied the problem of similarity measures between intuitionistic fuzzy sets, neutrosophic sets and vague soft sets. In this paper we have introduced some new cosine similarity measures for neutrosophic soft sets and derived some of their properties. A decision making method based on this similarity measure is constructed.[7], [8], [9]

2. PRELIMINARIES

Definition 2.1 ([1]). A neutrosophic set A on the universe of discourse X is defined as $A = \langle x, T_A(x), I_A(x), F_A(x) \rangle, x \in X$ where $T, I, F : X \rightarrow [0, 1]$ and $0 \leq T_A(x) + I_A(x) + F_A(x) \leq 3$, where T represents the truth value, I represents the indeterministic value and F represents the false value.

Definition 2.2. [1] Let X be a non empty set, and let $A = \langle x, T_A(x), I_A(x), F_A(x) \rangle$ and $B = \langle x, T_B(x), I_B(x), F_B(x) \rangle$ be neutrosophic sets. Then A is a subset of B , if $\forall x \in X, T_A(x) \leq T_B(x), I_A(x) \leq I_B(x)$ and $F_A(x) \geq F_B(x)$.

Definition 2.3 ([1]). Let X be a non empty set, and let $A = \langle x, T_A(x), I_A(x), F_A(x) \rangle$ and $B = \langle x, T_B(x), I_B(x), F_B(x) \rangle$ be neutrosophic sets. Then

$$A \cup B = \langle x, \max(T_A(x), T_B(x)), \max(I_A(x), I_B(x)), \min(F_A(x), F_B(x)) \rangle,$$

$$A \cap B = \langle x, \min(T_A(x), T_B(x)), \min(I_A(x), I_B(x)), \max(F_A(x), F_B(x)) \rangle.$$

Definition 2.4 ([1]). Let U be a the initial universal set and E be a set of parameters. Consider a non-empty set $A, A \subseteq E$. Let $P(U)$ denote the set of all neutrosophic sets of U . The collection (F, A) is termed to be the neutrosophic soft set over U , where F is a mapping given by $F:A \rightarrow P(U)$. (Neutrosophic soft set is denoted by NSS).

Definition 2.5 ([1]). A neutrosophic soft set (F, A) over the universe U is said to be empty neutrosophic soft set with respect to the parameter A , if $T_{F(e)} = 0, I_{F(e)} = 0, F_{F(e)} = 1, \forall x \in U, \forall e \in A$. It is denoted by $\tilde{0}$.

Definition 2.6 ([1]). A neutrosophic soft set (F, A) over the universe U is said to be universe neutrosophic soft set with respect to the parameter A , if $T_{F(e)} = 1, I_{F(e)} = 1, F_{F(e)} = 0, \forall x \in U, \forall e \in A$. It is denoted by $\tilde{1}$.

Definition 2.7 ([1]). A neutrosophic soft set (F, A) is said to be a subset of neutrosophic soft set (G, B) , if $A \subseteq B$ and $F(e) \subseteq G(e) \forall e \in E, u \in U$. We denote it by $(F, A) \subseteq (G, B)$.

Definition 2.8 ([1]). The complement of neutrosophic soft set (F, A) denoted by $(F, A)^c$ is defined as $(F, A)^c = (F^c, \neg A)$, where $F^c : \neg A \rightarrow P(U)$ is a mapping given by

$$F^c(\alpha) = \langle x, T_{F^c}(x) = F_F(x), I_{F^c}(x) = 1 - I_F(x), F_{F^c}(x) = T_F(x) \rangle.$$

Definition 2.9. [1] The union of two neutrosophic soft sets (F, A) and (G, B) over (U, E) is neutrosophic soft set, where $C = A \cup B, \forall e \in C$

$$H(e) = \begin{cases} F(e) & \text{if } e \in A - B \\ G(e) & \text{if } e \in B - A \\ F(e) \cup G(e) & \text{if } e \in A \cap B \end{cases}$$

and is written as $(F,A)\tilde{\cup}(G,B) = (H,C)$.

Definition 2.10 ([1]). The intersection of two neutrosophic soft sets (F,A) and (G,B) over (U,E) is neutrosophic soft set, where $C = A \cap B, \forall e \in C H(e) = F(e) \cap G(e)$ and is written as $(F,A)\tilde{\cap}(G,B) = (H,C)$.

Definition 2.11 ([2]). An interval valued neutrosophic set (IVNS in short) on a universe X is an object of the form, where $T_A(x) = X \rightarrow \text{Int}([0, 1]), I_A(x) = X \rightarrow \text{Int}([0, 1])$ and $F_A(x) = X \rightarrow \text{Int}([0, 1]), \text{Int}([0,1])$ stands for the set of all closed subinterval of $[0,1]$ satisfies the condition : $\forall x \in X,$

$$\sup T_A(x) + \sup I_A(x) + \sup F_A(x) \leq 3.$$

Definition 2.12 ([2]). For an arbitrary set $A \subseteq [0, 1],$ we define $\underline{A} = \inf A$ and $\overline{A} = \sup A.$

Definition 2.13 ([2]). Let U be an initial universe and E be a set of parameters. IVNS (U) denotes the set of all interval valued neutrosophic sets of $U.$ Let $A \subseteq E.$ A pair (F,A) is an interval valued neutrosophic soft set over $U,$ where F is a mapping given by $F: A \rightarrow \text{IVNS}(U).$

Note: Interval valued neutrosophic soft set/sets is denoted by IVNSS/IVNSSs.

Definition 2.14 ([2]). Let U be an initial universe and E be a set of parameters. Suppose that $A,B \subseteq E, (F,A)$ and (G,B) be two IVNSSs, we say that (F,A) is an interval valued neutrosophic soft subset of $(G,B),$ if

- (i) $A \subseteq B,$
- (ii) $e \in A, F(e)$ is an interval valued neutrosophic soft subset of $G(e),$ that is, for all $x \in U$ and $e \in A,$

$$\underline{T}_{F(e)}(x) \leq \underline{T}_{G(e)}(x), \overline{T}_{F(e)}(x) \leq \overline{T}_{G(e)}(x),$$

$$\underline{I}_{F(e)}(x) \leq \underline{I}_{G(e)}(x), \overline{I}_{F(e)}(x) \leq \overline{I}_{G(e)}(x),$$

$$\underline{F}_{F(e)}(x) \geq \underline{F}_{G(e)}(x), \overline{F}_{F(e)}(x) \geq \overline{F}_{G(e)}(x).$$

And it is denoted by $(F,A) \subseteq (G,B).$

Similarly, (F,A) is said to be an interval valued neutrosophic soft super set of $(G,B),$ if (G,B) is an interval valued neutrosophic soft subset of $(F,A),$ we denote it by $(F,A) \supseteq (G,B).$

Definition 2.15 ([2]). The complement of an IVNSS (F,A) is denoted by $(F,A)^c$ and is defined as $(F,A)^c = (F^c, \neg A)$ where $F^c : \neg A \rightarrow \text{IVNSS}(U)$ is a mapping given by

$$F^c(e) = \langle F_{F(\neg e)}(x), (I_{F(\neg e)}(x))^c, T_{F(\neg e)}(x) \rangle \text{ for all } x \in U \text{ and } e \in \neg A,$$

where $(I_{F(\neg e)}(x))^c = [1 - \underline{I}_{F(e)}(x), 1 - \overline{I}_{F(e)}(x)].$

3. COSINE SIMILARITY MEASURE NEUTROSOPHIC SOFT SET AND INTERVAL VALUED NEUTROSOPHIC SOFT SET

Definition 3.1. Let $U = \{x_1, x_2, \dots, x_n\}$ be the universal set of elements and $E = \{e_1, e_2, \dots, e_m\}$ be the universal set of parameters. Let (F,A) and (G,B) be two neutrosophic soft sets over $U.$ Then we define the cosine similarity measures between (F,A) and (G,B) as follows.

$$CS_1((F,A), (G,B))$$

$$\begin{aligned}
 &= \frac{1}{mn} \sum_{i=1}^m \sum_{j=1}^n \cos \left[\frac{\pi}{2} (|T_{F(e_i)}(x_j) - T_{G(e_i)}(x_j)|) \vee (|I_{F(e_i)}(x_j) - I_{G(e_i)}(x_j)|) \vee \right. \\
 &\qquad \qquad \qquad \left. (|F_{F(e_i)}(x_j) - F_{G(e_i)}(x_j)|) \right], \\
 &\quad CS_2((F, A), (G, B)) \\
 &= \frac{1}{mn} \sum_{i=1}^m \sum_{j=1}^n \cos \left[\frac{\pi}{6} (|T_{F(e_i)}(x_j) - T_{G(e_i)}(x_j)|) + (|I_{F(e_i)}(x_j) - I_{G(e_i)}(x_j)|) + \right. \\
 &\qquad \qquad \qquad \left. (|F_{F(e_i)}(x_j) - F_{G(e_i)}(x_j)|) \right],
 \end{aligned}$$

where the symbol \vee is the maximum operation. The two similarity measures satisfy the axiomatic requirements of similarity measures.

Proposition 3.2. *Let $U = \{x_1, x_2, \dots, x_n\}$ be the universal set of elements and $E = \{e_1, e_2, \dots, e_m\}$ be the universal set of parameters. Then for two neutrosophic soft sets (F, A) and (G, B) over U the cosine similarity measure $CS_k((F, A), (G, B))$ ($k=1, 2$) should satisfy the following properties (1)–(4).*

- (1) $0 \leq CS_k((F, A), (G, B)) \leq 1$.
- (2) $CS_k((F, A), (G, B)) = 1$ if and only if $(F, A) = (G, B)$.
- (3) $CS_k((F, A), (G, B)) = CS_k((G, B), (F, A))$.
- (4) If (H, C) is a neutrosophic soft set over U and $(F, A) \subseteq (G, B) \subseteq (H, C)$, then

$$CS_k((F, A), (H, C)) \leq CS_k((F, A), (G, B))$$

and

$$CS_k((F, A), (H, C)) \leq CS_k((G, B), (H, C)).$$

Proof. (1) Since the truth membership degree, indeterminacy- membership degree and falsity- membership degree in neutrosophic set and the value of cosine function are within $[0, 1]$, the similarity measure based on cosine function is also within $[0, 1]$. Thus $0 \leq CS_k((F, A), (G, B)) \leq 1$ for $k = 1, 2$.

- (2) For any two neutrosophic soft sets, $(F, A) = (G, B)$ implies

$$T_{F(e_i)}(x_j) = T_{G(e_i)}(x_j), I_{F(e_i)}(x_j) = I_{G(e_i)}(x_j), F_{F(e_i)}(x_j) = F_{G(e_i)}(x_j)$$

for all $i = 1, 2, \dots, m$ and $j = 1, 2, \dots, n$. Thus

$$|T_{F(e_i)}(x_j) - T_{G(e_i)}(x_j)| = |I_{F(e_i)}(x_j) - I_{G(e_i)}(x_j)| = |F_{F(e_i)}(x_j) - F_{G(e_i)}(x_j)| = 0.$$

So $CS_k((F, A), (G, B)) = 1$ for $k = 1, 2$.

- Conversely, if $CS_k((F, A), (G, B)) = 1$ for $k = 1, 2$, this implies

$$|T_{F(e_i)}(x_j) - T_{G(e_i)}(x_j)| = 0, |I_{F(e_i)}(x_j) - I_{G(e_i)}(x_j)| = 0, |F_{F(e_i)}(x_j) - F_{G(e_i)}(x_j)| = 0$$

since $\cos(0) = 1$. Then these equalities indicate

$$T_{F(e_i)}(x_j) = T_{G(e_i)}(x_j), I_{F(e_i)}(x_j) = I_{G(e_i)}(x_j), F_{F(e_i)}(x_j) = F_{G(e_i)}(x_j)$$

for all $i = 1, 2, \dots, m$ and $j = 1, 2, \dots, n$. Hence $(F, A) = (G, B)$.

- (3) Clearly $CS_k((F, A), (G, B)) = CS_k((G, B), (F, A))$.
- (4) If $(F, A) \subseteq (G, B) \subseteq (H, C)$, then

$$T_{F(e_i)}(x_j) \leq T_{G(e_i)}(x_j) \leq T_{H(e_i)}(x_j),$$

$$I_{F(e_i)}(x_j) \leq I_{G(e_i)}(x_j) \leq I_{H(e_i)}(x_j),$$

$$F_{F(e_i)}(x_j) \geq F_{G(e_i)}(x_j) \geq F_{H(e_i)}(x_j)$$

for all $i = 1, 2, \dots, m$ and $j = 1, 2, \dots, n$.

Then we have the following inequalities :

$$\begin{aligned} |T_{F(e_i)}(x_j) - T_{G(e_i)}(x_j)| &\leq |T_{F(e_i)}(x_j) - T_{H(e_i)}(x_j)|, \\ |T_{G(e_i)}(x_j) - T_{H(e_i)}(x_j)| &\leq |T_{F(e_i)}(x_j) - T_{H(e_i)}(x_j)|, \\ |I_{F(e_i)}(x_j) - I_{G(e_i)}(x_j)| &\leq |I_{F(e_i)}(x_j) - I_{H(e_i)}(x_j)|, \\ |I_{G(e_i)}(x_j) - I_{H(e_i)}(x_j)| &\leq |I_{F(e_i)}(x_j) - I_{H(e_i)}(x_j)|, \\ |F_{F(e_i)}(x_j) - F_{G(e_i)}(x_j)| &\leq |F_{F(e_i)}(x_j) - F_{H(e_i)}(x_j)|, \\ |F_{G(e_i)}(x_j) - F_{H(e_i)}(x_j)| &\leq |F_{F(e_i)}(x_j) - F_{H(e_i)}(x_j)|. \end{aligned}$$

Thus

$$CS_k((F, A), (H, C)) \leq CS_k((F, A), (G, B))$$

and

$$CS_k((F, A), (H, C)) \leq CS_k((G, B), (H, C))$$

for $k = 1, 2$, since cosine function is a decreasing function within the interval $\left[0, \frac{\pi}{2}\right]$. \square

Definition 3.3. Let $U = \{x_1, x_2, \dots, x_n\}$ be the universal set of elements and $E = \{e_1, e_2, \dots, e_m\}$ be the universal set of parameters and (F, A) and (G, B) be two neutrosophic soft sets over U . Now we consider the weights w_j of x_j ($j = 1, 2, \dots, n$) then the weighted cosine similarity measures between (F, A) and (G, B) is defined as follows:

$$\begin{aligned} &WCS_1((F, A), (G, B)) \\ &= \frac{1}{m} \sum_{i=1}^m \sum_{j=1}^n w_j \cos \left[\frac{\pi}{2} (|T_{F(e_i)}(x_j) - T_{G(e_i)}(x_j)|) \vee (|I_{F(e_i)}(x_j) - I_{G(e_i)}(x_j)|) \vee \right. \\ &\qquad \qquad \qquad \left. (|F_{F(e_i)}(x_j) - F_{G(e_i)}(x_j)|) \right], \\ &WCS_2((F, A), (G, B)) \\ &= \frac{1}{m} \sum_{i=1}^m \sum_{j=1}^n w_j \cos \left[\frac{\pi}{6} (|T_{F(e_i)}(x_j) - T_{G(e_i)}(x_j)|) + (|I_{F(e_i)}(x_j) - I_{G(e_i)}(x_j)|) + \right. \\ &\qquad \qquad \qquad \left. (|F_{F(e_i)}(x_j) - F_{G(e_i)}(x_j)|) \right], \end{aligned}$$

where $w_j \in [0, 1]$, $j = 1, 2, \dots, n$ and $\sum_{j=1}^n w_j = 1$.

In particular if we take $w_j = \frac{1}{n}$, $j = 1, 2, 3, \dots, n$, then

$$WCS_k((F, A), (G, B)) = CS_k((F, A), (G, B)),$$

$k = 1, 2$.

Definition 3.4. Let $U = \{x_1, x_2, \dots, x_n\}$ be the universal set of elements and $E = \{e_1, e_2, \dots, e_m\}$ be the universal set of parameters, and (F, A) and (G, B) be two interval valued neutrosophic soft sets over U . Then the cosine similarity measures between (F, A) and (G, B) is defined as

$$CS_3((F, A), (G, B))$$

$$\begin{aligned}
 &= \frac{1}{mn} \sum_{i=1}^m \sum_{j=1}^n \cos \left[\frac{\pi}{4} ((|\underline{T}_{F(e_i)}(x_j) - \underline{T}_{G(e_i)}(x_j)|) \vee (|\underline{I}_{F(e_i)}(x_j) - \underline{I}_{G(e_i)}(x_j)|) \vee \right. \\
 &\quad \left. (|\underline{F}_{F(e_i)}(x_j) - \underline{F}_{G(e_i)}(x_j)|) + (|\overline{T}_{F(e_i)}(x_j) - \overline{T}_{G(e_i)}(x_j)|) \vee \right. \\
 &\quad \left. (|\overline{I}_{F(e_i)}(x_j) - \overline{I}_{G(e_i)}(x_j)|) \vee (|\overline{F}_{F(e_i)}(x_j) - \overline{F}_{G(e_i)}(x_j)|) \right)]. \\
 &CS_4((F, A), (G, B)) \\
 &= \frac{1}{mn} \sum_{i=1}^m \sum_{j=1}^n \cos \left[\frac{\pi}{12} ((|\underline{T}_{F(e_i)}(x_j) - \underline{T}_{G(e_i)}(x_j)|) + (|\underline{I}_{F(e_i)}(x_j) - \underline{I}_{G(e_i)}(x_j)|) + \right. \\
 &\quad \left. (|\underline{F}_{F(e_i)}(x_j) - \underline{F}_{G(e_i)}(x_j)|) + (|\overline{T}_{F(e_i)}(x_j) - \overline{T}_{G(e_i)}(x_j)|) + \right. \\
 &\quad \left. (|\overline{I}_{F(e_i)}(x_j) - \overline{I}_{G(e_i)}(x_j)|) + (|\overline{F}_{F(e_i)}(x_j) - \overline{F}_{G(e_i)}(x_j)|) \right)],
 \end{aligned}$$

where the symbol \vee is the maximum operation. The cosine similarity measures $CS_k((F, A), (G, B))$, $k = 3, 4$ satisfy the following properties.

Proposition 3.5. *Let $U = \{x_1, x_2, \dots, x_n\}$ be the universal set of elements and $E = \{e_1, e_2, \dots, e_m\}$ be the universal set of parameters. Then for two interval valued neutrosophic soft sets (F, A) and (G, B) over U the cosine similarity measure $CS_k((F, A), (G, B))$ ($k=3,4$) should satisfy the following properties (1)–(4).*

- (1) $0 \leq CS_k((F, A), (G, B)) \leq 1$.
- (2) $CS_k((F, A), (G, B)) = 1$ if and only if $(F, A) = (G, B)$.
- (3) $CS_k((F, A), (G, B)) = CS_k((G, B), (F, A))$.
- (4) If (H, C) is a interval valued neutrosophic soft set over U and $(F, A) \subseteq (G, B) \subseteq (H, C)$, then

$$CS_k((F, A), (H, C)) \leq CS_k((F, A), (G, B))$$

and

$$CS_k((F, A), (H, C)) \leq CS_k((G, B), (H, C)).$$

Proof. (1) Since the truth membership degree, indeterminacy- membership degree and falsity- membership degree in neutrosophic set and the value of cosine function are within $[0,1]$, the similarity measure based on cosine function is also within $[0,1]$. Then $0 \leq CS_k((F, A), (G, B)) \leq 1$ for $k = 3, 4$.

(2) For any two interval valued neutrosophic soft sets, $(F, A) = (G, B)$ implies

$$T_{F(e_i)}(x_j) = T_{G(e_i)}(x_j), I_{F(e_i)}(x_j) = I_{G(e_i)}(x_j), F_{F(e_i)}(x_j) = F_{G(e_i)}(x_j)$$

for all $i = 1, 2, \dots, m$ and $j = 1, 2, \dots, n$. Then

$$|\underline{T}_{F(e_i)}(x_j) - \underline{T}_{G(e_i)}(x_j)| = |\underline{I}_{F(e_i)}(x_j) - \underline{I}_{G(e_i)}(x_j)| = |\underline{F}_{F(e_i)}(x_j) - \underline{F}_{G(e_i)}(x_j)| = 0$$

and

$$|\overline{T}_{F(e_i)}(x_j) - \overline{T}_{G(e_i)}(x_j)| = |\overline{I}_{F(e_i)}(x_j) - \overline{I}_{G(e_i)}(x_j)| = |\overline{F}_{F(e_i)}(x_j) - \overline{F}_{G(e_i)}(x_j)| = 0.$$

Thus $CS_k((F, A), (G, B)) = 1$ for $k = 3, 4$.

Conversely, if $CS_k((F, A), (G, B)) = 1$ for $k = 3, 4$, then, since $\cos(0) = 1$,

$$|\underline{T}_{F(e_i)}(x_j) - \underline{T}_{G(e_i)}(x_j)| = 0,$$

$$|\underline{I}_{F(e_i)}(x_j) - \underline{I}_{G(e_i)}(x_j)| = 0,$$

$$|\underline{F}_{F(e_i)}(x_j) - \underline{F}_{G(e_i)}(x_j)| = 0,$$

$$|\overline{T}_{F(e_i)}(x_j) - \overline{T}_{G(e_i)}(x_j)| = |\overline{I}_{F(e_i)}(x_j) - \overline{I}_{G(e_i)}(x_j)| = |\overline{F}_{F(e_i)}(x_j) - \overline{F}_{G(e_i)}(x_j)| = 0.$$

Then these equalities indicate

$$T_{F(e_i)}(x_j) = T_{G(e_i)}(x_j), I_{F(e_i)}(x_j) = I_{G(e_i)}(x_j), F_{F(e_i)}(x_j) = F_{G(e_i)}(x_j)$$

for all $i = 1, 2, \dots, m$ and $j = 1, 2, \dots, n$. Thus $(F, A) = (G, B)$.

(3) Clearly $CS_k((F, A), (G, B)) = CS_k((G, B), (F, A))$.

(4) If $(F, A) \subseteq (G, B) \subseteq (H, C)$, then

$$\underline{T}_{F(e_i)}(x_j) \leq \underline{T}_{G(e_i)}(x_j) \leq \underline{T}_{H(e_i)}(x_j),$$

$$\overline{T}_{F(e_i)}(x_j) \leq \overline{T}_{G(e_i)}(x_j) \leq \overline{T}_{H(e_i)}(x_j),$$

$$\underline{I}_{F(e_i)}(x_j) \leq \underline{I}_{G(e_i)}(x_j) \leq \underline{I}_{H(e_i)}(x_j),$$

$$\overline{I}_{F(e_i)}(x_j) \leq \overline{I}_{G(e_i)}(x_j) \leq \overline{I}_{H(e_i)}(x_j),$$

$$\underline{F}_{F(e_i)}(x_j) \geq \underline{F}_{G(e_i)}(x_j) \geq \underline{F}_{H(e_i)}(x_j),$$

$$\overline{F}_{F(e_i)}(x_j) \geq \overline{F}_{G(e_i)}(x_j) \geq \overline{F}_{H(e_i)}(x_j),$$

for all $i = 1, 2, \dots, m$ and $j = 1, 2, \dots, n$. Thus we have the following inequalities:

$$|\underline{T}_{F(e_i)}(x_j) - \underline{T}_{G(e_i)}(x_j)| \leq |\underline{T}_{F(e_i)}(x_j) - \underline{T}_{H(e_i)}(x_j)|,$$

$$|\underline{T}_{G(e_i)}(x_j) - \underline{T}_{H(e_i)}(x_j)| \leq |\underline{T}_{F(e_i)}(x_j) - \underline{T}_{H(e_i)}(x_j)|,$$

$$|\overline{T}_{F(e_i)}(x_j) - \overline{T}_{G(e_i)}(x_j)| \leq |\overline{T}_{F(e_i)}(x_j) - \overline{T}_{H(e_i)}(x_j)|,$$

$$|\overline{T}_{G(e_i)}(x_j) - \overline{T}_{H(e_i)}(x_j)| \leq |\overline{T}_{F(e_i)}(x_j) - \overline{T}_{H(e_i)}(x_j)|,$$

$$|\underline{I}_{F(e_i)}(x_j) - \underline{I}_{G(e_i)}(x_j)| \leq |\underline{I}_{F(e_i)}(x_j) - \underline{I}_{H(e_i)}(x_j)|,$$

$$|\underline{I}_{G(e_i)}(x_j) - \underline{I}_{H(e_i)}(x_j)| \leq |\underline{I}_{F(e_i)}(x_j) - \underline{I}_{H(e_i)}(x_j)|,$$

$$|\overline{I}_{F(e_i)}(x_j) - \overline{I}_{G(e_i)}(x_j)| \leq |\overline{I}_{F(e_i)}(x_j) - \overline{I}_{H(e_i)}(x_j)|,$$

$$|\overline{I}_{G(e_i)}(x_j) - \overline{I}_{H(e_i)}(x_j)| \leq |\overline{I}_{F(e_i)}(x_j) - \overline{I}_{H(e_i)}(x_j)|,$$

$$|\underline{F}_{F(e_i)}(x_j) - \underline{F}_{G(e_i)}(x_j)| \leq |\underline{F}_{F(e_i)}(x_j) - \underline{F}_{H(e_i)}(x_j)|,$$

$$|\underline{F}_{G(e_i)}(x_j) - \underline{F}_{H(e_i)}(x_j)| \leq |\underline{F}_{F(e_i)}(x_j) - \underline{F}_{H(e_i)}(x_j)|,$$

$$|\overline{F}_{F(e_i)}(x_j) - \overline{F}_{G(e_i)}(x_j)| \leq |\overline{F}_{F(e_i)}(x_j) - \overline{F}_{H(e_i)}(x_j)|,$$

$$|\overline{F}_{G(e_i)}(x_j) - \overline{F}_{H(e_i)}(x_j)| \leq |\overline{F}_{F(e_i)}(x_j) - \overline{F}_{H(e_i)}(x_j)|.$$

Thus

$$CS_k((F, A), (H, C)) \leq CS_k((F, A), (G, B))$$

and

$$CS_k((F, A), (H, C)) \leq CS_k((G, B), (H, C))$$

for $k = 3, 4$, since cosine function is a decreasing function within the interval $\left[0, \frac{\pi}{2}\right]$. \square

Definition 3.6. Let $U = \{x_1, x_2, \dots, x_n\}$ be the universal set of elements and $E = \{e_1, e_2, \dots, e_m\}$ be the universal set of parameters, and (F, A) and (G, B) be two interval valued neutrosophic soft sets over U . Then the weighted cosine similarity measures between (F, A) and (G, B) is defined as

$$\begin{aligned}
 & CS_3((F, A), (G, B)) \\
 &= \frac{1}{m} \sum_{i=1}^m \sum_{j=1}^n w_j \cos \left[\frac{\pi}{4} ((|\underline{T}_{F(e_i)}(x_j) - \underline{T}_{G(e_i)}(x_j)|) \vee (|\underline{I}_{F(e_i)}(x_j) - \underline{I}_{G(e_i)}(x_j)|) \vee \right. \\
 &\qquad\qquad\qquad (|\underline{F}_{F(e_i)}(x_j) - \underline{F}_{G(e_i)}(x_j)|) + (|\overline{T}_{F(e_i)}(x_j) - \overline{T}_{G(e_i)}(x_j)|) \vee \\
 &\qquad\qquad\qquad \left. (|\overline{I}_{F(e_i)}(x_j) - \overline{I}_{G(e_i)}(x_j)|) \vee (|\overline{F}_{F(e_i)}(x_j) - \overline{F}_{G(e_i)}(x_j)|) \right), \\
 & CS_4((F, A), (G, B)) \\
 &= \frac{1}{m} \sum_{i=1}^m \sum_{j=1}^n w_j \cos \left[\frac{\pi}{12} ((|\underline{T}_{F(e_i)}(x_j) - \underline{T}_{G(e_i)}(x_j)|) + (|\underline{I}_{F(e_i)}(x_j) - \underline{I}_{G(e_i)}(x_j)|) + \right. \\
 &\qquad\qquad\qquad (|\underline{F}_{F(e_i)}(x_j) - \underline{F}_{G(e_i)}(x_j)|) + (|\overline{T}_{F(e_i)}(x_j) - \overline{T}_{G(e_i)}(x_j)|) + \\
 &\qquad\qquad\qquad \left. (|\overline{I}_{F(e_i)}(x_j) - \overline{I}_{G(e_i)}(x_j)|) + (|\overline{F}_{F(e_i)}(x_j) - \overline{F}_{G(e_i)}(x_j)|) \right),
 \end{aligned}$$

where $w_j \in [0, 1]$, $j=1, 2, \dots, n$ and $\sum_{j=1}^n w_j = 1$.

In particular, if we take $w_j = \frac{1}{n}$, $j=1, 2, 3, \dots, n$, then

$$WCS_k((F, A), (G, B)) = CS_k((F, A), (G, B)), \quad k=3, 4.$$

4. APPLICATION OF COSINE SIMILARITY MEASURE OF NEUTROSOPHIC SOFT SET

Example 4.1. Consider areas of a state are affected by flood. A team of three members $U = \{m_1, m_2, m_3\}$ from Rehabilitation department inspect the flood affected areas and the relief measures recommended by them are described by the parameter set $E = \{e_1, e_2, e_3, e_4\}$ where $e_1 =$ relief for crop loss, $e_2 =$ relief for ecological damage, $e_3 =$ relief for livestock and $e_4 =$ provision for alternate livelihood. Based on these recommendations the government has to allocate funds according to the level of damage. Algorithm:

Step 1. An interval valued neutrosophic soft set (F, A) over U is constructed based on previous records of relief measures in similar situation.

Step 2. An interval valued neutrosophic soft set (G, B) over U based on the recommendations of the team visiting Area I is constructed.

Step 3. Cosine similarity measures between (F, A) and (G, B) is calculated.

Step 4. An interval valued neutrosophic soft set (H, C) over U based on the recommendations of the team visiting Area II is constructed.

Step 5. Cosine similarity measures between (F, A) and (H, C) is calculated.

Step 6. Estimate result by using the similarity value. An interval valued (F, A) over U with $A = E$ based on the previous records of relief measures in similar situation is given by.

$$\begin{aligned}
 (F, A) = & \{F(e_1)(m_1) = ([0.6, 0.7], [0.3, 0.4], [0.2, 0.3]), F(e_2)(m_1) = ([0.4, 0.5], [0.3, 0.4], [0.2, 0.3]), \\
 & F(e_3)(m_1) = ([0.5, 0.6], [0.4, 0.5], [0.1, 0.2]), F(e_4)(m_1) = ([0.5, 0.6], [0.2, 0.3], [0.3, 0.4]), \\
 & F(e_1)(m_2) = ([0.5, 0.6], [0.3, 0.4], [0.2, 0.3]), F(e_2)(m_2) = ([0.4, 0.5], [0.2, 0.3], [0.5, 0.6]), \\
 & F(e_3)(m_2) = ([0.5, 0.6], [0.1, 0.2], [0.3, 0.4]), F(e_4)(m_2) = ([0.5, 0.6], [0.3, 0.5], [0.4, 0.5]), \\
 & F(e_1)(m_3) = ([0.7, 0.8], [0.4, 0.5], [0.2, 0.3]), F(e_2)(m_3) = ([0.5, 0.6], [0.3, 0.4], [0.2, 0.3]),
 \end{aligned}$$

$F(e_3)(m_3) = ([0.6, 0.7], [0.4, 0.5], [0.3, 0.4]), F(e_4)(m_3) = ([0.6, 0.7], [0.2, 0.3], [0.4, 0.5])$.

An interval valued (G,B) over U with B = E based on recommendations of the expert team visiting Area I.

(G,B)=

$\{G(e_1)(m_1) = ([0.6, 0.7], [0.5, 0.6], [0.4, 0.5]), G(e_2)(m_1) = ([0.3, 0.4], [0.5, 0.6], [0.4, 0.5]),$
 $G(e_3)(m_1) = ([0.3, 0.4], [0.4, 0.5], [0.5, 0.6]), G(e_4)(m_1) = ([0.2, 0.3], [0.3, 0.4], [0.5, 0.6]),$
 $G(e_1)(m_2) = ([0.8, 0.9], [0.4, 0.5], [0.2, 0.3]), G(e_2)(m_2) = ([0.5, 0.6], [0.1, 0.2], [0.3, 0.4]),$
 $G(e_3)(m_2) = ([0.5, 0.6], [0.1, 0.2], [0.2, 0.3]), G(e_4)(m_2) = ([0.3, 0.4], [0.3, 0.4], [0.2, 0.3]),$
 $G(e_1)(m_3) = ([0.5, 0.6], [0.4, 0.5], [0.3, 0.4]), G(e_2)(m_3) = ([0.4, 0.5], [0.2, 0.3], [0.5, 0.6]),$
 $G(e_3)(m_3) = ([0.2, 0.3], [0.4, 0.5], [0.4, 0.5]), G(e_4)(m_3) = ([0.2, 0.3], [0.5, 0.6], [0.3, 0.4])\}.$

An interval valued (H,C) over U with C= E based on recommendations of the expert team visiting Area II.

(H,C)=

$\{H(e_1)(m_1) = ([0.4, 0.5], [0.2, 0.3], [0.1, 0.2]), H(e_2)(m_1) = ([0.3, 0.4], [0.2, 0.3], [0.1, 0.2]),$
 $H(e_3)(m_1) = ([0.6, 0.7], [0.2, 0.3], [0.1, 0.2]), H(e_4)(m_1) = ([0.1, 0.2], [0.3, 0.4], [0.6, 0.7]),$
 $H(e_1)(m_2) = ([0.4, 0.5], [0.2, 0.3], [0.5, 0.6]), H(e_2)(m_2) = ([0.2, 0.3], [0.3, 0.4], [0.4, 0.5]),$
 $H(e_3)(m_2) = ([0.1, 0.2], [0.5, 0.6], [0.4, 0.5]), H(e_4)(m_2) = ([0.5, 0.6], [0.3, 0.4], [0.1, 0.2]),$
 $H(e_1)(m_3) = ([0.2, 0.3], [0.4, 0.5], [0.3, 0.4]), H(e_2)(m_3) = ([0.1, 0.2], [0.3, 0.4], [0.4, 0.5]),$
 $H(e_3)(m_3) = ([0.2, 0.3], [0.4, 0.5], [0.4, 0.5]), H(e_4)(m_3) = ([0.3, 0.4], [0.5, 0.6], [0.6, 0.7])\}.$

Using Definition 3.4, we get

$$CS_3((F, A), (G, B)) = 0.904$$

and

$$CS_3((F, A), (H, C)) = 0.871.$$

We have $CS_3((F, A), (G, B)) > CS_3((F, A), (H, C))$. Hence we conclude that Area I is severely affected by flood.

REFERENCES

- [1] I. Arockiarani, I. R. Sumathi and J. Martina Jency, Fuzzy Neutrosophic Soft Topological Spaces, International journal of mathematical archives 4 (10) (2013) 225–238.
- [2] I. Arockiarani, I. R. Sumathi, Some results on interval valued Fuzzy neutrosophic soft set, International journal of innovative research and studies 3 (5) (2014) 386–405.
- [3] W. L. Hung and M. S. Yang, Similarity measures of intuitionistic fuzzy sets based on Hausdorff distance, Pattern Recognition Letters 25 (2004) 1603–1611.
- [4] Jun Ye, Entropy, Cosine similarity measures for intuitionistic fuzzy sets and their applications, Mathematical and computer modelling 53 (1-2) (2011) 91–97.
- [5] Jun Ye, Vector similarity measures of simplified neutrosophic sets and their application in multicriteria Decision Making, International Journal of Fuzzy Systems 16 (2) (2014) 204–211.
- [6] P. Majumdar and S. K. Samanta, On similarity and entropy of neutrosophic sets, J. Intell. Fuzzy Systems 26 (2014) 1245–1252.
- [7] A. A. Salama and Florentin Smarandache, Neutrosophic Crisp Set Theory, 2015 USA Book, Educational. Education Publishing 1313 Chesapeake, Avenue, Columbus, Ohio 43212, USA.
- [8] A. A. Salama and Said Broumi, Roughness of Neutrosophic Sets, Elixir Appl. Math. 74 (2014) 26833–26837.
- [9] A. A. Salama, Basic Structure of Some Classes of Neutrosophic Crisp Nearly Open Sets and Possible Application to GIS Topology, Neutrosophic Sets and Systems 7 (2015) 18–22.
- [10] A. A. Salama, Mohamed Abdelfattah and Mohamed Eisa, Distances, Hesitancy Degree and Flexible Querying via Neutrosophic Sets, International Journal of Computer Applications, 101 (10) (2014) 7–12.

- [11] A. A. Salama, Haithem A. El-Ghareeb, Ayman M. Mainie and Florentin Smarandache, Introduction to Develop Some Software Programs for dealing with Neutrosophic Sets, *Neutrosophic Sets and Systems* 4 (2014) 51–52.
- [12] A. A. Salama, F. Smarandache, and M. Eisa, Introduction to Image Processing via Neutrosophic Technique, *Neutrosophic Sets and Systems* 5 (2014) 59–63.
- [13] A. A. Salama, Mohamed Abdelfattah and Mohamed Eisa, A Novel Model for Implementing Security over Mobile Ad-hoc Networks using Intuitionistic Fuzzy Function, *International Journal of Emerging Technologies in Computational and Applied Sciences (IJETCAS)*, 7 (1) (2014) 1–7.
- [14] A. A. Salama, Mohamed Eisa and M. M. Abdelmoghny, Neutrosophic Relations Database, *International Journal of Information Science and Intelligent System* 3 (1) (2014) 33–46 .
- [15] A. A. Salama, Haitham A. El-Ghareeb, Ayman M. Manie and M. M. Lotfy, Utilizing Neutrosophic Set in Social Network Analysis e-Learning Systems, *International Journal of Information Science and Intelligent System* 3 (2) (2014) 61–72.
- [16] A. A. Salama, Mohamed Eisa, S. A. ELhafeez and M. M. Lotfy, Review of Recommender Systems Algorithms Utilized in Social Networks based e-Learning Systems and Neutrosophic System, *Neutrosophic Sets and Systems* 8 (2015) 35–44.
- [17] A. A. Salama and Florentin Smarandache, Neutrosophic Ideal Theory Neutrosophic Local Function and Generated Neutrosophic Topology, In *Neutrosophic Theory and Its Applications, Collected Papers, Volume 1, EuropaNova, Bruxelles* (2014) 213–218.
- [18] A. Salama, Said Broumi and Florentin Smarandache, Some Types of Neutrosophic Crisp Sets and Neutrosophic Crisp Relations, *I.J. Information Engineering and Electronic Business* 2014.
- [19] S. A. Alblowi, A. A. Salama and Mohamed Eisa, New concepts of neutrosophic sets, *international journal of mathematics and computer applications research (ijmcar)* 4 (1) (2014) 59–66.
- [20] A. A. Salama, Neutrosophic Crisp Points and Neutrosophic Crisp Ideals, *Neutrosophic Sets and Systems* 1 (1) (2013) 50–54.
- [21] A. A. Salama and F. Smarandache, Filters via Neutrosophic Crisp Sets, *Neutrosophic Sets and Systems*, 1 (1) (2013) 34–38.
- [22] A. A. Salama, F. Smarandache and S. A. Alblowi, The Characteristic Function of a Neutrosophic Set, *Neutrosophic Sets and Systems* 3 (2014) 14–17.
- [23] Florentin Smarandache, *Neutrosophy: Neutrosophic Probability, Set, and Logic*, Amer. Res. Press, Rehoboth, USA 105 p., 1998.
- [24] F. Smarandache, *A Unifying Field in Logics, Neutrosophy: Neutrosophic Probability, Set and Logic*, American Research Press 2005.
- [25] F. Smarandache, *Neutrosophy and Neutrosophic Logic*, First International Conference On Neutrosophy, Neutrosophic Logic, Set, Probability and Statistics University of New Mexico, Gallup, NM 87301, USA 2002.
- [26] C. Wang and A. Qu, Entropy, similarity measure and distance measure of vague soft sets and their relations, *Inform. Sci.* 244 (2013) 92–106.

I. R. SUMATHI (sumathi_raman2005@yahoo.co.in)

Department of Mathematics, Kumaraguru College of Technology, Coimbatore, Tamilnadu, India

I. AROCKIARANI (stell11960@gmail.com)

Department of Mathematics, Nirmala College for women, Coimbatore, Tamilnadu, India