


Wijsman rough I-convergence limit point of triple sequences defined by a metric function

AYHAN ESI, NAGARAJAN SUBRAMANIAN, AYTEN ESI



Reprinted from the
 Annals of Fuzzy Mathematics and Informatics
 Vol. 15, No. 1, February 2018

Wijsman rough I-convergence limit point of triple sequences defined by a metric function

AYHAN ESI, NAGARAJAN SUBRAMANIAN, AYTEN ESI

Received 31 October 2017; Revised 23 November 2017; Accepted 8 December 2017

ABSTRACT. We introduce and study some basic properties of Wijsman rough I -convergent of triple sequence and also study the set of all rough I -limits of a triple sequence.

2010 AMS Classification: 40F05, 40J05, 40G05

Keywords: Triple sequences, Wijsman rough convergence, Strongly admissible ideal, Cluster points.

Corresponding Author: Ayhan Esi (aesi23@hotmail.com)

1. INTRODUCTION

The idea of statistical convergence was introduced by Steinhaus [13] and also independently by Fast [8] for real or complex sequences. Statistical convergence is a generalization of the usual notion of convergence, which parallels the theory of ordinary convergence.

Let K be a subset of the set of positive integers $\mathbb{N} \times \mathbb{N} \times \mathbb{N}$, and let us denote the set $\{(m, n, k) \in K : m \leq u, n \leq v, k \leq w\}$ by K_{uvw} . Then the natural density of K is given by $\delta(K) = \lim_{uvw \rightarrow \infty} \frac{|K_{uvw}|}{uvw}$, where $|K_{uvw}|$ denotes the number of elements in K_{uvw} . Clearly, a finite subset has natural density zero, and we have $\delta(K^c) = 1 - \delta(K)$ where $K^c = \mathbb{N} \setminus K$ is the complement of K . If $K_1 \subseteq K_2$, then $\delta(K_1) \leq \delta(K_2)$.

Throughout the paper, \mathbb{R} denotes the real of three dimensional space with metric (X, d) . Consider a triple sequence $x = (x_{mnk})$ such that $x_{mnk} \in \mathbb{R}, m, n, k \in \mathbb{N}$.

A triple sequence $x = (x_{mnk})$ is said to be statistically convergent to $0 \in \mathbb{R}$, written as $st - \lim x = 0$, provided that the set

$$\{(m, n, k) \in \mathbb{N}^3 : |x_{mnk}, 0| \geq \epsilon\}$$

has natural density zero for any $\epsilon > 0$. In this case, 0 is called the statistical limit of the triple sequence x .

If a triple sequence is statistically convergent, then for every $\epsilon > 0$, infinitely many terms of the sequence may remain outside the ϵ -neighborhood of the statistical limit, provided that the natural density of the set consisting of the indices of these terms is zero. This is an important property that distinguishes statistical convergence from ordinary convergence. Because the natural density of a finite set is zero, we can say that every ordinary convergent sequence is statistically convergent.

If a triple sequence $x = (x_{mnk})$ satisfies some property P for all m, n, k except a set of natural density zero, then we say that the triple sequence x satisfies P for almost all (m, n, k) and we abbreviate this by a.a. (m, n, k) .

Let $(x_{m_i n_j k_\ell})$ be a sub sequence of $x = (x_{mnk})$. If the natural density of the set $K = \{(m_i, n_j, k_\ell) \in \mathbb{N}^3 : (i, j, \ell) \in \mathbb{N}^3\}$ is different from zero, then $(x_{m_i n_j k_\ell})$ is called a non-thin subsequence of a triple sequence x .

$\xi \in \mathbb{R}$ is called a statistical cluster point of a triple sequence $x = (x_{mnk})$ provided that the natural density of the set

$$\{(m, n, k) \in \mathbb{N}^3 : |x_{mnk} - \xi| < \epsilon\}$$

is different from zero for every $\epsilon > 0$. We denote the set of all statistical cluster points of the sequence x by Γ_x .

A triple sequence $x = (x_{mnk})$ is said to be statistically analytic if there exists a positive number M such that

$$\delta \left(\left\{ (m, n, k) \in \mathbb{N}^3 : |x_{mnk}|^{1/(m+n+k)} \geq M \right\} \right) = 0$$

The theory of statistical convergence has been discussed in trigonometric series, summability theory, measure theory, turnpike theory, approximation theory, fuzzy set theory and so on. For example one may refer to Tripathy and Goswami [15, 16, 17, 18].

The idea of rough convergence was introduced by Phu [10], who also introduced the concepts of rough limit points and roughness degree. The idea of rough convergence occurs very naturally in numerical analysis and has interesting applications. Aytar [2] extended the idea of rough convergence into rough statistical convergence using the notion of natural density just as usual convergence was extended to statistical convergence. Pal et al. [9] extended the notion of rough convergence using the concept of ideals which automatically extends the earlier notions of rough convergence and rough statistical convergence.

Let (X, ρ) be a metric space. For any non empty closed subsets $A, A_{mnk} \subset X$ ($m, n, k \in \mathbb{N}$), we say that the triple sequence (A_{mnk}) is Wijsman statistical convergent to A if the triple sequence $(d(x, A_{mnk}))$ is statistically convergent to $d(x, A)$, i.e., for $\epsilon > 0$ and for each $x \in X$

$$\lim_{rst} \frac{1}{rst} |\{m \leq r, n \leq s, k \leq t : |d(x, A_{mnk}) - d(x, A)| \geq \epsilon\}| = 0.$$

In this case, we write $St - \lim_{mnk} A_{mnk} = A$ or $A_{mnk} \rightarrow A (WS)$. The triple sequence (A_{mnk}) is bounded if $\sup_{mnk} d(x, A_{mnk}) < \infty$ for each $x \in X$.

In this paper, we introduce the notion of Wijsman rough statistical convergence of triple sequences. Defining the set of Wijsman rough statistical limit points of a triple sequence, we obtain to Wijsman statistical convergence criteria associated

with this set. Later, we prove that this set of Wijsman statistical cluster points and the set of Wijsman rough statistical limit points of a triple sequence.

A triple sequence (real or complex) can be defined as a function $x : \mathbb{N} \times \mathbb{N} \times \mathbb{N} \rightarrow \mathbb{R}(\mathbb{C})$, where \mathbb{N} , \mathbb{R} and \mathbb{C} denote the set of natural numbers, real numbers and complex numbers respectively. The different types of notions of triple sequence was introduced and investigated at the initial by Sahiner et al. [11, 12], Esi et al. [5, 6, 7], Dutta et al. [4], Subramanian et al. [14], Debnath et al. [3], Aiyub et al. [1] and many others.

2. DEFINITIONS AND PRELIMINARIES

Definition 2.1. A triple sequence $x = (x_{mnk})$ of real numbers is said to be statistically convergent to $l \in \mathbb{R}^3$ if for any $\epsilon > 0$ we have $d(A(\epsilon)) = 0$, where

$$A(\epsilon) = \{(m, n, k) \in \mathbb{N}^3 : |x_{mnk} - l| \geq \epsilon\}.$$

Definition 2.2. A triple sequence $x = (x_{mnk})$ is said to be statistically convergent to $l \in \mathbb{R}^3$, written as $st - \lim x = l$, provided that the set

$$\{(m, n, k) \in \mathbb{N}^3 : |x_{mnk} - l| \geq \epsilon\}$$

has natural density zero for every $\epsilon > 0$.

In this case, l is called the statistical limit of the sequence x .

Definition 2.3. Let $x = (x_{mnk})_{m,n,k \in \mathbb{N} \times \mathbb{N} \times \mathbb{N}}$ be a triple sequence in a metric space $(X, |., .|)$ and r be a non-negative real number. A triple sequence $x = (x_{mnk})$ is said to be r -convergent to $l \in X$, denoted by $x \rightarrow^r l$, if for any $\epsilon > 0$ there exists $N_\epsilon \in \mathbb{N} \times \mathbb{N} \times \mathbb{N}$ such that for all $m, n, k \geq N_\epsilon$ we have

$$|x_{mnk} - l| < r + \epsilon$$

In this case, l is called an r -limit of x .

Remark 2.4. We consider r -limit set x which is denoted by LIM_x^r and is defined by

$$LIM_x^r = \{l \in X : x \rightarrow^r l\}.$$

Definition 2.5. A triple sequence $x = (x_{mnk})$ is said to be r -convergent, if $LIM_x^r \neq \emptyset$ and r is called a rough convergence degree of x . If $r = 0$, then it is ordinary convergence of triple sequence.

Definition 2.6. Let $x = (x_{mnk})$ be a triple sequence in a metric space $(X, |., .|)$ and r be a non-negative real number is said to be r -statistically convergent to l , denoted by $x \rightarrow^{r-st} l$, if for any $\epsilon > 0$ we have $d(A(\epsilon)) = 0$, where

$$A(\epsilon) = \{(m, n, k) \in \mathbb{N} \times \mathbb{N} \times \mathbb{N} : |x_{mnk} - l| \geq r + \epsilon\}.$$

In this case, l is called r -statistical limit of x . If $r = 0$, then it is ordinary statistical convergence of triple sequence.

Definition 2.7. A class I of subsets of a nonempty set X is said to be an ideal in X , provided that

- (i) $\emptyset \in I$,
- (ii) $A, B \in I$ implies $A \cup B \in I$,

- (iii) $A \in I, B \subset A$ implies $B \in I$.
 I is called a nontrivial ideal, if $X \notin I$.

Definition 2.8. A nonempty class F of subsets of a nonempty set X is said to be a filter in X , provided that

- (i) $\emptyset \in F$,
(ii) $A, B \in F$ implies $A \cap B \in F$,
(iii) $A \in F, A \subset B$ implies $B \in F$.

Definition 2.9. Let I be a non trivial ideal in X and let $X \neq \emptyset$. Then the class

$$F(I) = \{M \subset X : M = X \setminus A \text{ for some } A \in I\}$$

is a filter on X , called the filter associated with I .

Definition 2.10. A non trivial ideal I in X is called admissible, if $\{x\} \in I$, for each $x \in X$.

Note 2.11. If I is an admissible ideal, then usual convergence in X implies I convergence in X .

Remark 2.12. If I is an admissible ideal, then usual rough convergence implies rough I – convergence.

Definition 2.13. Let $x = (x_{mnk})$ be a triple sequence in a metric space $(X, |.,.|)$ and r be a non-negative real number is said to be rough ideal convergent or rI – convergent to l , denoted by $x \rightarrow^{rI} l$, if for any $\epsilon > 0$ we have

$$\{(m, n, k) \in \mathbb{N} \times \mathbb{N} \times \mathbb{N} : |x_{mnk} - l| \geq r + \epsilon\} \in I.$$

In this case, l is called rI – limit of x and a triple sequence $x = (x_{mnk})$ is called rough I – convergent to l with r as roughness of degree. If $r = 0$, then it is ordinary I – convergent.

Note 2.14. Generally, a triple sequence $y = (y_{mnk})$ is not I – convergent in usual sense and $|x_{mnk} - y_{mnk}| \leq r$ for all $(m, n, k) \in \mathbb{N} \times \mathbb{N} \times \mathbb{N}$ or

$$\{(m, n, k) \in \mathbb{N} \times \mathbb{N} \times \mathbb{N} : |x_{mnk} - y_{mnk}| \geq r\} \in I,$$

for some $r > 0$. Then the triple sequence $x = (x_{mnk})$ is rI – convergent.

Note 2.15. It is clear that rI – limit of x is not necessarily unique.

Definition 2.16. Consider rI – limit set of x , which is denoted by

$$I - LIM_x^r = \{L \in X : x \rightarrow^{rI} l\},$$

then the triple sequence $x = (x_{mnk})$ is said to be rI – convergent, if $I - LIM_x^r \neq \emptyset$ and r is called a rough I – convergence degree of x .

Definition 2.17. A triple sequence $x = (x_{mnk}) \in X$ is said to be I – analytic, if there exists a positive real number M such that

$$\{(m, n, k) \in \mathbb{N} \times \mathbb{N} \times \mathbb{N} : |x_{mnk}|^{1/(m+n+k)} \geq M\} \in I.$$

Definition 2.18. A point $L \in X$ is said to be an I - accumulation point of a triple sequence $x = (x_{mnk})$ in a metric space (X, d) , if for each $\epsilon > 0$, the set

$$\{(m, n, k) \in \mathbb{N}^3 : d(x_{mnk}, l) = |x_{mnk} - l| < \epsilon\} \notin I.$$

We denote the set of all I - accumulation points of x by $I(\Gamma_x)$.

Definition 2.19. A triple sequence $x = (x_{mnk})$ is said to be Wijsman r - convergent to A denoted by $A_{mnk} \rightarrow^r A$, provided that

$$\forall \epsilon > 0 \exists (m_\epsilon, n_\epsilon, k_\epsilon) \in \mathbb{N}^3 : m \geq m_\epsilon, n \geq n_\epsilon, k \geq k_\epsilon \implies \lim_{rst} \frac{1}{rst} |\{m \leq r, n \leq s, k \leq t : |d(x, A_{mnk}) - d(x, A)| < r + \epsilon\}| = 0.$$

The set

$$LIM^r A = \{L \in \mathbb{R}^3 : A_{mnk} \rightarrow^r A\}$$

is called the Wijsman r - limit set of the triple sequences.

Definition 2.20. A triple sequence $x = (x_{mnk})$ is said to be Wijsman r - convergent if $LIM^r A \neq \emptyset$. In this case, r is called the Wijsman convergence degree of the triple sequence $x = (x_{mnk})$. For $r = 0$, we get the ordinary convergence.

Definition 2.21. A triple sequence (x_{mnk}) is said to be Wijsman r - statistically convergent to A , denoted by $A_{mnk} \rightarrow^{rst} A$, provided that the set

$$\lim_{rst} \frac{1}{rst} |\{(m, n, k) \in \mathbb{N}^3 : |d(x, A_{mnk}) - d(x, A)| \geq r + \epsilon\}| = 0$$

has natural density zero for every $\epsilon > 0$, or equivalently, if the condition

$$st - \limsup |d(x, A_{mnk}) - d(x, A)| \leq r$$

is satisfied.

In addition, we can write $A_{mnk} \rightarrow^{rst} A$ if and only if the inequality

$$\lim_{rst} \frac{1}{rst} |\{m \leq r, n \leq s, k \leq t : |d(x, A_{mnk}) - d(x, A)| < r + \epsilon\}| = 0$$

holds for every $\epsilon > 0$ and almost all (m, n, k) . Here r is called the Wijsman roughness of degree. If we take $r = 0$, then we obtain the ordinary Wijsman statistical convergence of triple sequence.

In a similar fashion to the idea of classic Wijsman rough convergence, the idea of Wijsman rough statistical convergence of a triple sequence spaces can be interpreted as follows:

Assume that a triple sequence $y = (y_{mnk})$ is Wijsman statistically convergent and cannot be measured or calculated exactly; one has to do with an approximated (or Wijsman statistically approximated) triple sequence $x = (x_{mnk})$ satisfying $|d(x - y, A_{mnk}) - d(x - y, A)| \leq r$ for all m, n, k (or for almost all (m, n, k)), i.e.,

$$\delta \left(\lim_{rst} \frac{1}{rst} |\{m \leq r, n \leq s, k \leq t : |d(x - y, A_{mnk}) - d(x - y, A)| > r\}| \right) = 0.$$

Then the triple sequence x is not statistically convergent any more, but as the inclusion

$$(2.1) \quad \lim_{rst} \frac{1}{rst} \{ |d(y, A_{mnk}) - d(y, A)| \geq \epsilon \} \supseteq \lim_{rst} \frac{1}{rst} \{ |d(x, A_{mnk}) - d(x, A)| \geq r + \epsilon \}$$

holds and we have

$$\delta \left(\lim_{rst} \frac{1}{rst} |\{ (m, n, k) \in \mathbb{N}^3 : |y_{mnk} - l| \geq \epsilon \}| \right) = 0,$$

i.e., we get

$$\delta \left(\lim_{rst} \frac{1}{rst} |\{ m \leq r, n \leq s, k \leq t : |d(x, A_{mnk}) - d(x, A)| \geq r + \epsilon \}| \right) = 0,$$

i.e., the triple sequence spaces x is Wijsman r - statistically convergent in the sense of definition 2.21.

In general, the Wijsman rough statistical limit of a triple sequence may not unique for the Wijsman roughness degree $r > 0$. So we have to consider the so called Wijsman r - statistical limit set of a triple sequence $x = (x_{mnk})$, which is defined by

$$st - LIM^r A_{mnk} = \{ L \in \mathbb{R} : A_{mnk} \xrightarrow{rst} A \}.$$

The triple sequence x is said to be Wijsman r - statistically convergent provided that $st - LIM^r A_{mnk} \neq \emptyset$. It is clear that if $st - LIM^r A_{mnk} \neq \emptyset$ for a triple sequence $x = (x_{mnk})$ of real numbers, then we have

$$(2.2) \quad st - LIM^r A_{mnk} = [st - \limsup A_{mnk} - r, st - \liminf A_{mnk} + r]$$

We know that $LIM^r = \emptyset$ for an unbounded triple sequence $x = (x_{mnk})$. But such a triple sequence might be Wijsman rough statistically convergent. For instance, define

$$d(x, A_{mnk}) = \begin{cases} (-1)^{mnk}, & \text{if } (m, n, k) \neq (i, j, \ell)^2 (i, j, \ell \in \mathbb{N}), \\ (mnk), & \text{otherwise} \end{cases}.$$

in \mathbb{R} . Because the set $\{1, 64, 739, \dots\}$ has natural density zero, we have

$$st - LIM^r A_{mnk} = \begin{cases} \phi, & \text{if } r < 1, \\ [1 - r, r - 1], & \text{otherwise} \end{cases}$$

and $LIM^r A_{mnk} = \phi$ for all $r \geq 0$.

As can be seen by the example above, the fact that $st - LIM^r A_{mnk} \neq \emptyset$ does not imply $LIM^r A_{mnk} \neq \emptyset$. Because a finite set of natural numbers has natural density zero, $LIM^r A_{mnk} \neq \emptyset$ implies $st - LIM^r A_{mnk} \neq \emptyset$. Therefore, we get $LIM^r A_{mnk} \subseteq st - LIM^r A_{mnk}$. This obvious fact means

$$\{r \geq 0 : LIM^r A_{mnk} \neq \emptyset\} \subseteq \{r \geq 0 : st - LIM^r A_{mnk} \neq \emptyset\}$$

in this language of sets and yields immediately

$$\inf \{r \geq 0 : LIM^r A_{mnk} \neq \emptyset\} \geq \inf \{r \geq 0 : st - LIM^r A_{mnk} \neq \emptyset\}.$$

Moreover, it also yields directly $diam(LIM^r A_{mnk}) \leq diam(st - LIM^r A_{mnk})$.

Throughout the paper, we let $(X; \rho)$ be a separable metric space, $I \subseteq 2^{\mathbb{N}^3}$ be an admissible ideal and $A; A_{mnk}$ be any non-empty closed subsets of X .

Definition 2.22. A triple sequence (x_{mnk}) is said to be Wijsman $r - I$ convergent to A , if for every $\epsilon > 0$ and for each $x \in X$,

$$A(x, \epsilon) = \{(m, n, k) \in \mathbb{N}^3 : |d(x, A_{mnk}) - d(x, A)| \geq r + \epsilon\} \in I$$

Definition 2.23. A triple sequence (x_{mnk}) is said to be Wijsman $r - I$ statistical convergent to A , if for every $\epsilon > 0, \delta > 0$ and for each $x \in X$,

$$\left\{ (r, s, t) \in \mathbb{N}^3 : \frac{1}{rst} |\{(r, s, t) \leq (m, n, k) : |d(x, A_{mnk}) - d(x, A)| \geq r + \epsilon\}| \geq \delta \right\} \in I.$$

In this case, we write $A_{mnk} \rightarrow^{s(Iw)} A$.

Definition 2.24. Let θ be a lacunary sequence. A triple sequence (x_{mnk}) is said to be Wijsman strongly $r - I$ convergent to A , if for every $\epsilon > 0$ and for each $x \in X$,

$$\left\{ (r, s, t) \in \mathbb{N}^3 : \frac{1}{h_{rst}} \sum_{(m,n,k) \in I_{rst}} |d(x, A_{mnk}) - d(x, A)| \geq r + \epsilon \right\} \in I.$$

In this case, we write $A_{mnk} \rightarrow^{N_\theta(Iw)} A$.

3. MAIN RESULTS

Theorem 3.1. If $I - LIM^r |d(x, A_{mnk}) - d(x, A)| \neq \emptyset$ for a triple sequence $|d(x, A_{mnk}) - d(x, A)|$ of real numbers, and

$$I - LIM^r |d(x, A_{mnk}) - d(x, A)| = [I - \limsup |d(x, A_{mnk}) - d(x, A)| - r, I - \liminf |d(x, A_{mnk}) - d(x, A)| + r]$$

then

$$\text{diam}(LIM^r |d(x, A_{mnk}) - d(x, A)|) \leq \text{diam}(I - LIM^r |d(x, A_{mnk}) - d(x, A)|).$$

Proof. We know that $I - LIM^r |d(x, A_{mnk}) - d(x, A)| = \emptyset$ for an unbounded triple sequence $|d(x, A_{mnk}) - d(x, A)|$. But such a sequence might be Wijsman rough I -convergent. For instance, let I be the I_d of \mathbb{N} and define

$$|d(x, A_{mnk}) - d(x, A)| = \begin{cases} \cos(mnk)\pi, & \text{if } (m, n, k) \neq (ij\ell)^2 (i, j, \ell \in \mathbb{N}), \\ (mnk), & \text{otherwise} \end{cases},$$

in \mathbb{R}^3 . Because the set $\{1, 64, 739, \dots\}$ belong to I , we have

$$I - LIM^r |d(x, A_{mnk}) - d(x, A)| = \begin{cases} \phi, & \text{if } r < 1, \\ [1 - r, r - 1], & \text{otherwise} \end{cases},$$

and $LIM^r |d(x, A_{mnk}) - d(x, A)| = \emptyset$, for all $r \geq 0$. The fact that $I - LIM^r |d(x, A_{mnk}) - d(x, A)| \neq \emptyset$ does not imply $LIM^r |d(x, A_{mnk}) - d(x, A)| \neq \emptyset$. Because I is a admissible ideal

$$LIM^r |d(x, A_{mnk}) - d(x, A)| \neq \emptyset \implies I - LIM^r |d(x, A_{mnk}) - d(x, A)| \neq \emptyset,$$

i.e., if $x = (x_{mnk}) \in LIM^r |d(x, A_{mnk}) - d(x, A)|, (x_{mnk}) \in I - LIM^r |d(x, A_{mnk}) - d(x, A)|$, for each triple sequences. Also, if we define all the Wijsman rough convergence sequences by LIM^r and rough I -convergence sequences by $I - LIM^r$, then we get

$$LIM^r \subseteq I - LIM^r.$$

$$\begin{aligned} \{r \geq 0 : LIM^r |d(x, A_{mnk}) - d(x, A)| \neq \phi\} \subseteq \\ \{r \geq 0 : I - LIM^r |d(x, A_{mnk}) - d(x, A)| \neq \phi\}. \end{aligned}$$

Hence the sets yields immediately

$$\begin{aligned} \inf \{r \geq 0 : LIM^r |d(x, A_{mnk}) - d(x, A)| \neq \phi\} \geq \\ \{r \geq 0 : I - LIM^r |d(x, A_{mnk}) - d(x, A)| \neq \phi\}, \end{aligned}$$

for each triple sequences. Moreover, it also yield directly

$$diam(LIM^r |d(x, A_{mnk}) - d(x, A)|) \leq diam(I - LIM^r |d(x, A_{mnk}) - d(x, A)|).$$

□

Note: The Wijsman rough I - limit of a triple sequence is unique for the roughness degree $r > 0$.

We state the following result without proof, since it can be established using standard technique.

Theorem 3.2. *If $I \subset 3^{\mathbb{N}}$ be an Wijsman rough strongly admissible ideal and $|d(x, A_{mnk}) - d(x, A)|$ be a triple sequence then we have $I(\Lambda_{|d(x, A_{mnk}) - d(x, A)|}) \subseteq I(\Gamma_{|d(x, A_{mnk}) - d(x, A)|})$.*

Proof. Let $c \in I(\Lambda_{|d(x, A_{mnk}) - d(x, A)|})$. If $c \notin LIM^r |d(x, A_{mnk}) - d(x, A)|$ then there exists a set $M = \{(m, n, k) \in \mathbb{N}^3 : (u_m v_n w_k)\} \notin I$, such that

$$(3.1) \quad \{(m, n, k) \in \mathbb{N}^3 : |d(x_{u_m v_n w_k}, A_{mnk}) - d(x_{u_m v_n w_k}, A)|, \xi|^{1/(m+n+k)} \geq r + \epsilon\} \notin I.$$

Let $\epsilon > 0$. Then by (3.1) there exists $(r_0 s_0 t_0) \in \mathbb{N}$ such that $u_m \geq r_0, v_n \geq s_0, w_k \geq t_0$, we have

$$\begin{aligned} \{(m, n, k) \in \mathbb{N}^3 : |d(x_{u_m v_n w_k}, A_{mnk}) - d(x_{u_m v_n w_k}, A)|, \xi|^{1/(m+n+k)} \geq r + \epsilon\} \supset \\ M \setminus \{(m, n, k) \in \mathbb{N}^3 : (u_m v_n w_k), \text{ either } u_m \leq (r_0 - 1) \text{ or } v_m \leq (s_0 - 1) \text{ or } w_m \leq (t_0 - 1)\}. \end{aligned}$$

Since I is Wijsman rough strongly admissible, so

$$\{(m, n, k) \in \mathbb{N}^3 : |d(x, A_{mnk}) - d(x, A)|, c| \geq r + \epsilon\} \notin I.$$

This implies $c \in I(\Gamma_{|d(x, A_{mnk}) - d(x, A)|})$. Hence

$$I(\Lambda_{|d(x, A_{mnk}) - d(x, A)|}) \subseteq I(\Gamma_{|d(x, A_{mnk}) - d(x, A)|}).$$

□

Theorem 3.3. *If $I \subset 3^{\mathbb{N}}$ be an Wijsman rough strongly admissible ideal, $I - LIM^r |d(x, A_{mnk}) - d(x, A)| \neq \emptyset$ for a triple sequence $x = (x_{mnk})$ of real numbers and (i) $I - \limsup |d(x, A_{mnk}) - d(x, A)| = \alpha$ if and only if for any $\epsilon > 0$,*

$$\{(m, n, k) \in \mathbb{N}^3 : |d(x, A_{mnk}) - d(x, A)|, \alpha| \geq r + \epsilon\} \notin I$$

and

$$\{(m, n, k) \in \mathbb{N}^3 : |d(x, A_{mnk}) - d(x, A)|, \alpha| \geq \epsilon\} \in I.$$

(ii) $I - \liminf |d(x, A_{mnk}) - d(x, A)| = \beta$ if and only if for any $\epsilon > 0$,

$$\{(m, n, k) \in \mathbb{N}^3 : |d(x, A_{mnk}) - d(x, A)|, \beta \leq r + \epsilon\} \notin I$$

and

$$\{(m, n, k) \in \mathbb{N}^3 : |d(x, A_{mnk}) - d(x, A)|, \beta \leq \epsilon\} \in I.$$

Theorem 3.4. If $I \subset 3^{\mathbb{N}}$ be an Wijsman rough strongly admissible ideal, $I - LIM^r |d(x, A_{mnk}) - d(x, A)| \neq \emptyset$ for a triple sequence $x = (x_{mnk})$ of real numbers then

$$I - \liminf |d(x, A_{mnk}) - d(x, A)| \leq I - \limsup |d(x, A_{mnk}) - d(x, A)|$$

holds.

Proof. It can be established using the techniques of Theorem 3 ([15]), so proof is omitted. \square

Theorem 3.5. If $I \subset 3^{\mathbb{N}}$ be an Wijsman rough strongly admissible ideal, $I - LIM^r |d(x, A_{mnk}) - d(x, A)| \neq \emptyset$ for a triple sequence $x = (x_{mnk})$ of real numbers then

$$\begin{aligned} I - LIM^r |d(x, A_{mnk}) - d(x, A)| &\leq I - \liminf |d(x, A_{mnk}) - d(x, A)| + r \\ &\leq I - \limsup |d(x, A_{mnk}) - d(x, A)| + r \leq I - LIM^r |d(x, A_{mnk}) - d(x, A)|. \end{aligned}$$

Proof. We first prove that

$$\begin{aligned} I - LIM^r |d(x, A_{mnk}) - d(x, A)| - \liminf |d(x, A_{mnk}) - d(x, A)| \\ \leq I - \limsup |d(x, A_{mnk}) - d(x, A)| + r. \end{aligned}$$

If $I - LIM^r |d(x, A_{mnk}) - d(x, A)| - \liminf |d(x, A_{mnk}) - d(x, A)| = -\infty$, then it is obvious. Let $I - LIM^r |d(x, A_{mnk}) - d(x, A)| - \liminf = r > -\infty$. Then

$$r_1 = \sup_{uvw} r$$

where $r = \inf \{m \geq u, n \geq v, k \geq w : |d(x, A_{mnk}) - d(x, A)|\}$. Then

$$\begin{aligned} \{(m, n, k) \in \mathbb{N}^3 : |d(x, A_{mnk}) - d(x, A)| < r_1\} \subset \\ \{(m, n, k) \in \mathbb{N}^3 : (m, n, k), \text{ either } m \leq (u-1) \text{ or } n \leq (v-1) \text{ or } k \leq (w-1)\}. \end{aligned}$$

The fact that $I - LIM^r |d(x, A_{mnk}) - d(x, A)| \neq \emptyset$ does not imply $LIM^r |d(x, A_{mnk}) - d(x, A)| \neq \emptyset$. Since I is strongly admissible ideal

$$LIM^r |d(x, A_{mnk}) - d(x, A)| \neq \emptyset \implies I - LIM^r |d(x, A_{mnk}) - d(x, A)| \neq \emptyset,$$

we have

$$\{(m, n, k) \in \mathbb{N}^3 : (m, n, k), \text{ either } m \leq (u-1) \text{ or } n \leq (v-1) \text{ or } k \leq (w-1)\} \in I,$$

so, then there exists $\epsilon > 0$ such that

$$\{(m, n, k) \in \mathbb{N}^3 : |d(x, A_{mnk}) - d(x, A)| < r_1 + \epsilon\} \in I.$$

Now, let $r_2 = I - \liminf |d(x, A_{mnk}) - d(x, A)| = \inf A_1(\epsilon)$. Since $a \in I - LIM^r |d(x, A_{mnk}) - d(x, A)|$, we have $A_1(\epsilon) \in I$ for every $\epsilon > 0$, where

$$A_1(\epsilon) = \{a \in \mathbb{R} : \{(m, n, k) \in \mathbb{N}^3 : |d(x, A_{mnk}) - d(x, A)|, a \geq a + \epsilon\} \notin I\}.$$

Now if $r_2 < r_1$, then there exists $a' \in A_1(\epsilon)$ such that $r_2 \leq a' < r_1$, we have,

$$\begin{aligned} \{r \geq 0 : LIM^r |d(x, A_{mnk}) - d(x, A)| \neq \phi\} \subseteq \\ \{r \geq 0 : I - LIM^r |d(x, A_{mnk}) - d(x, A)| \neq \emptyset\}. \end{aligned}$$

Hence the sets yields immediately

$$\begin{aligned} \inf \{r \geq 0 : LIM^r |d(x, A_{mnk}) - d(x, A)| \neq \phi\} \geq \\ \{r \geq 0 : I - LIM^r |d(x, A_{mnk}) - d(x, A)| \neq \emptyset\}. \end{aligned}$$

for each triple sequences. However,

$$\begin{aligned} \{(m, n, k) \in \mathbb{N}^3 : |d(x, A_{mnk}) - d(x, A)| < a' + \epsilon\} \subset \\ \{(m, n, k) \in \mathbb{N}^3 : |d(x, A_{mnk}) - d(x, A)| < r_1 + \epsilon\} \in I \end{aligned}$$

which yields $a' \notin A_1(\epsilon)$, which is a contradiction. Then $r_2 \geq r_1$ for all (u, v, w) . Hence it also yield directly $r_1 \leq r_2$, i.e.,

$$I - LIM^r |d(x, A_{mnk}) - d(x, A)| - \liminf |d(x, A_{mnk}) - d(x, A)| \leq I - \liminf x.$$

Similarly we can show

$$I - \limsup |d(x, A_{mnk}) - d(x, A)| \leq I - LIM^r x - \limsup |d(x, A_{mnk}) - d(x, A)|.$$

□

Competing Interests: The authors declare that there is not any conflict of interests regarding the publication of this manuscript.

REFERENCES

- [1] M. Aiyub, A. Esi and N. Subramanian, The triple entire difference ideal of fuzzy real numbers over fuzzy p - metric spaces defined by Musielak Orlicz function, J. Intell. Fuzzy Systems 33 (2017) 1505–1512.
- [2] S. Aytar, Rough statistical convergence, Numer. Funct. Anal. Optim. 29 (3-4) (2008) 291–303.
- [3] S. Debnath, B. Sarma and B. C. Das, Some generalized triple sequence spaces of real numbers, J. Nonlinear Anal. Optim. 6 (1) (2015) 71–78.
- [4] A. J. Dutta A. Esi and B. C. Tripathy, Statistically convergent triple sequence spaces defined by Orlicz function, J. Math. Anal. 4 (2) (2013) 16–22.
- [5] A. Esi, On some triple almost lacunary sequence spaces defined by Orlicz functions, Research and Reviews: Discrete Mathematical Structures 1 (2) (2014) 16–25.
- [6] A. Esi and M. Necdet Catalbas, Almost convergence of triple sequences, Global Journal of Mathematical Analysis 2 (1) (2014) 6–10.
- [7] A. Esi and E. Savas, On lacunary statistically convergent triple sequences in probabilistic normed space, Appl. Math. Inf. Sci. 9 (5) (2015) 2529–2534.
- [8] H. Fast, Sur la convergence statistique, Colloq. Math. 2 (1951) 241–244.
- [9] S. K. Pal, D. Chandra and S. Dutta, Rough ideal convergence, Hacet. J. Math. Stat. 42 (6) (2013) 633–640.
- [10] H. X. Phu, Rough convergence in normed linear spaces, Numer. Funct. Anal. Optim. 22 (1-2) (2001) 199–222.
- [11] A. Sahiner, M. Gurdal and F. K. Duden, Triple sequences and their statistical convergence, Selçuk J. Appl. Math., 8 (2) (2007) 49–55.
- [12] A. Sahiner, B. C. Tripathy, Some I -related properties of triple sequences, Selçuk J. Appl. Math. 9 (2) (2008) 9–18.
- [13] H. Steinhaus, Sur la convergence ordinaire et la convergence asymptotique, Colloq. Math. 2 (1951) 73–74.

- [14] N. Subramanian and A. Esi, The generalized tripled difference of χ^3 sequence spaces, Global Journal of Mathematical Analysis 3 (2) (2015) 54–60.
- [15] B. C. Tripathy and R. Goswami, Vector valued multiple sequences defined by Orlicz functions, Bol. Soc. Parana. Mat. 33 (1) (2015) 67–79.
- [16] B. C. Tripathy and R. Goswami, Multiple sequences in probabilistic normed spaces, Afr. Mat. 26 (5-6) (2015) 753–760.
- [17] B. C. Tripathy and R. Goswami, Fuzzy real valued P -absolutely summable multiple sequences in probabilistic normed spaces, Afr. Mat. 26 (7-8) (2015) 1281–1289.
- [18] B. C. Tripathy and R. Goswami, Statistically convergent multiple sequences in probabilistic normed spaces, Politehn. Univ. Bucharest Sci. Bull. Ser. A Appl. Math. Phys. 78 (4) (2016) 83–94.

AYHAN ESI (aesi23@hotmail.com)

Department of Mathematics, Adiyaman University, 02040, Adiyaman, Turkey

NAGARAJAN SUBRAMANIAN (nsmaths@yahoo.com)

Department of Mathematics, SASTRA University, Thanjavur-613 401, India

AYTEN ESI (aytenesi@yahoo.com)

Department of Mathematics, Adiyaman University, 02040, Adiyaman, Turkey