Annals of Fuzzy Mathematics and Informatics Volume 21, No. 1, (February 2021) pp. 29–37 ISSN: 2093–9310 (print version) ISSN: 2287–6235 (electronic version) http://www.afmi.or.kr https://doi.org/10.30948/afmi.2021.21.1.29



© Research Institute for Basic Science, Wonkwang University http://ribs.wonkwang.ac.kr

Concept lattice on cubic sets

@FMI QFMI **◎ F** M I 0 0 0 TF I @ F M I @ F M I ©ℾМӏ © **⊮** M I @ **⊮** M I © **⊮** M I $\odot \mathbb{F} \mathbb{M} \mathbb{I}$ © **F** M Ⅰ $\textcircled{0} \mathbb{F} \mathbb{M} \mathbb{I}$ $\textcircled{0} \mathbb{F} \mathbb{M} \mathbb{I}$ $\textcircled{\ }\mathbb{F}\mathbb{M}\mathbb{I}$ $\textcircled{0} \mathbb{F} \mathbb{M} \mathbb{I}$ $@ \mathbb{F} \mathbb{M} \mathbb{I}$ $@ \mathbb{F} \mathbb{M} \mathbb{I}$ $@ \mathbb{F} \mathbb{M}$ $@ \mathbb{F} \mathbb{M}$ T 00 \mathbb{F} 0 \mathbb{F} 0 \mathbb{F} \mathbb{N} IVI \mathbb{F} $(\mathbf{0})$ \mathbb{M}



XUEYOU CHEN

Annals of Fuzzy Mathematics and Informatics Volume 21, No. 1, (February 2021) pp. 29–37 ISSN: 2093–9310 (print version) ISSN: 2287–6235 (electronic version) http://www.afmi.or.kr https://doi.org/10.30948/afmi.2021.21.1.29



© Research Institute for Basic Science, Wonkwang University http://ribs.wonkwang.ac.kr

Concept lattice on cubic sets

XUEYOU CHEN

Received 27 August 2020; Revised 16 September 2020; Accepted 28 September 2020

ABSTRACT. In 2012, Jun et al. introduced the notion of a cubic set, which is a generalization of Zadeh's fuzzy set. In 1999, Ganter and Wille put forward formal concept analysis theory(FCA), formal context, concept lattice are two main notions and tools. In the paper, we define formal context, Galois connection, concept lattice on cubic sets, and generalize formal concept analysis theory on cubic sets structure.

2020 AMS Classification: 03E72, 06A15, 08A72

Keywords: Galios connection, Concept lattice, Cubic set.

Corresponding Author: Xueyou Chen (math-chen@qq.com)

1. INTRODUCTION

In 1965, Zadeh [1] initiated fuzzy sets. As a generalization, in 2012, Jun et al. [2] introduced the notion of a cubic set. After then, Kang and Kim [3] defined a mapping of cubic sets, Kim et al. [4] investigated a cubic relation between cubic sets. The related contents may be referred [5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18].

Ganter and Wille [19] put forward formal concept analysis theory(FCA), which is an order-theoretical analysis of scientific data. Formal context, concept lattice are two main notions and tools. In [20], from the point of view of graded approach, Bělohlávek established **L**-order, **L**-context and **L**-concept lattice. Djouadi and Prade [21] investigated the interval-valued formal concept analysis and Brito et al. [22] considered fuzzy formal concept analysis. We also discussed approximate concept lattice in fuzzy setting and the relation to fuzzy rough sets [23, 24]. More references see International Conference Homepages: Concept Lattices and Their Applications.

In the paper, we introduce the notions of formal context, concept lattice on cubic sets, combine formal concept analysis and cubic sets together. The contents are arranged into two parts, Section 3: Concept lattice on Cubic sets. In Section 2, we give an overview of concept lattice and cubic sets, which surveys Preliminaries

2. Preliminaries

The section is devoted to some main notions for each area, i.e., formal concept analysis [19, 20] fuzzy sets [1], interval-valued fuzzy sets [25, 26] and cubic sets [2, 4].

2.1. Formal Concept Analysis. FCA is an order-theoretic method for the mathematical analysis of scientific data, pioneered by Wille and others [19], has attracted a growing number of researchers and practitioners. We introduce by formalizing the notion of (formal) context.

A context is a triple (G, M, I) consisting of two sets G and M and a relation I between them. The elements of G are called the *objects* and the *elements* of M are called the attributes. We write gIm or $(g, m) \in I$ to show that the object g has the attribute m.

For a set $A \subseteq G$ of objects define $A' = \{m \in M \mid gIm \text{ for all } g \in A\}$. Correspondingly, for a set $B \subseteq M$ of attributes define $B' = \{g \in G \mid gIm \text{ for all } m \in B\}$. A concept of the context (G, M, I) is a pair (A, B) where $A \subseteq G, B \subseteq M, A' = B$ and B' = A. We call A the *extent* and B the *intent* of the concept (A, B).

Suppose (A_1, B_1) , (A_2, B_2) are two concepts of the context (G, M, I), we define an order \leq , where $(A_1, B_1) \leq (A_2, B_2)$ if $A_1 \subseteq A_2$ (which is equivalent to $B_1 \supseteq B_2$).

Let $\mathcal{C}(G, M, I, \leq)$ be the set of all concepts of the context (G, M, I) with the order \leq . It forms a complete lattice (Concept Lattice) in which join and meet are given by:

$$\bigvee_{j \in J} (A_j, B_j) = ((\bigcup_{j \in J} A_j)'', \bigcap_{j \in J} B_j), \quad \bigwedge_{j \in J} (A_j, B_j) = (\bigcap_{j \in J} A_j, (\bigcup_{j \in J} B_j)'').$$

2.2. **Fuzzy sets.** We introduce some basic definitions about fuzzy sets, fuzzy relations [1].

For a set X, a mapping $\lambda : X \to I$ is called a *fuzzy set* in X, where I = [0, 1]. The collection of all fuzzy sets in X is denoted by I^X . In particular, 0 and 1 denote the fuzzy empty set and the fuzzy whole set in X, respectively.

For any λ , $\mu \in I^X$, the join (\vee) and meet (\wedge) of λ and μ , denoted by $\lambda \vee \mu$ and $\lambda \wedge \mu$, are defined as follows: for each $x \in X$,

$$(\lambda \lor \mu)(x) = \max\{\lambda(x), \mu(x)\}, \qquad (\lambda \land \mu)(x) = \min\{\lambda(x), \mu(x)\}.$$

For any family $(\lambda_j)_{j \in J}$ of fuzzy sets in X, the join (\vee) and meet (\wedge) of $(\lambda_j)_{j \in J}$, denoted by $\bigvee_{j \in J} \lambda_j$ and $\bigwedge_{j \in J} \lambda_j$, are defined as follows: for each $x \in X$,

$$(\bigvee_{j\in J}\lambda_j)(x) = \sup_{j\in J}\lambda_j(x) \qquad (\bigwedge_{j\in J}\lambda_j)(x) = \inf_{j\in J}\lambda_j(x).$$

For two sets X, Y, $r \in I^{X \times Y}$ is called a *fuzzy relation from* X to Y.

2.3. Interval-valued fuzzy sets. In the section, we introduce some basic definitions related to interval-valued fuzzy sets and interval-valued fuzzy relations (See [25, 26, 27]).

The set of all closed subintervals of I is denoted by [I], and members of [I] are called *interval numbers* and are denoted by \tilde{a} , \tilde{b} , \tilde{c} , etc., where $\tilde{a} = [a^-, a^+]$ and $0 \le a^- \le a^+ \le 1$. In particular, if $a^- = a^+$, then we write as $\tilde{a} = \mathbf{a}$.

We define relations \leq and = on [I] as follows:

$$(\forall \ \widetilde{a}, \ \widetilde{b} \in [I])(\widetilde{a} \preceq \widetilde{b} \Longleftrightarrow a^{-} \leq b^{-} \text{ and } a^{+} \leq b^{+}),$$
$$(\forall \ \widetilde{a}, \ \widetilde{b} \in [I])(\widetilde{a} = \widetilde{b} \Longleftrightarrow \widetilde{a} \succeq \widetilde{b} \text{ and } \widetilde{a} \preceq \widetilde{b}), i.e.,$$
$$(\forall \ \widetilde{a}, \ \widetilde{b} \in [I])(\widetilde{a} = \widetilde{b} \Longleftrightarrow a^{-} = b^{-} \text{ and } a^{+} = b^{+}).$$

To say $\widetilde{a} < \widetilde{b}$, we mean $\widetilde{a} \preceq \widetilde{b}$ and $\widetilde{a} \neq \widetilde{b}$.

For any $\tilde{a}, \tilde{b} \in [I]$, their minimum and maximum, denoted by $\tilde{a} \wedge \tilde{b}$ and $\tilde{a} \vee \tilde{b}$ are defined as follows:

$$\widetilde{a} \wedge \widetilde{b} = [a^- \wedge b^-, a^+ \wedge b^+], \qquad \widetilde{a} \vee \widetilde{b} = [a^- \vee b^-, a^+ \vee b^+].$$

Let $(\tilde{a}_j)_{j\in J} \subset [I]$. Then its inf and sup, denoted by $\bigwedge_{j\in J} \tilde{a}_j$ and $\bigvee_{j\in J} \tilde{a}_j$, are defined as follows:

$$\bigwedge_{j\in J} \widetilde{a}_j = [\bigwedge_{j\in J} a_j^-, \bigwedge_{j\in J} a_j^+], \qquad \bigvee_{j\in J} \widetilde{a}_j = [\bigvee_{j\in J} a_j^-, \bigvee_{j\in J} a_j^+].$$

For a nonempty set X, a mapping $A: X \to [I]$ is called an *interval-valued fuzzy* set (briefly, an IVF set) in X. Let $[I]^X$ denote the set of all IVF sets in X. For each $A \in [I]^X$ and $x \in X$, $A(x) = [A^-(x), A^+(x)]$ is called the *degree of membership* of an element x to A, where $A^-, A^+ \in I^X$ are called a *lower fuzzy set* and an *upper fuzzy* set in X, respectively. For each $A \in [I]^X$, we write $A = [A^-, A^+]$. In particular, $\widetilde{0}$ and $\widetilde{1}$ denote the interval-valued fuzzy empty set and the interval-valued fuzzy whole set in X, respectively. We define relations \subset and = on $[I]^X$ as follows:

$$(\forall A, B \in [I]^X)(A \subset B \iff (x \in X)(A(x) \preceq B(x)),$$
$$(\forall A, B \in [I]^X)(A = B \iff (x \in X)(A(x) = B(x)).$$

For any $(A_j)_{j\in J} \subset [I]^X$, its intersection $\bigcap_{j\in J} A_j$ and union $\bigcup_{j\in J} A_j$ are defined, respectively as follows: for each $x \in X$,

$$(\bigcap_{j\in J} A_j)(x) = \bigwedge_{j\in J} A_j(x), \qquad (\bigcup_{j\in J} A_j)(x) = \bigvee_{j\in J} A_j(x).$$

For two sets X, Y, $R \in [I]^{X \times Y}$ is called an *interval-valued fuzzy relation* (briefly, IVF relation) from X to Y.

2.4. Cubic sets and cubic relations. As a generalization, the notions of a cubic fuzzy set and a cubic fuzzy relation are introduced in [2, 4].

Definition 2.1 ([2]). Let X be a nonempty set. Then a complex mapping $\mathcal{A} = \langle A, \lambda \rangle \colon X \to [I] \times I$ is called a *cubic set* in X.

In special, a cubic set $\mathcal{A} = \langle A, \lambda \rangle$ in which $A(x) = \mathbf{0}$ and $\lambda(x) = 0$ (resp. $A(x) = \mathbf{1}$ and $\lambda(x) = 1$) for each $x \in X$ is denoted by $\hat{0}$ (resp. $\hat{1}$). In this case, $\hat{0}$ (resp. $\hat{1}$) will be called a cubic empty (resp. whole) set in X.

We denote the set of all cubic sets in X as $([I] \times I)^X$.

Definition 2.2 ([2]). Let $\mathcal{A} = \langle A, \lambda \rangle$, $\mathcal{B} = \langle B, \mu \rangle \in ([I] \times I)^X$. Then we define the following relations:

- (i) (Equality) $\mathcal{A} = \mathcal{B} \Leftrightarrow \mathcal{A} = B$ and $\lambda = \mu$,
- (ii) (P-order) $\mathcal{A} \sqsubset \mathcal{B} \Leftrightarrow \mathcal{A} \subset \mathcal{B} \text{ and } \lambda \leq \mu$,

(iii) (R-order) $\mathcal{A} \Subset \mathcal{B} \Leftrightarrow \mathcal{A} \subset \mathcal{B}$ and $\lambda \geq \mu$

Definition 2.3 ([2]). Let $\mathcal{A} = \langle A, \lambda \rangle$, $\mathcal{B} = \langle B, \mu \rangle \in ([I] \times I)^X$ and let $(\mathcal{A}_j)_{j \in J} = (\langle A_j, \lambda_j \rangle)_{j \in J} \subset ([I] \times I)^X$. Then the complement \mathcal{A}^c of \mathcal{A} , P-union \sqcup , P-intersection \sqcap , R-union \Downarrow and R-intersection \Cap are defined as follows, respectively: for each $x \in X$,

- (i) (Complement) $\mathcal{A}^{c}(x) = \langle A^{c}(x), \lambda^{c}(x) \rangle$,
- (ii) (P-union) $(\mathcal{A} \sqcup \mathcal{B})(x) = \langle (A \cup B)(x), (\lambda \lor \mu)(x) \rangle$,
 - $(\sqcup_{j\in J}\mathcal{A}_j)(x) = < (\bigcup_{j\in J}A_j)(x), (\bigvee_{j\in J}\lambda_j)(x) >,$
- (iii) (P-intersection) $(\mathcal{A} \sqcap \mathcal{B})(x) = \langle (A \cap B)(x), (\lambda \land \mu)(x) \rangle$,
- $(\bigcap_{j\in J}\mathcal{A}_{j})(x) = < (\bigcap_{j\in J}A_{j})(x), (\bigwedge_{j\in J}\lambda_{j})(x) >,$ (iv) (R-union) $(\mathcal{A} \sqcup \mathcal{B})(x) = < (\mathcal{A} \cup \mathcal{B})(x), (\lambda \land \mu)(x) >,$ $(\amalg_{j\in J}\mathcal{A}_{j})(x) = < (\bigcup_{j\in J}A_{j})(x), (\bigwedge_{j\in J}\lambda_{j})(x) >,$ (v) (R-intersection) $(\mathcal{A} \cap \mathcal{B})(x) = < (\mathcal{A} \cap \mathcal{B})(x), (\lambda \lor \mu)(x) >,$

$$(\bigcap_{j \in J} \mathcal{A}_j)(x) = < (\bigcap_{j \in J} \mathcal{A}_j)(x), (\bigvee_{j \in J} \lambda_j)(x) >$$

In special, for cubic sets $\langle \tilde{a}, \lambda \rangle, \langle \tilde{b}, \mu \rangle, \langle \tilde{a}_j, \lambda_j \rangle, \langle \tilde{b}_j, \mu_j \rangle$ $(j \in J)$, we also adopt the above symbols:

$$\begin{split} &< \widetilde{a}, \lambda > \Box < \widetilde{b}, \mu > \Leftrightarrow \widetilde{a} \preceq \widetilde{b} \text{ and } \lambda \leq \mu, \\ &< \widetilde{a}, \lambda > \Box < \widetilde{b}, \mu > = < \widetilde{a} \lor \widetilde{b}, \lambda \lor \mu >, \\ &\subset \widetilde{a}, \lambda > \Box < \widetilde{b}, \mu > = < \widetilde{a} \lor \widetilde{b}, \lambda \lor \mu >, \\ &\sqcup_{j \in J} < \widetilde{a}_j, \lambda_j > = < \bigvee_{j \in J} \widetilde{a}_j, \bigvee_{j \in J} \lambda_j >, \\ &< \widetilde{a}, \lambda > \Box < \widetilde{b}, \mu > = < \widetilde{a} \land \widetilde{b}, \lambda \land \mu >, \\ &\Box_{j \in J} < \widetilde{a}_j, \lambda_j > = < \bigvee_{j \in J} \widetilde{a}_j, \bigvee_{j \in J} \lambda_j >, \\ &< \widetilde{a}, \lambda > \Subset < \widetilde{b}, \mu > \Leftrightarrow \widetilde{a} \preceq \widetilde{b} \text{ and } \lambda \geq \mu. \end{split}$$

Definition 2.4 ([4]). Let $\mathcal{A} = \langle A, \lambda \rangle \in ([I] \times I)^X$, let $\tilde{a} \in [I]$ with $a^+ > 0$ and let $\alpha \in I$ with $\alpha > 0$. Then $\mathcal{A} = \langle A, \lambda \rangle$ is called a *cubic point in* X with the support $x \in X$ and the value $\langle \tilde{a}, \alpha \rangle$, denoted by $x_{\langle \tilde{a}, \alpha \rangle}$, if for each $y \in X$,

$$x_{<\tilde{a},\alpha>} = \begin{cases} <\tilde{a},\alpha> & \text{if } y=x\\ <\mathbf{0},0> & \text{otherwise} \end{cases}$$

The set of all cubic points in X is denoted by M_X .

Definition 2.5 ([4]). Let $x_{<\tilde{a},\alpha>} \in M_X$ and let $\mathcal{A} = \langle A, \lambda \rangle \in ([I] \times I)^X$.

- (i) $x_{<\tilde{a},\alpha>}$ is said to belong to \mathcal{A} by P-order type, denoted by $x_{<\tilde{a},\alpha>} \in_P \mathcal{A}$, if $\tilde{a} \leq A(x)$ and $\alpha \leq \lambda(x)$, i.e., $x_{\tilde{a}} \in A$ and $x_{\alpha} \in \lambda$.
- (ii) $x_{<\tilde{a},\alpha>}$ is said to belong to \mathcal{A} by R-order type, denoted by $x_{<\tilde{a},\alpha>} \in_R \mathcal{A}$, if $\tilde{a} \leq A(x)$ and $\alpha \geq \lambda(x)$, i.e., $x_{\tilde{a}} \in A$ and $x_{1-\alpha} \in \lambda^c$.

Proposition 2.6 ([4]). Let $x_{\langle \tilde{a}, \alpha \rangle} \in M_X$ and let $\mathcal{A} = \langle A, \lambda \rangle$, $\mathcal{B} = \langle B, \mu \rangle \in ([I] \times I)^X$.

- (1) $\mathcal{A} \sqsubset \mathcal{B}$ if and only if $x_{\langle \tilde{a}, \alpha \rangle} \in_P \mathcal{B}$, for each $x_{\langle \tilde{a}, \alpha \rangle} \in_P \mathcal{A}$.
- (2) $\mathcal{A} \subseteq \mathcal{B}$ if and only if $x_{\langle \tilde{a}, \alpha \rangle} \in_R \mathcal{B}$, for each $x_{\langle \tilde{a}, \alpha \rangle} \in_R \mathcal{A}$.

Proposition 2.7 ([4]). Let $\mathcal{A} = \langle A, \lambda \rangle$, $\mathcal{B} = \langle B, \mu \rangle \in ([I] \times I)^X$, let $(\mathcal{A}_j)_{j \in J} = (\langle A_j, \lambda_j \rangle)_{j \in J} \subset ([I] \times I)^X$ and let $x_{\langle \tilde{a}, \alpha \rangle} \in M_X$.

- (1) If $x_{\langle \tilde{a}, \alpha \rangle} \in_P \mathcal{A}$ or $x_{\langle \tilde{a}, \alpha \rangle} \in_P \mathcal{B}$, then $x_{\langle \tilde{a}, \alpha \rangle} \in_P \mathcal{A} \sqcup \mathcal{B}$.
- (1)' If there is $j \in J$ such that $x_{\langle \tilde{a}, \alpha \rangle} \in_P \mathcal{A}_j$, then $x_{\langle \tilde{a}, \alpha \rangle} \in_P \sqcup_{j \in J} \mathcal{A}_j$.
- (2) If $x_{<\tilde{a},\alpha>} \in_R \mathcal{A}$ or $x_{<\tilde{a},\alpha>} \in_R \mathcal{B}$, then $x_{<\tilde{a},\alpha>} \in_R \mathcal{A} \cup \mathcal{B}$.
- (2)' If there is $j \in J$ such that $x_{\leq \tilde{a}, \alpha >} \in_R \mathcal{A}_j$, then $x_{\leq \tilde{a}, \alpha >} \in_R \bigcup_{i \in J} \mathcal{A}_i$.

3. Concept lattice on cubic sets

In the section, we introduce the notions of Galois connection, formal context, concept lattice on cubic sets. First, Galois connection is the basic structure related to concept lattice, closure operator, etc..

Definition 3.1. A Galois connection between two cubic sets X and Y is a pair (f, g) of mappings

 $\begin{aligned} f: ([I] \times I)^X &\to ([I] \times I)^Y, \qquad g: ([I] \times I)^Y \to ([I] \times I)^X, \text{ satisfying} \\ (1) \quad \mathcal{A}_1 \sqsubset \mathcal{A}_2 \Rightarrow f(\mathcal{A}_2) \sqsubset f(\mathcal{A}_1), \qquad (2) \quad \mathcal{B}_1 \sqsubset \mathcal{B}_2 \Rightarrow g(\mathcal{B}_2) \sqsubset g(\mathcal{B}_1), \\ (3) \quad \mathcal{A} \sqsubset gf(\mathcal{A}), \qquad (4) \quad \mathcal{B} \sqsubset fg(\mathcal{B}), \\ \text{for any } \mathcal{A}, \mathcal{A}_1, \mathcal{A}_2 \in ([I] \times I)^X \text{ and } \mathcal{B}, \mathcal{B}_1, \mathcal{B}_2 \in ([I] \times I)^Y. \end{aligned}$

In the theory of formal concept analysis, formal context is one of important notions and tools. On cubic sets, a formal context is the triple structure (X, Y, \mathcal{R}) , where X is the sets of all objects, Y the sets of all attributes, \mathcal{R} is a cubic relation between them. See the follow example.

Example 3.2. Let $X = \{a, b, c\}, Y = \{p, q\}$ be two sets, a cubic relation \mathcal{R} between X and Y given by the following table:

$\mathcal{R} = < R, r >$	p	q
a	< [0, 1], 1 >	< [0.4, 0.8], 0.4 >
b	< [0.2, 0.6], 0.4 >	< [0.2, 1], 1 >
С	< [0, 0.5], 0.5 >	< [0.5, 1], 0.5 >
Table 3.1		

Suppose \mathcal{R} is a cubic relation on $X \times Y$, we obtain a Galois connection $(\uparrow_{\mathcal{R}}, \downarrow_{\mathcal{R}})$, in simple way, (\uparrow, \downarrow) .

Definition 3.3. Suppose \mathcal{R} is a cubic relation on $X \times Y$, for any $\mathcal{A} \in ([I] \times I)^X$, $\mathcal{B} \in ([I] \times I)^Y$, and for every $x \in X$, $y \in Y$, two cubic sets are defined.

$$\mathcal{A}^{\uparrow}(y) = \sqcap_{x \in X} \bigsqcup_{\substack{y_{< \widetilde{b}, \mu >} \in M_{Y}}} \{< \widetilde{b}, \mu > |< \widetilde{b}, \mu > \sqcap \mathcal{A}(x) \sqsubset \mathcal{R}(x, y)\}$$
$$\mathcal{B}^{\downarrow}(x) = \sqcap_{y \in Y} \bigsqcup_{\substack{x_{< \widetilde{a}, \lambda >} \in M_{X}}} \{< \widetilde{a}, \lambda > |< \widetilde{a}, \lambda > \sqcap \mathcal{B}(y) \sqsubset \mathcal{R}(x, y)\}$$

If $\mathcal{A}^{\uparrow} = \mathcal{B}$, and $\mathcal{A} = \mathcal{B}^{\downarrow}$, then $(\mathcal{A}, \mathcal{B})$ is called a *cubic concept* of (X, Y, \mathcal{R}) , \mathcal{A} is the extent and \mathcal{B} the intent of the concept $(\mathcal{A}, \mathcal{B})$. The set of all cubic concepts is denoted by $C(X, Y, \mathcal{R})$. i.e.,

 $C(X,Y,\mathcal{R}) = \{(\mathcal{A},\mathcal{B}) \mid \mathcal{A}^{\uparrow} = \mathcal{B}, \mathcal{A} = \mathcal{B}^{\downarrow}\}$

We define a partial order \sqsubset on $C(X, Y, \mathcal{R})$, for $(\mathcal{A}_1, \mathcal{B}_1), (\mathcal{A}_2, \mathcal{B}_2) \in C(X, Y, \mathcal{R})$, $(\mathcal{A}_1, \mathcal{B}_1) \sqsubset (\mathcal{A}_2, \mathcal{B}_2) \Leftrightarrow \mathcal{A}_1 \sqsubset \mathcal{A}_2$ (which is equivalent to $\mathcal{B}_2 \sqsubset \mathcal{B}_1$). Clearly $(\mathcal{A}_1, \mathcal{B}_1) = (\mathcal{A}_2, \mathcal{B}_2) \Leftrightarrow \mathcal{A}_1 = \mathcal{A}_2$ (which is equivalent to $\mathcal{B}_2 = \mathcal{B}_1$).

Next, we introduce a example as follows:

Example 3.4. Following Example 3.2, for a cubic set $\mathcal{A} = \hat{1}_X$ on X, we obtain \mathcal{A}^{\uparrow} ,

$$\begin{split} \mathcal{A}^{\top}(p) &= \sqcap_{x \in X} \bigsqcup_{p_{<\tilde{b}, \mu >} \in M_{Y}} \{ < b, \mu > | < b, \mu > \sqcap \mathcal{A}(x) \sqsubset \mathcal{R}(x, p) \} \\ &= (\bigsqcup_{p_{<\tilde{b}, \mu >} \in M_{Y}} \{ < \tilde{b}, \mu > | < \tilde{b}, \mu > \sqcap < \mathbf{1} \ , 1 > \sqsubset < [0, 1], 1 > \}) \\ &\sqcap (\bigsqcup_{p_{<\tilde{b}, \mu >} \in M_{Y}} \{ < \tilde{b}, \mu > | < \tilde{b}, \mu > \sqcap < \mathbf{1} \ , 1 > \sqsubset < [0.2, 0.6], 0.4 > \}) \\ &\sqcap (\bigsqcup_{p_{<\tilde{b}, \mu >} \in M_{Y}} \{ < \tilde{b}, \mu > | < \tilde{b}, \mu > \sqcap < \mathbf{1} \ , 1 > \sqsubset < [0, 0.5], 0.5 >) \\ &= < [0, 1], 1 > \sqcap < [0.2, 0.6], 0.4 > \sqcap < [0, 0.5], 0.5 > \\ &= < [0, 0.5], 0.4 >, \end{split}$$

In the same way, we obtain $\mathcal{A}^{\uparrow}(q) = < [0.2, 0.8], 0.4 >$. Then \mathcal{A}^{\uparrow} is a cubic set on Y.

Furthermore, let $\mathcal{B} = \mathcal{A}^{\uparrow}$, we obtain \mathcal{B}^{\downarrow} , $\mathcal{B}^{\downarrow}(a) = \langle \mathbf{1}, 1 \rangle$, $\mathcal{B}^{\downarrow}(b) = \langle \mathbf{1}, 1 \rangle$, $\mathcal{B}^{\downarrow}(c) = \langle \mathbf{1}, 1 \rangle$, i.e., $\mathcal{B}^{\downarrow} = \hat{1}_X$. Clearly \mathcal{B}^{\downarrow} is a cubic set on X, and $\mathcal{B}^{\downarrow} = \mathcal{A}$.

By Definition 3.3, we know that $(\mathcal{A}, \mathcal{B})$ is a cubic concept of the cubic formal context (X, Y, \mathcal{R}) .

Third, we prove the set $C(X, Y, \mathcal{R})$ forms a complete lattice.

$$\begin{aligned} \text{Proposition 3.5. Suppose } x \in X, y \in Y, we have \\ & (x_{<1,1>})^{\uparrow}(y) = (y_{<1,1>})^{\downarrow}(x) \end{aligned}$$

$$\begin{aligned} \text{Proof. (1)} \quad (x_{<1,1>})^{\uparrow}(y) &= \sqcap_{z \in X} \bigsqcup_{y_{<\tilde{b},\mu>}} \{<\tilde{b}, \mu > |<\tilde{b}, \mu > \sqcap x_{<1,1>}(z) \sqsubset \mathcal{R}(z,y)\} \\ &= (\sqcap_{z \neq x} \bigsqcup_{y_{<\tilde{b},\mu>}} \{<\tilde{b}, \mu > |<\tilde{b}, \mu > \sqcap x_{<1,1>}(z) \sqsubset \mathcal{R}(z,y)\}) \\ \quad \sqcap(\bigsqcup_{y_{<\tilde{b},\mu>}} \{<\tilde{b}, \mu > |<\tilde{b}, \mu > \sqcap x_{<1,1>}(x) \sqsubset \mathcal{R}(x,y)\}) \\ &= (\sqcap_{z \neq x} \bigsqcup_{y_{<\tilde{b},\mu>}} \{<\tilde{b}, \mu > |<\tilde{b}, \mu > \sqcap < \mathbf{0}, 0 > \sqsubset \mathcal{R}(z,y)\}) \\ \quad \sqcap(\bigsqcup_{y_{<\tilde{b},\mu>}} \{<\tilde{b}, \mu > |<\tilde{b}, \mu > \sqcap < \mathbf{1}, 1 > \sqsubset \mathcal{R}(x,y)\}) \\ &= (\sqcap_{z \neq x} \{<\mathbf{1}, 1 > \}) \sqcap(\bigsqcup_{y_{<\tilde{b},\mu>}} \{<\tilde{b}, \mu > |<\tilde{b}, \mu > \sqsubset \mathcal{R}(x,y)\}) \\ &= (1, 1 > \sqcap(\bigsqcup_{y_{<\tilde{b},\mu>}} \{<\tilde{b}, \mu > |<\tilde{b}, \mu > \sqsubset \mathcal{R}(x,y)\}) \\ &= <\mathbf{1}, 1 > \sqcap(\bigsqcup_{y_{<\tilde{b},\mu>}} \{<\tilde{b}, \mu > |<\tilde{b}, \mu > \sqsubset \mathcal{R}(x,y)\}) \\ &= <\mathbf{1}, 1 > \sqcap(\sqcup_{y_{<\tilde{b},\mu>}} \{<\tilde{b}, \mu > |<\tilde{b}, \mu > \sqsubset \mathcal{R}(x,y)\}) \end{aligned}$$

$$= \bigcup_{\substack{y_{<\widetilde{b},\mu>}}} \{<\widetilde{b},\mu> \mid <\widetilde{b},\mu> \sqsubset \mathcal{R}(x,y)\}$$

= $\mathcal{R}(x,y),$
(2) In the similar way, we obtain $(y_{<1,1>})^{\downarrow}(x) = \mathcal{R}(x,y).$
So $x_{<1,1>}(y) = \mathcal{R}(x,y) = y_{<1,1>}(x).$

Now, we consider the surpremum, the infimum of a family cubic concepts of $C(X, Y, \mathcal{R})$.

Proposition 3.6. Suppose (A_j, B_j) $(j \in J)$ is a family of cubic concepts of $C(X, Y, \mathcal{R})$, then

$$(\bigsqcup_{j\in J}\mathcal{A}_j)^{\uparrow} = \sqcap_{j\in J}\mathcal{A}_j^{\uparrow}, \qquad (\bigsqcup_{j\in J}\mathcal{B}_j)^{\downarrow} = \sqcap_{j\in J}\mathcal{B}_j^{\downarrow}.$$

 $\begin{array}{l} \textit{Proof.} \ (1) \ \text{ We first to prove } (\bigsqcup_{j \in J} \mathcal{A}_j)^{\uparrow} = \sqcap_{j \in J} \mathcal{A}_j^{\uparrow}.\\\\ \text{ For every } y_{<\widetilde{b},\mu>} \in (\bigsqcup_{j \in J} \mathcal{A}_j)^{\uparrow}\\ \Rightarrow \forall x \in X, \quad <\widetilde{b},\mu>\sqcap(\bigsqcup_{j \in J} \mathcal{A}_j)(x) \sqsubset \mathcal{R}(x,y)\\ \Rightarrow \forall x \in X, \quad <\widetilde{b},\mu>\sqcap \bigsqcup_{j \in J} \mathcal{A}_j(x) \sqsubset \mathcal{R}(x,y)\\ \Rightarrow \forall j \in J, \quad \forall x \in X, \quad <\widetilde{b},\mu>\sqcap \mathcal{A}_j(x) \sqsubset \mathcal{R}(x,y)\\ \Rightarrow \forall j \in J, \quad y_{<\widetilde{b},\mu>} \in \mathcal{A}_j^{\uparrow}\\ \Rightarrow \forall j \in J, \quad y_{<\widetilde{b},\mu>} \in \sqcap_{j \in J} \mathcal{A}_j^{\uparrow}.\\\\ \text{ Then } (\bigsqcup_{j \in J} \mathcal{A}_j)^{\uparrow} \sqsubset \sqcap_{j \in J} \mathcal{A}_j^{\uparrow}.\end{array}$

 $j \in J$ On the other hand,

For every $y_{<\widetilde{b},\mu>} \in \sqcap_{j\in J} \mathcal{A}_j^{\uparrow}$

$$\begin{split} &\Rightarrow \forall j \in J, \quad y_{<\widetilde{b},\mu>} \in \mathcal{A}_{j}^{\uparrow} \\ &\Rightarrow \forall j \in J, \quad \forall x \in X, \quad <\widetilde{b},\mu> \sqcap \mathcal{A}_{j}(x) \sqsubset \mathcal{R}(x,y) \\ &\Rightarrow \forall x \in X, \quad <\widetilde{b},\mu> \sqcap \bigsqcup_{j \in J} \mathcal{A}_{j}(x) \sqsubset \mathcal{R}(x,y) \\ &\Rightarrow \forall x \in X, \quad <\widetilde{b},\mu> \sqcap \bigsqcup_{j \in J} \mathcal{A}_{j})(x) \sqsubset \mathcal{R}(x,y) \\ &\Rightarrow \forall x \in X, \quad <\widetilde{b},\mu> \sqcap (\bigsqcup_{j \in J} \mathcal{A}_{j})(x) \sqsubset \mathcal{R}(x,y) \\ &\Rightarrow y_{<\widetilde{b},\mu>} \in (\bigsqcup_{j \in J} \mathcal{A}_{j})^{\uparrow}. \end{split}$$

Thus $\sqcap_{j \in J} \mathcal{A}_j^{\uparrow} \sqsubset (\bigsqcup_{j \in J} \mathcal{A}_j)^{\uparrow}$. So the equality holds. (2) In the similar way.

Since $\mathcal{A}_{j}^{\uparrow} = \mathcal{B}_{j}$, we obtain $(\bigsqcup_{j \in J} \mathcal{A}_{j})^{\uparrow} = \sqcap_{j \in J} \mathcal{A}_{j}^{\uparrow} = \sqcap_{j \in J} \mathcal{B}_{j}$. 35 In the similar way, we prove $(\sqcap_{j\in J}\mathcal{B}_j)^{\downarrow} = \bigsqcup_{j\in J}\mathcal{A}_j$. Then $((\bigsqcup_{j\in J}\mathcal{A}_j)^{\uparrow\downarrow}, \sqcap_{j\in J}\mathcal{B}_j)$ is a cubic concept of (X, Y, \mathcal{R}) .

Moreover, we also obtain $(\prod_{j \in J} \mathcal{A}_j, (\bigsqcup_{j \in J} \mathcal{B}_j)^{\downarrow\uparrow})$ is a cubic concept of (X, Y, \mathcal{R}) . In $C(X, Y, \mathcal{R})$, we define $((\bigsqcup_{j \in J} \mathcal{A}_j)^{\uparrow\downarrow}, \prod_{j \in J} \mathcal{B}_j)$ as the surpremum of the family

 $(\mathcal{A}_j, \mathcal{B}_j)$ $(j \in J)$ cubic concepts of (X, Y, \mathcal{R}) , and $(\prod_{j \in J} \mathcal{A}_j, (\bigsqcup_{j \in J} \mathcal{B}_j)^{\downarrow\uparrow})$ as the infimum of the family $(\mathcal{A}_j, \mathcal{B}_j)$ $(j \in J)$ cubic concepts of (X, Y, \mathcal{R}) . So we obtain,

Proposition 3.7. $(C(X,Y,\mathcal{R}), \sqsubset)$ forms a complete lattice with respect the operators \bigsqcup and \sqcap , where

$$\bigsqcup_{j\in J}(\mathcal{A}_j,\mathcal{B}_j) = ((\bigsqcup_{j\in J}\mathcal{A}_j)^{\uparrow\downarrow}, \sqcap_{j\in J}\mathcal{B}_j), \qquad \sqcap_{j\in J}(\mathcal{A}_j,\mathcal{B}_j) = (\sqcap_{j\in J}\mathcal{A}_j, (\bigsqcup_{j\in J}\mathcal{B}_j)^{\downarrow\uparrow}).$$

By Definition 3.3 and the above results, we obtain

Proposition 3.8. Suppose \mathcal{R} is a cubic relation between X and Y, then $(\uparrow_{\mathcal{R}}, \downarrow_{\mathcal{R}})$ is a Galois connection induced by \mathcal{R} .

Conversely, suppose (\uparrow, \downarrow) is a Galois connection on cubic sets, we also define a cubic relation \mathcal{R} as: for any $x \in X, y \in Y$, $\mathcal{R}(x, y) = (x_{<1,1>})^{\uparrow}(y) = (y_{<1,1>})^{\downarrow}(x)$, and (\uparrow, \downarrow) is precisely generated by \mathcal{R} , i.e., $(\uparrow, \downarrow) = (\uparrow_{\mathcal{R}}, \downarrow_{\mathcal{R}})$.

Note: In the paper, we introduce the notions of Galois connection, Concept lattice on cubic sets with respect to the P-order, P-union, P-intersection. The above results hold with respect to the R-order, R-union, R-intersection.

CONCLUSION

In the paper, we define formal context, Galois connection, concept lattice on cubic sets, thus combine formal concept analysis and cubic sets together, provide a new platform to further study.

Acknowledgements. The author would like to thank the referees for their valuable suggestions and comments, which have helped improve this paper significantly.

References

- [1] L. A. Zadeh, Fuzzy sets, Information and Control 8 (1965) 338–353.
- [2] Y. B. Jun, C. S. Kim and K. O. Yang, Cubic sets, Ann. Fuzzy Math. Inform. 4 (1) (2012) 83–98.
- [3] J. G. Kang and C. S. Kim, Mappings of Cubic sets, Commun. Korean Math. Soc. 31 (3) (2016) 423–431.
- [4] J. H. Kim, P. K. Lim, J. G. Lee and K. Hur, Cubic relations, Ann. Fuzzy Math. Inform. 19 (1) (2020) 21–43.
- Y. B. Jun and K. J. Lee, Closed cubic ideals and cubic 0-subalgebras in BCK/BCI-algebras, Appl. Math. Sci. 4 (65–68) (2009) 3395–3402.
- [6] Y. B. Jun, K. J. Lee and M. S. Kang, Cubic structures applied to ideals of BCI-algebras, Comput. Math. Appl. 62 (9) (2011) 3334–3342.
- [7] Y. B. Jun, S. T. Jun and M. S. Kim, Cubic subgroups, Ann. Fuzzy Math. Inform. 2 (1) (2011) 9–15.
- [8] Y. B. Jun and A. Khan, Cubic ideals in semigroups, Honam Math. J 35(4) (2013) 607-623.

- M. Akram, N. Yaqoob and M. Gulistan, Cubic KU-subalgebras, Int, J. Pure. Appl. Math. 89 (5) (2013) 659–665.
- [10] N. Yaqoob, S. M. Mostafa and M. A. Ansari, On cubic KU-ideals on KU-algebras, ISRN algebra (2013) Art, ID 935905, 10 pages.
- [11] S. S. Ahn and J. M. Ko, Cubic subalgebras and filters of CI-algebras, Honam Math. J 36 (1) (2014) 43–54.
- [12] Y. B. Jun, S. Smarandache and C. S. Kim, Neutrosophic cubic sets, New Mathematics and Natural Computation (2015) 1–15.
- [13] T. Senapati, C. S. Kim, M. Bhowmik and M. Pal, Cubic subalgebras and cubic closed ideals of B-algebras, Fuzzy Inf. Eng. 7 (2) (2015) 129–149.
- [14] V. Chinnadurai, K. Suganya and A. Swaminathan, Some characterizations on cubic sets, Int. J. Multidisciplinary Research and Modern Education II (2016) 258–264.
- [15] T. Mahmood, S. Abdulla, S. R. Rashid and M. Bila, Multicriteria decision making based on cubic set, J. of New theory 16 (2017) 1–9.
- [16] Xue-Ling Ma, Jianming Zhan, Madad Khan, Muhammad Gulistan and Naveed Yaqoob, Generalized cubic relations in H_v-LA-semigroups, Journal of Discrete Mathematical Sciences and Cryptography 21 (3) (2018) 607–630.
- [17] Y. B. Jun, S. Z. Song and S. J. Kim Cubic interval-valued intuitionistic fuzzy sets and their application in BCK/BCI-algebras, Axioms 2018, 7, 7; doi:10.3390/axioms7010007.
- [18] J. G. Lee, K. Hur and Xueyou Chen, A study on cubic *H*-relations in a topological universe view-point, Mathematics 8 (2020),doi:10.3390/8040482.
- [19] B. Ganter and R. Wille, Formal concept analysis, Springer-Verlag 1999.
- [20] R. Bělohlávek, Fuzzy relational systems, Foundations and Principles, Kluwer, New York 2002.
- [21] Y. Djouadi and H. Prade, Interval-valued fuzzy formal concept analysis, Lecture Notes in Artificial Intelligence Vol 5722 (2009) 592–601.
- [22] A. Brito, L. Barros, E Laureano, F. Bertato and M. Coniglio, Fuzzy formal concept analysis Communications in Computer and Information Science 831 (2018) 192–205.
- [23] Xueyou Chen and Qingguo Li, Formal topology, Chu space and approximable concept, Proc. of CLA 2005, https://sunsite.informatik.rwth-aachen.de/Publications/CEUR-WS/VOL-162/
- [24] Xueyou Chen, Qingguo Li and Zike Deng, Chu space and approximable concept lattice in fuzzy setting, Proceedings of Fuzz-IEEE 2007, IEEE international conference of fuzzy systems, London, UK, July 23–26. PP 1–6 (2007), ISSN:1098-7584.
- [25] L. A. Zadeh, The concept of a linguistic variable and its application to approximate reasoning-I, Inform. Sci. 8 (1975) 199–249.
- [26] M. B. Gorzalczany, A method of inference in approximate reasoning based on interval-valued fuzzy sets, Fuzzy sets and Systems 21 (1987) 1–17.
- [27] K. Hur, J. G. Lee and J. Y. Choi, Interval-valued fuzzy relations, JKIIS 19 (3) (2009) 425–432.

Xueyou Chen (math-chen@qq.com)

School of Mathematics, Shandong University of Technology, Zibo, Shandong 255049, P. R. CHINA