

The diagram illustrates the recursive construction of a fractal curve using four basic moves: @, F, M, and I. The moves are color-coded: @ is black, F is red, M is green, and I is blue. The construction starts with a single '@' move, which branches into two paths. Each path then branches into two more paths, resulting in a total of eight paths. The moves are arranged in a way that shows how the fractal curve is built up from these basic moves.

Reprinted from the
Annals of Fuzzy Mathematics and Informatics
Vol. 28, No. 2, October 2024

Separation axioms in interval-valued soft topological spaces

J. I. BAEK, G. ŞENEL, S. JAFARI, S. H. HAN, K. HUR

Received 1 April 2024; Revised 20 May 2024; Accepted 27 May 2024

ABSTRACT. Our purpose of the research is to investigate two aspects: First, we introduce the concept of separation axioms in an interval-valued soft topological space, and obtain some of its properties and give some examples. Second, we propose new separation axioms in an interval-valued soft topological space by using interval-valued α -open sets basic topological structures based on interval-valued soft set, and study some of their basic properties. Furthermore, we deal with hereditary problems of each separation axiom.

2020 AMS Classification: 54A40, 54C08, 54C10.

Keywords: Interval-valued soft topological space, $IVST_i(j)$ -space ($i=0,1,2,3,4$; $j=i,ii$), Interval-valued α -open [resp. closed] set, $IVS\alpha T_i(j)$ -space ($i=0,1,2,3,4$; $j=i,ii$).

Corresponding Author: S. Jafari (jafaripersia@gmail.com,
saeidjafari@topositus.com)
S. H. Han (shhan235@wku.ac.kr)

1. INTRODUCTION

In 1999, Molodtsov [1] introduced the concept of soft sets which has rich potential for practical applications in several domains as a tool for dealing with uncertainties. After then, Maji et al. [2] proposed some basic operations on soft sets and studied some of their properties (See [3, 4, 5] for the further researches). Moreover, many researchers have applied the notion of soft sets to various fields, for example abstract algebra (See [6, 7, 8, 9]), logical algebra (See [10, 11, 12]), decision making problems (See [13, 14, 15]), topology (See [16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26]), topological group (See [27, 28, 29, 30, 31, 32]), etc. In particular, Bayramov et al. [33] introduced the concepts of interval-valued fuzzy soft topological spaces, interval-valued fuzzy soft neighborhoods, interval-valued fuzzy soft continuities, etc. and studied some of their properties.

Topology is an important area of mathematics with many applications in the domains of computer and physical science. In particular, separation axioms in topology not only provide a systematic way to classify and study topological spaces but also ensure that the spaces behave in predictable and desirable ways for further mathematical exploration and application. Moreover, understanding the separation properties of a space is often the first step in analyzing its structure and behavior in both theoretical and applied settings. Then their necessity and importance cannot be overstated. In 2021, Lee et al. [34] investigated interval-valued soft topological structures as a generalization of soft topologies.

We intend to study in the following two aspects: First, we define separation axioms in an interval-valued soft topological space by modifying separation axioms introduced by Shabir and Naz [16]. Second, we introduce the concept of separation axioms in an interval-valued soft topological space by modifying separation axioms proposed by Akdag and Ozkan [35]. In order to realize our aim, this paper is composed of six sections. In Section 2, we recall some basic concepts required in each section. In Section 3, we introduce the notions of separation axioms in an interval-valued soft topological space in the sense of Shabir and Naz [16], and obtain its basic properties and give some examples. In Section 4, we deal with hereditary properties of separation axioms discussed in Section 3. In Section 5, we propose the concept of interval-valued soft α -open set by modifying soft α -open set introduced by Akdag and Ozkan [36] and study some of its properties. In Section 6, we define separation axioms in an interval-valued soft topological space in the viewpoint of Akdag and Ozkan [35] and discuss some of their properties.

2. PRELIMINARIES

In this section, we recall basic concepts needed in next sections. Throughout this paper, let $X, Y, Z, \dots$ denote non-empty universe sets, let $E, E', E'', \dots$ denote non-empty sets of parameters and let 2^X denote the power set of X .

Definition 2.1 ([37, 38]). The form

$$[A^-, A^+] = \{B \subset X : A^- \subset B \subset A^+\}$$

is called an *interval-valued set* (briefly, IVS) or *interval set* in X , if $A^-, A^+ \subset X$ and $A^- \subset A^+$. In this case, A^- [resp. A^+] represents the set of minimum [resp. maximum] memberships of elements of X to A . $[\emptyset, \emptyset]$ [resp. $[X, X]$] is called the *interval-valued empty* [resp. *whole*] set in X and will be denoted by $\tilde{\emptyset}$ [resp. $\tilde{X}$]. We will denote the set of all IVSs in X as $IVS(X)$.

It is obvious that $[A, A] \in IVS(X)$ for a classical subset A of X . Then we consider an IVS in X as the generalization of a classical subset of X . Furthermore, if $A = [A^-, A^+] \in IVS(X)$, then

$$\chi_A = [\chi_{A^-}, \chi_{A^+}]$$

is an interval-valued fuzzy set in X introduced by Zadeh [39], where χ_A denotes the characteristic function of A . Thus we can consider an interval-valued fuzzy set as the generalization of an IVS.

Definition 2.2 ([37, 38]). Let $A, B \in IVS(X)$. Then

- (i) we say that A *contained in* B , denoted by $A \subset B$, if $A^- \subset B^-$ and $A^+ \subset B^+$,
- (ii) we say that A *equals to* B , denoted by $A = B$, if $A \subset B$ and $B \subset A$,
- (iii) the *complement* of A , denoted A^c , is an interval-valued set in X defined by:

$$A^c = [(A^+)^c, (A^-)^c],$$

- (iv) the *union* of A and B , denoted by $A \cup B$, is an interval-valued set in X defined by:

$$A \cup B = [A^- \cup B^-, A^+ \cup B^+],$$

- (v) the *intersection* of A and B , denoted by $A \cap B$, is an interval-valued set in X defined by:

$$A \cap B = [A^- \cap B^-, A^+ \cap B^+].$$

Definition 2.3 ([37]). Let $a \in X$ and $A \in IVS(X)$. Then the notation $[\{a\}, \{a\}]$ [resp. $[\emptyset, \{a\}]$] is called an *interval-valued* [resp. *vanishing*] *point* in X and denoted by a_1 [resp. a_0]. We denote the set of all interval-valued points in X as $IV_P(X)$.

- (i) We say that a_1 *belongs to* A , denoted by $a_1 \in A$, if $a \in A^-$.
- (ii) We say that a_0 *belongs to* A , denoted by $a_0 \in A$, if $a \in A^+$.

Definition 2.4 ([37]). Let τ be a non-empty family of IVSs on X . Then τ is called an *interval-valued topology* (briefly, IVT) on X , if it satisfies the following axioms:

- (IVO₁) $\tilde{\emptyset}, \tilde{X} \in \tau$,
- (IVO₂) $A \cap B \in \tau$ for any $A, B \in \tau$,
- (IVO₃) $\bigcup_{j \in J} A_j \in \tau$ for any family $(A_j)_{j \in J}$ of members of τ .

In this case, the pair (X, τ) is called an *interval-valued topological space* (briefly, IVTS) and each member of τ is called an *interval-valued open set* (briefly, IVOS) in X . An IVS A is called an *interval-valued closed set* (briefly, IVCS) in X , if $A^c \in \tau$.

It is obvious that $\{\tilde{\emptyset}, \tilde{X}\}$ is an IVT on X , and is called the interval-valued indiscrete topology on X and denoted by $\tau_{IV,0}$. Also $IVS(X)$ is an IVT on X , and is called the *interval-valued discrete topology* on X and is denoted by $\tau_{IV,1}$. The pair $(X, \tau_{IV,0})$ [resp. $(X, \tau_{IV,1})$] is called the *interval-valued indiscrete* [resp. *discrete*] *space*.

We denote the set of all IVTs on X as $IVT(X)$. For an IVTS X , we denote the set of all IVOSs [resp. IVCSs] in X as $IVO(X)$ [resp. $IVC(X)$].

Definition 2.5 ([1, 17]). An F_A is called a *soft set* over X , if $F_A : A \rightarrow 2^X$ is a mapping such that $F_A(e) = \emptyset$ for each $e \notin A$, where $A \in 2^E$.

In other words, a soft set over X is a parametrized family of subsets of X . For each $e \in A$, $F_A(e)$ may be considered as the set of e -approximate elements of the soft set F_A . It is clear that a soft set is not a set. We will denote the set of all soft sets over X as $SS(X)$ while $SS(X)_E$ will denote the set of all soft sets over with respect to a fixed set E of parameters.

It has been well-known [1] that every Zadeh's fuzzy set A may be considered as the soft set $F_{[0,1]}$.

Definition 2.6 ([2, 17]). Let $F_A, F_B \in SS(X)$ and $A, B \in 2^E$. Then we say that:

- (i) F_A is a *soft subset* of F_B , denoted by $F_A \widetilde{\subset} F_B$, if $A \subset B$ and $F_A(e) \subset F_B(e)$ for each $e \in A$,
- (ii) F_A is a *soft super set* of F_B , denoted by $F_A \widetilde{\supset} F_B$, if $F_B \widetilde{\subset} F_A$,
- (iii) F_A and F_B are *soft equal*, if $F_A \widetilde{\subset} F_B$ and $F_A \widetilde{\supset} F_B$.

Definition 2.7 ([2, 3]). Let $F_A \in SS(X)$. Then F_A is called:

- (i) a *null soft set* or a *relative null soft set* (with respect to A), denoted by $\emptyset_A$, if $F_A(e) = \emptyset$ for each $e \in A$,
- (ii) an *absolute soft set* or a *relative whole soft set* (with respect to A), denoted by X_A , if $F_A(e) = X$ for each $e \in A$.

We will denote the null [resp. absolute] soft set in $SS_E(X)$ by X_E [resp. $\emptyset_E$].

Definition 2.8 ([16]). Let τ be a collection of members of $SS_E(X)$. Then τ is called a *soft topology* on X , if it satisfies the following conditions:

- (i) $\emptyset_E, X_E \in \tau$,
- (ii) $A \cap B \in \tau$ for any $A, B \in \tau$,
- (iii) $\bigcup_{j \in J} A_j \in \tau$ for each $(A_j)_{j \in J} \subset \tau$, where J denotes an index set.

Each member of τ is called a *soft open set in X* and the complement of each member of τ is called a *closed soft set in X* . The triple (X, τ, E) is called a *soft topological space* over X .

Definition 2.9 ([34]). Let $A \in 2^E$. Then an $\mathbf{F}_A = [F_A^-, F_A^+]$ is called an *interval-valued soft set* (briefly, IVSS) over X , if $\mathbf{F}_A : A \rightarrow IVS(X)$ is a mapping such that $\mathbf{F}_A(e) = \widetilde{\emptyset}$ for each $e \notin A$, i.e., $F_A^-, F_A^+ \in SS(X)$ such that $F_A^-(e) \subset F_A^+(e)$ for each $e \in A$.

In other words, an IVSS over X is a parametrized family of IVSSs of X . For each $e \in A$, $\mathbf{F}_A(e) = [F_A^-(e), F_A^+(e)]$ may be considered as an interval-valued set of e -approximate elements of the IVSS $\mathbf{F}_A$. We denote the set of all IVSSs over X as $IVSS(X)$.

It is obvious that if $F_A \in SS(X)$, then $[F_A, F_A] \in IVSS(X)$. Now we can see that an IVSS is the generalization of a soft set. Moreover, if $\mathbf{F}_A \in IVSS(X)$, then clearly, $\chi_{\mathbf{F}_A}$ is an interval-valued fuzzy soft set (briefly, IVFSS) over X introduced by Yang et al. [40]. Thus an IVSS is the special case of an IVFSS.

Definition 2.10 ([34]). Let $A \in 2^E$ and $\mathbf{F}_A \in IVSS(X)$. $\mathbf{F}_A$ is called:

- (i) a *relative null interval-valued soft set* (with respect to A), denoted by $\widetilde{\emptyset}_A$, if $\mathbf{F}_A(e) = \widetilde{\emptyset}$ for each $e \in A$,
- (ii) a *relative whole interval-valued soft set* (with respect to A), denoted by $\widetilde{X}_A$, if $\mathbf{F}_A(e) = \widetilde{X}$ for each $e \in A$.

We denote the set of all IVSSs over X with respect to the fixed parameter set A as $IVSS_A(X)$.

Now we will denote the members of $IVSS_E(X)$ by $\mathbf{A}, \mathbf{B}, \mathbf{C}, \dots$. In fact, $\mathbf{A}, \mathbf{B}, \mathbf{C} : E \rightarrow IVS(X)$. In particular, the interval-valued soft empty [resp. whole] set over X with respect to E , denoted by $\widetilde{\emptyset}_E$ [resp. $\widetilde{X}_E$], is the IVS in X defined by $\widetilde{\emptyset}_E(e) = \widetilde{\emptyset}$ [resp. $\widetilde{X}_E(e) = \widetilde{X}$] for each $e \in E$.

Definition 2.11 ([34]). Let $\mathbf{A}, \mathbf{B} \in IVSS_E(X)$. We say that

- (i) $\mathbf{A}$ is an *interval-valued soft subset* of $\mathbf{B}$, denoted by $\mathbf{A} \subset \mathbf{B}$, if $\mathbf{A}(e) \subset \mathbf{B}(e)$ for each $e \in E$,
- (ii) $\mathbf{A}$ and $\mathbf{B}$ are *interval-valued soft equal*, denoted by $\mathbf{A} = \mathbf{B}$, if $\mathbf{A} \subset \mathbf{B}$ and $\mathbf{B} \subset \mathbf{A}$,
- (iii) the *interval-valued soft complement* of $\mathbf{A}$, denoted by $\mathbf{A}^c$, is the mapping $\mathbf{A}^c : E \rightarrow IVS(X)$ defined as: for each $e \in E$,

$$\mathbf{A}^c(e) = (\mathbf{A}(e))^c.$$

3. INTERVAL-VALUED SOFT SEPARATION AXIOMS

In this section, we introduce the concepts of separation axioms in interval-valued soft topological spaces, for example, $IVST_j(i)$ -space for $j = 0, 1, 2, 3, 4$ and $i = 1, 2$. And we study some of their properties and some relationships among them. Also, we give some examples.

Definition 3.1. Let $\mathbf{A} \in IVSS_E(X)$ and $x \in X$.

- (i) x_1 said to *belong* or *totally belong* to $\mathbf{A}$, denoted by $x_1 \in \mathbf{A}$, if $x \in A^-(e)$ for each $e \in E$.
- (ii) x_0 said to *belong* or *totally belong* to $\mathbf{A}$, denoted by $x_0 \in \mathbf{A}$, if $x \in A^+(e)$ for each $e \in E$.

Note that for any $x \in X$, $x_1 \notin \mathbf{A}$ [resp. $x_0 \notin \mathbf{A}$], if $x \notin A^-(e)$ [resp. $x \notin A^+(e)$] for some $e \in E$. It is obvious that if $x_1 \in \mathbf{A}$, then $x_0 \in \mathbf{A}$. But the converse is not true in general (See Example 3.2).

Example 3.2. Let $X = \{a, b, c, x, y, z\}$ be a universe set and $E = \{e, f, g\}$ a set of parameters. Consider the IVSS $\mathbf{A}$ defined by:

$$\mathbf{A}(e) = [\{a, b\}, \{a, b, c\}], \mathbf{A}(f) = [\{a\}, \{a, c, z\}], \mathbf{A}(g) = [\{a, c, x\}, \{a, c, x\}].$$

Then it can be easily checked that $a_1, a_0 \in \mathbf{A}$, $c_0 \in \mathbf{A}$ but $b_1, c_1, x_1, y_1, z_1 \notin \mathbf{A}$ and $b_0, x_0, y_0, z_0 \notin \mathbf{A}$.

Definition 3.3 ([34]). Let τ be a family of IVSSs over X with respect to E . Then τ is called an *interval-valued soft topology* (briefly, IVST) on X with respect to E , if it satisfies the following axioms:

- [IVSO₁] $\tilde{\emptyset}_E, \tilde{X}_E \in \tau$,
- [IVSO₂] $\mathbf{A} \cap \mathbf{B} \in \tau$ for any $\mathbf{A}, \mathbf{B} \in \tau$,
- [IVSO₃] $\bigcup_{j \in J} \mathbf{A}_j \in \tau$ for each $(\mathbf{A}_j)_{j \in J} \subset \tau$.

The triple (X, τ, E) is called an *interval-valued soft topological space* (briefly, IVSTS). Every member of τ is called an *interval-valued soft open set* (briefly, IVSOS) and the complement of an IVSOS is called an *interval-valued soft closed set* (briefly, IVSCS) in X . The set of all IVSOSs [resp. IVSCSs] in X is denoted by $IVSO(X)$ [resp. $IVSC(X)$]. It is obvious that $\{\tilde{\emptyset}_E, \tilde{X}_E\}, IVSS_E(X) \in IVST_E(X)$, where $IVST_E(X)$ denotes the set of all IVSTs on X with respect to E . In this case, $\{\tilde{\emptyset}_E, \tilde{X}_E\}$ [resp. $IVSS_E(X)$] is called an *interval-valued soft indiscrete* [resp. *discrete*] topology on X and is denoted by $\tilde{\tau}_0$ [resp. $\tilde{\tau}_1$]. We will denote the set of all IVSTSs over X with respect to E as $IVSTS_E(X)$ and denote the set of all IVSCSs in an IVSTS (X, τ, E) by τ^c . In fact,

$$\tau^c = \{\mathbf{A} \in IVSS_E(X) : \mathbf{A}^c \in \tau\}.$$

It is obvious that if $\tau \in IVST_E(X)$, then $\chi_\tau = \{\chi_U : U \in \tau\}$ is an interval-valued fuzzy soft topology (briefly, IVFST) on X defined by Ali et al. [41]. Thus an IVFST is the generalization of an IVST.

From Remark 4.3 in [34], we can consider (X, τ^-, τ^+) as a soft bi-topological space in the sense of Kelly [42] for each $\tau \in IVST_E(X)$, where

$$\tau^- = \{U^- \in IVS(X) : U \in \tau\}, \quad \tau^+ = \{U^+ \in IVS(X) : U \in \tau\}.$$

Result 3.4 (Proposition 4.5, [34]). *Let (X, τ, E) be an IVSTS and for each $e \in E$, let*

$$\tau_e = \{U(e) \in IVS(X) : U \in \tau\}.$$

Then τ_e is an interval-valued topology (briefly, IVT) on X introduced by Kim et al. [37]. In this case, τ_e will be called a parametric interval-valued topology and (X, τ_e) will be called a parametric interval-valued topological space.

Furthermore, we obtain two classical topologies on X for each IVSTS (X, τ, E) and each $e \in E$ given as follows (See Remark 4.6 (1), [34]):

$$\tau_e^- = \{A(e)^- \in 2^X : A(e) \in \tau_e\} \text{ and } \tau_e^+ = \{A(e)^+ \in 2^X : A(e) \in \tau_e\}.$$

In this case, τ_e^- and τ_e^+ will be called *parametric topologies* on X .

Definition 3.5. An IVSTS (X, τ, E) is called an:

- (i) *interval-valued soft T_0 (i)-space* (briefly, $IVST_0$ (i)-space), if for any $x, y \in X$ with $x \neq y$, there is $U, V \in \tau$ such that either $x_1 \in U, y_1 \notin U$ or $y_1 \in V, x_1 \notin V$,
- (ii) *interval-valued soft T_0 (ii)-space* (briefly, $IVST_0$ (ii)-space), if for any $x, y \in X$ with $x \neq y$, there is $U, V \in \tau$ such that either $x_0 \in U, y_0 \notin U$ or $y_0 \in V, x_0 \notin V$,
- (iii) *interval-valued soft T_1 (i)-space* (briefly, $IVST_1$ (i)-space), if for any $x, y \in X$ with $x \neq y$, there are $U, V \in \tau$ such that $x_1 \in U, y_1 \notin U$ and $y_1 \in V, x_1 \notin V$,
- (iv) *interval-valued soft T_1 (ii)-space* (briefly, $IVST_1$ (ii)-space), if for any $x, y \in X$ with $x \neq y$, there are $U, V \in \tau$ such that $x_0 \in U, y_0 \notin U$ and $y_0 \in V, x_0 \notin V$.

Remark 3.6. (1) Every $IVST_1$ (i) [resp. $IVST_1$ (ii)]-space is an $IVST_0$ (i) [resp. $IVST_0$ (ii)]-space. The converse is not true in general (See Example 3.7)

(2) Every $IVST_0$ (i) [resp. $IVST_1$ (i)]-space is an $IVST_0$ (ii) [resp. $IVST_1$ (ii)]-space. The converse is not true in general (See Example 3.7)

Example 3.7. Let $X = \{a, b, c\}$ be a universe set and $E = \{e, f\}$ a set of parameters.

- (1) Consider IVST τ_1 on X given by:

$$\tau_1 = \{\tilde{\varnothing}_E, \mathbf{A}, \mathbf{B}, \mathbf{C}, \mathbf{D}, \mathbf{E}, \mathbf{F}, \mathbf{G}, \tilde{X}_E\},$$

where $\mathbf{A}(e) = \mathbf{A}(f) = [\varnothing, \{c\}]$, $\mathbf{B}(e) = \mathbf{B}(f) = \{\varnothing, \{b, c\}\}$,
 $\mathbf{C}(e) = \mathbf{C}(f) = \{\{a\}, \{a, c\}\}$, $\mathbf{D}(e) = \mathbf{D}(f) = [\{c\}, \{b, c\}]$,
 $\mathbf{E}(e) = \mathbf{E}(f) = [\{a, b\}, X]$, $\mathbf{F}(e) = \mathbf{F}(f) = [\{a, c\}, X]$,
 $\mathbf{G}(e) = \mathbf{G}(f) = [\{a\}, X]$.

Then clearly, (X, τ_1, E) is an $IVST_0$ (i)-space. On the other hand, for $a \neq b \in X$, we cannot find $U \in \tau_1$ such that $a_1 \in U, b_1 \notin U$. Thus X is not an $IVST_1$ (i)-space.

- (2) Consider IVST τ_2 on X given by:

$$\tau_2 = \{\tilde{\varnothing}_E, \mathbf{A}, \mathbf{B}, \mathbf{C}, \mathbf{D}, \mathbf{E}, \mathbf{F}, \mathbf{G}, \mathbf{H}, \mathbf{I}, \mathbf{J}, \tilde{X}_E\},$$

where $\mathbf{A}(e) = \mathbf{A}(f) = [\emptyset, \{a\}]$, $\mathbf{B}(e) = \mathbf{B}(f) = \{\emptyset, \{b\}\}$, $\mathbf{C}(e) = \mathbf{C}(f) = \{\emptyset, \{c\}\}$,
 $\mathbf{D}(e) = \mathbf{D}(f) = [\{a\}, \{a, c\}]$, $\mathbf{E}(e) = \mathbf{E}(f) = [\{b\}, \{a, b\}]$,
 $\mathbf{F}(e) = \mathbf{F}(f) = [\{c\}, \{b, c\}]$, $\mathbf{G}(e) = \mathbf{G}(f) = [\{a, b\}, X]$,
 $\mathbf{H}(e) = \mathbf{H}(f) = [\{a, c\}, X]$, $\mathbf{I}(e) = \mathbf{I}(f) = [\{b, c\}, X]$,
 $\mathbf{J}(e) = \mathbf{J}(f) = [\emptyset, \{b, c\}]$.

Then we can easily check that (X, τ_2, E) is an $\text{IVST}_1(\text{i})$ -space and also an $\text{IVST}_1(\text{ii})$ -space

(3) Consider IVST τ_3 on X given by:

$$\tau_3 = \{\tilde{\mathcal{O}}_E, \mathbf{A}, \mathbf{B}, \mathbf{C}, \mathbf{D}, \mathbf{E}, \mathbf{F}, \mathbf{G}, \mathbf{H}, \tilde{X}_E\},$$

where $\mathbf{A}(e) = \mathbf{A}(f) = [\emptyset, \{a\}]$, $\mathbf{B}(e) = \mathbf{B}(f) = \{\{a\}, \{a\}\}$,
 $\mathbf{C}(e) = \mathbf{C}(f) = \{\{b\}, \{b\}\}$, $\mathbf{D}(e) = \mathbf{D}(f) = [\{c\}, \{a, c\}]$,
 $\mathbf{E}(e) = \mathbf{E}(f) = [\{a, b\}, \{a, b\}]$, $\mathbf{F}(e) = \mathbf{F}(f) = [\{a, c\}, \{a, c\}]$,
 $\mathbf{G}(e) = \mathbf{G}(f) = [\{b, c\}, X]$, $\mathbf{H}(e) = \mathbf{H}(f) = [\{b\}, \{a, b\}]$.

Observe that (X, τ_3, E) is an $\text{IVST}_0(\text{ii})$ -space but there are not $\mathbf{U}$, $\mathbf{U} \in \tau_3$ such that $a_0 \in \mathbf{U}$, $b_0 \notin \mathbf{U}$ and $b_0 \in \mathbf{V}$, $a_0 \notin \mathbf{V}$ for $a \neq b \in X$. Then X is not an $\text{IVST}_1(\text{ii})$ -space.

(4) Consider IVST τ_4 on X given by:

$$\tau_4 = \{\tilde{\mathcal{O}}_E, \mathbf{A}, \mathbf{B}, \mathbf{C}, \mathbf{D}, \mathbf{E}, \mathbf{F}, \mathbf{G}, \tilde{X}_E\},$$

where $\mathbf{A}(e) = \mathbf{A}(f) = [\emptyset, \{a\}]$, $\mathbf{B}(e) = \mathbf{B}(f) = \{\emptyset, \{b\}\}$,
 $\mathbf{C}(e) = \mathbf{C}(f) = \{\emptyset, \{c\}\}$, $\mathbf{D}(e) = \mathbf{D}(f) = [\emptyset, \{a, b\}]$,
 $\mathbf{E}(e) = \mathbf{E}(f) = [\emptyset, \{a, c\}]$, $\mathbf{F}(e) = \mathbf{F}(f) = [\emptyset, \{b, c\}]$,
 $\mathbf{G}(e) = \mathbf{G}(f) = [\emptyset, X]$.

Observe that (X, τ, E) is an $\text{IVST}_1(\text{ii})$ -space but there are not $\mathbf{U}$, $\mathbf{U} \in \tau_4$ such that $a_1 \in \mathbf{U}$, $b_1 \notin \mathbf{U}$ and $b_1 \in \mathbf{V}$, $a_1 \notin \mathbf{V}$ for $a \neq b \in X$. Then X is not an $\text{IVST}_1(\text{i})$ -space.

Remark 3.8. (1) If (X, τ, E) is an $\text{IVST}_0(\text{i})$ [resp. $\text{IVST}_1(\text{i})$]-space, then (X, τ^-, E) and (X, τ^+, E) are soft T_0 [resp. T_1]-spaces in the sense of Shabir and Naz [16].

(2) If (X, τ, E) is an $\text{IVST}_0(\text{ii})$ [resp. $\text{IVST}_1(\text{ii})$]-space, then (X, τ^+, E) is a soft T_0 [resp. T_1]-space in the sense of Shabir and Naz [16].

Definition 3.9. Let X be a set, E be a set of parameters and $x \in X$. An IVSS $\mathbf{x}_1$ [resp. $\mathbf{x}_0$] is defined by $\mathbf{x}_1(e) = x_1$ [resp. $\mathbf{x}_0(e) = x_0$] for each $e \in E$.

Proposition 3.10. Let (X, τ, E) be an IVSTS.

(1) If $\mathbf{x}_1$ is an IVSCS in X for each $x \in X$, then X is an $\text{IVST}_1(\text{i})$ -space.

(2) If $\mathbf{x}_0$ is an IVSCS in X for each $x \in X$, then X is an $\text{IVST}_1(\text{ii})$ -space.

Proof. (1) Suppose $\mathbf{x}_1$ is an IVSCS in X for each $x \in X$ and let $x \neq y \in X$. Then clearly, $\mathbf{x}_1^c = [X \setminus \{x\}, X \setminus \{x\}] \in \tau$ and $\mathbf{y}_1^c = [X \setminus \{y\}, X \setminus \{y\}] \in \tau$. Moreover, $x_1 \notin \mathbf{x}_1^c$, $y_1 \in \mathbf{x}_1^c$ and $y_1 \notin \mathbf{y}_1^c$, $x_1 \in \mathbf{y}_1^c$. Thus X is $\text{IVST}_1(\text{i})$ -space.

(2) The poof is similar to (1). □

Remark 3.11. The converses of Proposition 3.10 (1), (2) are not true in general (See Example 3.12).

Example 3.12. Let $X = \{x, y\}$ and $E = \{e, f\}$.

(1) Consider the collection τ_1 of IVSSs over X given by:

$$\tau_1 = \{\tilde{\mathcal{O}}_E, \mathbf{A}, \mathbf{B}, \mathbf{C}, \tilde{X}_E\},$$

where $\mathbf{A}(e) = \tilde{X}$, $\mathbf{A}(f) = [\{y\}, \{y\}]$, $\mathbf{B}(e) = [\{x\}, \{x\}]$, $\mathbf{B}(f) = \tilde{X}$,
 $\mathbf{C}(e) = [\{x\}, \{x\}]$, $\mathbf{C}(f) = [\{y\}, \{y\}]$.

Then clearly, (X, τ_1, E) is an IVSTS on X . Moreover, $\mathbf{A}, \mathbf{B} \in \tau$ such that $y_1 \in \mathbf{A}$, $x_1 \notin \mathbf{A}$ and $x_1 \in \mathbf{B}$, $y_1 \notin \mathbf{B}$. Thus (X, τ_1, E) is an $\text{IVST}_1(\text{i})$ -space.

On the other hand, since

$$\mathbf{x}_1(e) = \mathbf{x}_1(f) = x_1 = [\{x\}, \{x\}]$$

and

$$\mathbf{y}_1(e) = \mathbf{y}_1(f) = y_1 = [\{y\}, \{y\}],$$

we have

$$\mathbf{x}_1^c(e) = \mathbf{x}_1^c(f) = y_1 = [\{y\}, \{y\}]$$

and

$$\mathbf{y}_1^c(e) = \mathbf{y}_1^c(f) = x_1 = [\{x\}, \{x\}].$$

So $\mathbf{x}_1^c, \mathbf{y}_1^c \notin \tau_1$. Hence the converse of Proposition 3.10 (1) does not hold.

(2) Consider the collection τ_2 of IVSSs over X given by:

$$\tau_2 = \{\tilde{\mathcal{O}}_E, \mathbf{A}, \mathbf{B}, \mathbf{C}, \mathbf{D}, \tilde{X}_E\},$$

where $\mathbf{A}(e) = [\emptyset, X]$, $\mathbf{A}(f) = [\emptyset, \{y\}]$, $\mathbf{B}(e) = [\emptyset, \{x\}]$, $\mathbf{B}(f) = [\emptyset, X]$,
 $\mathbf{C}(e) = [\emptyset, \{x\}]$, $\mathbf{C}(f) = [\emptyset, \{y\}]$, $\mathbf{D}(e) = \mathbf{D}(f) = [\emptyset, X]$.

Then clearly, (X, τ_2, E) is an IVSTS on X . Moreover, $\mathbf{A}, \mathbf{B} \in \tau$ such that $y_0 \in \mathbf{A}$, $x_0 \notin \mathbf{A}$ and $x_0 \in \mathbf{B}$, $y_0 \notin \mathbf{B}$. Thus (X, τ_2, E) is an $\text{IVST}_1(\text{ii})$ -space.

On the other hand, since

$$\mathbf{x}_0(e) = \mathbf{x}_0(f) = x_0 = [\emptyset, \{x\}]$$

and

$$\mathbf{y}_0(e) = \mathbf{y}_0(f) = y_0 = [\emptyset, \{y\}],$$

we have

$$\mathbf{x}_1^c(e) = \mathbf{x}_1^c(f) = [\{y\}, X]$$

and

$$\mathbf{y}_1^c(e) = \mathbf{y}_1^c(f) = [\{x\}, X].$$

It follows that $\mathbf{x}_0^c, \mathbf{y}_0^c \notin \tau_2$. This means that the converse of Proposition 3.10 (2) does not hold.

Proposition 3.13. *Let (X, τ, E) be an IVSTS and $x \neq y \in X$.*

(1) *If there are $\mathbf{U}, \mathbf{V} \in \tau$ such that $x_1 \in \mathbf{U}$, $y_1 \in \mathbf{U}^c$ or $y_1 \in \mathbf{V}$, $x_1 \in \mathbf{V}^c$, then (X, τ, E) is an $\text{IVST}_0(\text{i})$ -space and for each $e \in E$, (X, τ_e) is an interval-valued $T_0(\text{i})$ -space in the sense of Lee et al. [43].*

(2) *If there are $\mathbf{U}, \mathbf{V} \in \tau$ such that $x_0 \in \mathbf{U}$, $y_0 \in \mathbf{U}^c$ or $y_0 \in \mathbf{V}$, $x_0 \in \mathbf{V}^c$, then (X, τ, E) is an $\text{IVST}_0(\text{ii})$ -space and for each $e \in E$, (X, τ_e) is an interval-valued $T_0(\text{ii})$ -space in the sense of Lee et al. [43].*

Proof. (1) Suppose there are $\mathbf{U}, \mathbf{V} \in \tau$ such that $x_1 \in \mathbf{U}$, $y_1 \in \mathbf{U}^c$ or $y_1 \in \mathbf{V}$, $x_1 \in \mathbf{V}^c$. If $y_1 \in \mathbf{U}^c$, then $y \in (U^-(e))^c$. Thus $y \notin U^-(e)$ for each $e \in E$. So $y_1 \notin \mathbf{U}$. Similarly, we can see that if $x_1 \in \mathbf{V}^c$, then $x_1 \notin \mathbf{V}$. Hence (X, τ, E) is an $\text{IVST}_0(\text{i})$ -space.

Now let $e \in E$. Then clearly, by Result 3.4, τ_e is an IVT on X . Since $x_1 \in \mathbf{U}$ and $y_1 \in \mathbf{U}^c$ or $y_1 \in \mathbf{V}$ and $x_1 \in \mathbf{V}^c$, $x_1 \in \mathbf{U}(e)$ and $y_1 \notin \mathbf{U}(e)$ or $y_1 \in \mathbf{V}(e)$ and $x_1 \notin \mathbf{V}(e)$. Thus (X, τ_e) is an interval-valued $T_0(\text{ii})$ -space.

(2) The proof is analogous to (1). $\square$

The following is an immediate consequence of Proposition 3.13 and Proposition 13 in [16].

Corollary 3.14. *Let (X, τ, E) be an IVSTS and $x \neq y \in X$.*

(1) *If there are $\mathbf{U}, \mathbf{V} \in \tau$ such that $x_1 \in \mathbf{U}$, $y_1 \in \mathbf{U}^c$ or $y_1 \in \mathbf{V}$, $x_1 \in \mathbf{V}^c$, then (X, τ_e^-) and (X, τ_e^+) are T_0 -spaces for each $e \in E$.*

(2) *If there are $\mathbf{U}, \mathbf{V} \in \tau$ such that $x_0 \in \mathbf{U}$, $y_0 \in \mathbf{U}^c$ or $y_0 \in \mathbf{V}$, $x_0 \in \mathbf{V}^c$, then (X, τ_e^+) is a T_0 -space for each $e \in E$.*

Proposition 3.15. *Let (X, τ, E) be an IVSTS and $x \neq y \in X$.*

(1) *If there are $\mathbf{U}, \mathbf{V} \in \tau$ such that $x_1 \in \mathbf{U}$, $y_1 \in \mathbf{U}^c$ and $y_1 \in \mathbf{V}$, $x_1 \in \mathbf{V}^c$, then (X, τ, E) is an $\text{IVST}_1(\text{i})$ -space and for each $e \in E$, (X, τ_e) is an interval-valued $T_1(\text{i})$ -space in the sense of Lee et al. [43].*

(2) *If there are $\mathbf{U}, \mathbf{V} \in \tau$ such that $x_0 \in \mathbf{U}$, $y_0 \in \mathbf{U}^c$ and $y_0 \in \mathbf{V}$, $x_0 \in \mathbf{V}^c$, then (X, τ, E) is an $\text{IVST}_0(\text{ii})$ -space and for each $e \in E$, (X, τ_e) is an interval-valued $T_1(\text{ii})$ -space in the sense of Lee et al. [43].*

Proof. The proofs are similar to Proposition 3.13. $\square$

The following is an immediate consequence of Proposition 3.15 and Proposition 14 in [16].

Corollary 3.16. *Let (X, τ, E) be an IVSTS and $x \neq y \in X$.*

(1) *If there are $\mathbf{U}, \mathbf{V} \in \tau$ such that $x_1 \in \mathbf{U}$, $y_1 \in \mathbf{U}^c$ and $y_1 \in \mathbf{V}$, $x_1 \in \mathbf{V}^c$, then (X, τ_e^-) and (X, τ_e^+) are T_1 -spaces for each $e \in E$.*

(2) *If there are $\mathbf{U}, \mathbf{V} \in \tau$ such that $x_0 \in \mathbf{U}$, $y_0 \in \mathbf{U}^c$ and $y_0 \in \mathbf{V}$, $x_0 \in \mathbf{V}^c$, then (X, τ_e^+) is an T_1 -space for each $e \in E$.*

Remark 3.17. (X, τ, E) is an $\text{IVST}_1(\text{i})$ [resp. $\text{IVST}_1(\text{ii})$]-space but (X, τ_e) may not be an interval-valued $T_1(\text{i})$ [resp. $T_1(\text{ii})$]-space (See Example 3.18).

Example 3.18. Let (X, τ_1, E) [resp. (X, τ_2, E)] be the $\text{IVST}_1(\text{i})$ [resp. $\text{IVST}_1(\text{ii})$]-space given in Example 3.12. Then we have

$$\tau_{1_e} = \{\tilde{\emptyset}, [\{x\}, \{x\}], \tilde{X}\}, \tau_{1_f} = \{\tilde{\emptyset}, [\{y\}, \{y\}], \tilde{X}\}$$

$$[\text{resp. } \tau_{2_e} = \{\tilde{\emptyset}, [\emptyset, X], [\emptyset, \{x\}], \tilde{X}\}, \tau_{2_f} = \{\tilde{\emptyset}, [\emptyset, X], [\emptyset, \{y\}], \tilde{X}\}].$$

Thus we can easily check that neither τ_{1_e} nor τ_{1_f} [resp. neither τ_{2_e} nor τ_{2_f}] is an interval-valued $T_1(\text{i})$ [resp. $T_1(\text{ii})$].

Definition 3.19. An IVSTS (X, τ, E) is called an:

(i) *interval-valued soft $T_2(\text{i})$ -space* (briefly, $\text{IVST}_2(\text{i})$ -space), if for any $x, y \in X$ with $x \neq y$, there are $\mathbf{U}, \mathbf{V} \in \tau$ such that $x_1 \in \mathbf{U}$, $y_1 \in \mathbf{V}$ and $\mathbf{U} \cap \mathbf{V} = \tilde{\emptyset}_E$,

(ii) *interval-valued soft T_2 (ii)-space* (briefly, $IVST_2$ (ii)-space), if for any $x, y \in X$ with $x \neq y$, there are $\mathbf{U}, \mathbf{V} \in \tau$ such that $x_0 \in \mathbf{U}$, $y_0 \in \mathbf{V}$ and $\mathbf{U} \cap \mathbf{V} = \tilde{\emptyset}_E$.

Remark 3.20. (1) Every $IVST_2$ (i) [resp. $IVST_2$ (ii)]-space is an $IVST_1$ (i) [resp. $IVST_1$ (ii)]-space. The converse is not true in general (See Example 3.21).

(2) Every $IVST_2$ (i)-space is an $IVST_2$ (ii)-space. The converse is not true in general (See Example 3.21).

Example 3.21. (1) Let (X, τ_1, E) [resp. (X, τ_2, E)] be the $IVST_1$ (i) [resp. $IVST_1$ (ii)]-space given as Example 3.12 (1) [resp. (2)]. Then we cannot have $\mathbf{U}, \mathbf{V} \in \tau_1$ such that $x_1 \in \mathbf{U}$, $y_1 \notin \mathbf{U}$ and $y_1 \in \mathbf{V}$, $x_1 \notin \mathbf{V}$ and $\mathbf{U} \cap \mathbf{V} = \tilde{\emptyset}_E$ [resp. $\mathbf{U}, \mathbf{V} \in \tau_2$ such that $x_0 \in \mathbf{U}$, $y_0 \notin \mathbf{U}$ and $y_0 \in \mathbf{V}$, $x_0 \notin \mathbf{V}$ and $\mathbf{U} \cap \mathbf{V} = \tilde{\emptyset}_E$]. Thus (X, τ_1, E) [resp. (X, τ_2, E)] is not an $IVST_2$ (i) [resp. $IVST_2$ (ii)]-space.

(2) Let $X = \{x, y\}$, $E = \{e, f\}$ and consider the $IVST$ τ given by:

$$\tau = \{\tilde{\emptyset}_E, \mathbf{A}, \mathbf{B}, \mathbf{C}, \tilde{X}_E\},$$

where $\mathbf{A}(e) = \mathbf{A}(f) = [\emptyset, \{x\}]$, $\mathbf{B}(e) = \mathbf{B}(f) = [\emptyset, \{y\}]$, $\mathbf{C}(e) = \mathbf{C}(f) = [\emptyset, X]$. Then observe that (X, τ, E) is an $IVST_2$ (ii)-space. On the other hand, there are not $\mathbf{U}, \mathbf{V} \in \tau_1$ such that $x_1 \in \mathbf{U}$, $y_1 \notin \mathbf{U}$ and $y_1 \in \mathbf{V}$, $x_1 \notin \mathbf{V}$ and $\mathbf{U} \cap \mathbf{V} = \tilde{\emptyset}_E$. Thus X is not an $IVST_2$ (i)-space.

Remark 3.22. (1) If (X, τ, E) is an $IVST_2$ (i)-space, then (X, τ^-, E) and (X, τ^+, E) are soft T_2 -spaces in the sense of Shabir and Naz [16].

(2) If (X, τ, E) is an $IVST_2$ (ii)-space, then (X, τ^+, E) is a soft T_2 -space in the sense of Shabir and Naz [16].

Proposition 3.23. Let (X, τ, E) be an $IVSTS$.

(1) If (X, τ, E) is an $IVST_2$ (i)-space, then for each $e \in E$, (X, τ_e) is an interval-valued T_2 (i)-space in the sense of Lee et al. [43].

(2) (X, τ, E) is an $IVST_2$ (ii)-space, then for each $e \in E$, (X, τ_e) is an interval-valued T_2 (ii)-space in the sense of Lee et al. [43].

Proof. (1) Suppose (X, τ, E) is an $IVST_2$ (i)-space, let $x \neq y \in X$ and $e \in E$. Then there are $\mathbf{U}, \mathbf{V} \in \tau$ such that $x_1 \in \mathbf{U}$, $y_1 \in \mathbf{V}$ and $\mathbf{U} \cap \mathbf{V} = \tilde{\emptyset}_E$. Thus $x_1 \in \mathbf{U}(e)$, $y_1 \in \mathbf{V}(e)$ and $\mathbf{U}(e) \cap \mathbf{V}(e) = \tilde{\emptyset}$. By Result 3.4, $\mathbf{U}(e), \mathbf{V}(e) \in \tau_e$. So (X, τ_e) is an interval-valued T_2 (i)-space.

(2) The proof is analogous to (1). $\square$

The following is an immediate consequence of Proposition 3.23 and Proposition 17 in [16].

Proposition 3.24. Let (X, τ, E) be an $IVSTS$.

(1) If (X, τ, E) is an $IVST_2$ (i)-space, then (X, τ_e^-) and (X, τ_e^+) are T_2 -space for each $e \in E$.

(2) (X, τ, E) is an $IVST_2$ (ii)-space, then (X, τ_e^+) is a T_2 -space for each $e \in E$.

Proof. (1) Suppose (X, τ, E) is an $IVST_2$ (i)-space, let $x \neq y \in X$ and let $e \in E$. Then there are $\mathbf{U}, \mathbf{V} \in \tau$ such that $x_1 \in \mathbf{U}$, $y_1 \in \mathbf{V}$ and $\mathbf{U} \cap \mathbf{V} = \tilde{\emptyset}_E$. Thus Definition 3.1 and Result 3.4, $x \in U^-(e)$, $y \in V^-(e)$, $U^-(e) \cap V^-(e) = \emptyset$ and $U^-(e), V^-(e) \in \tau_e^-$. So (X, τ_e^-) is a T_2 -space. Note that $\tau_e^- \subset \tau_e^+$. Hence (X, τ_e^+) is a T_2 -space.

(2) The proof is similar to (1). $\square$

Definition 3.25. Let (X, τ, E) be an IVSTS and $\mathbf{A}$ an IVSCS in X . Then (X, τ, E) is called:

- (i) an *interval-valued soft regular (i)-space* (briefly, IVSR(i)-space), if for each $x \in X$ and each $\mathbf{A} \in \tau^c$ with $x_1 \notin \mathbf{A}$, there are $\mathbf{U}, \mathbf{V} \in \tau$ such that $x_1 \in \mathbf{U}$, $\mathbf{A} \subset \mathbf{V}$ and $\mathbf{U} \cap \mathbf{V} = \tilde{\emptyset}_E$,
- (ii) an *interval-valued soft regular (ii)-space* (briefly, IVSR(ii)-space), if for each $x \in X$ and each $\mathbf{A} \in \tau^c$ with $x_0 \notin \mathbf{A}$, there are $\mathbf{U}, \mathbf{V} \in \tau$ such that $x_0 \in \mathbf{U}$, $\mathbf{A} \subset \mathbf{V}$ and $\mathbf{U} \cap \mathbf{V} = \tilde{\emptyset}_E$.

Definition 3.26. An IVSTS (X, τ, E) is called:

- (i) an *interval-valued soft T_3 (i)-space* (briefly, IVST₃(i)-space), if it is an IVSR(i) and IVST₁(i)-space,
- (ii) an *interval-valued soft T_3 (ii)-space* (briefly, IVST₃(ii)-space), if it is an IVSR(ii) and IVST₁(ii)-space.

Remark 3.27. (1) Every IVST₃(i)-space is an IVST₃(ii)-space but the converse is not true in general (See Example 3.28).

(2) An IVST₃(i) [resp. IVST₃(ii)]-space may not be an IVST₂(i) [resp. IVST₂(ii)]-space (See Example 3.28).

(3) If (X, τ, E) is an IVST₃(i) [resp. IVST₃(ii)]-space, then (X, τ_e) may not be an interval-valued T_3 (i) [resp. T_3 (ii)]-space for each $e \in E$ in the sense of Lee et al. [43] Furthermore, (X, τ_e^-) and (X, τ_e^+) may not be T_3 -spaces (See Example 3.28).

Example 3.28. (1) Let $X = \{a, b, c, d\}$, let $E = \{e, f\}$ and consider the IVST τ_1 on X given by:

$$\tau_1 = \{\tilde{\emptyset}_E, \mathbf{A}_1, \mathbf{A}_2, \dots, \mathbf{A}_{15}, \tilde{X}_E\},$$

where $\mathbf{A}_1(e) = \mathbf{A}_1(f) = [\{a, b\}, \{a, b, c\}]$, $\mathbf{A}_2(e) = \mathbf{A}_2(f) = [\{c, d\}, \{a, c, d\}]$,
 $\mathbf{A}_3(e) = \mathbf{A}_3(f) = [\{c, d\}, \{a, b, d\}]$, $\mathbf{A}_4(e) = \mathbf{A}_4(f) = [\{b, c\}, \{b, c, d\}]$,
 $\mathbf{A}_5(e) = \mathbf{A}_5(f) = [\emptyset, \{a, c\}]$, $\mathbf{A}_6(e) = \mathbf{A}_6(f) = [\{a\}, \{a, b\}]$,
 $\mathbf{A}_7(e) = \mathbf{A}_7(f) = [\{b\}, \{b, c\}]$, $\mathbf{A}_8(e) = \mathbf{A}_8(f) = [\emptyset, \{a, d\}]$,
 $\mathbf{A}_9(e) = \mathbf{A}_9(f) = [\{c\}, \{c, d\}]$, $\mathbf{A}_{10}(e) = \mathbf{A}_{10}(f) = [\emptyset, \{b, d\}]$,
 $\mathbf{A}_{11}(e) = \mathbf{A}_{11}(f) = [\{a, b, d\}, X]$, $\mathbf{A}_{12}(e) = \mathbf{A}_{12}(f) = [\{a, b, c\}, X]$,
 $\mathbf{A}_{13}(e) = \mathbf{A}_{13}(f) = [\{a, c, d\}, X]$, $\mathbf{A}_{14}(e) = \mathbf{A}_{14}(f) = [\{b, c, d\}, X]$,
 $\mathbf{A}_{15}(e) = \mathbf{A}_{15}(f) = [\{a\}, \{a, b\}]$.

Observe that (X, τ_1, E) is an IVST₃(ii)-space but not an IVST₃(i)-space.

Now consider the IVST τ_2 on X given by:

$$\tau_2 = \{\tilde{\emptyset}_E, \mathbf{A}_1, \mathbf{A}_2, \dots, \mathbf{A}_{17}, \tilde{X}_E\},$$

where $\mathbf{A}_1(e) = \mathbf{A}_1(f) = [\{a\}, \{a, d\}]$, $\mathbf{A}_2(e) = \mathbf{A}_2(f) = [\{b\}, \{a, b\}]$,
 $\mathbf{A}_3(e) = \mathbf{A}_3(f) = [\{c\}, \{b, c\}]$, $\mathbf{A}_4(e) = \mathbf{A}_4(f) = [\{d\}, \{c, d\}]$,
 $\mathbf{A}_5(e) = \mathbf{A}_5(f) = [\emptyset, \{a\}]$, $\mathbf{A}_6(e) = \mathbf{A}_6(f) = [\emptyset, \{b\}]$,
 $\mathbf{A}_7(e) = \mathbf{A}_7(f) = [\emptyset, \{c\}]$, $\mathbf{A}_8(e) = \mathbf{A}_8(f) = [\emptyset, \{d\}]$,
 $\mathbf{A}_9(e) = \mathbf{A}_9(f) = [\{a, b\}, \{a, b, d\}]$, $\mathbf{A}_{10}(e) = \mathbf{A}_{10}(f) = [\{a, d\}, \{a, c, d\}]$,
 $\mathbf{A}_{11}(e) = \mathbf{A}_{11}(f) = [\{c, d\}, \{b, c, d\}]$, $\mathbf{A}_{12}(e) = \mathbf{A}_{12}(f) = [\{b, c\}, \{a, b, c\}]$,
 $\mathbf{A}_{13}(e) = \mathbf{A}_{13}(f) = [\{a, c\}, X]$, $\mathbf{A}_{14}(e) = \mathbf{A}_{14}(f) = [\{b, d\}, X]$,
 $\mathbf{A}_{15}(e) = \mathbf{A}_{15}(f) = [\emptyset, \{a, b\}]$, $\mathbf{A}_{16}(e) = \mathbf{A}_{16}(f) = [\emptyset, \{a, c\}]$,

$$\mathbf{A}_{17}(e) = \mathbf{A}_{17}(f) = [\emptyset, \{a, d\}].$$

Then observe that (X, τ_2, E) is an $\text{IVST}_3(\text{i})$ -space.

(2) Let $X = \{a, b, c\}$, $E = \{e, f\}$ and consider the IVST τ on X given by:

$$\tau = \{\tilde{\mathcal{O}}_E, \mathbf{A}_1, \mathbf{A}_2, \dots, \mathbf{A}_{32}, \tilde{X}_E\},$$

where $\mathbf{A}_1(e) = \tilde{X}$, $\mathbf{A}_1(f) = \tilde{\mathcal{O}}$, $\mathbf{A}_2(e) = [\{a\}, \{a\}]$, $\mathbf{A}_2(f) = \tilde{\mathcal{O}}$,
 $\mathbf{A}_3(e) = [\{b\}, \{b\}]$, $\mathbf{A}_3(f) = \tilde{\mathcal{O}}$, $\mathbf{A}_4(e) = [\{c\}, \{c\}]$, $\mathbf{A}_4(f) = \tilde{\mathcal{O}}$,
 $\mathbf{A}_5(e) = [\{a, b\}, \{a, b\}]$, $\mathbf{A}_5(f) = \tilde{\mathcal{O}}$, $\mathbf{A}_6(e) = [\{a, c\}, \{a, c\}]$, $\mathbf{A}_6(f) = \tilde{\mathcal{O}}$,
 $\mathbf{A}_7(e) = [\{b, c\}, \{b, c\}]$, $\mathbf{A}_7(f) = \tilde{\mathcal{O}}$, $\mathbf{A}_8(e) = \tilde{X}$, $\mathbf{A}_8(f) = [\{a\}, \{a\}]$,
 $\mathbf{A}_9(e) = \mathbf{A}_9(f) = [\{a\}, \{a\}]$, $\mathbf{A}_{10}(e) = [\{b\}, \{b\}]$, $\mathbf{A}_{10}(f) = [\{a\}, \{a\}]$,
 $\mathbf{A}_{11}(e) = [\{c\}, \{c\}]$, $\mathbf{A}_{11}(f) = [\{a\}, \{a\}]$, $\mathbf{A}_{12}(e) = [\{a, b\}, \{a, b\}]$,
 $\mathbf{A}_{12}(f) = [\{a\}, \{a\}]$, $\mathbf{A}_{13}(e) = [\{a, c\}, \{a, c\}]$, $\mathbf{A}_{13}(f) = [\{a\}, \{a\}]$,
 $\mathbf{A}_{14}(e) = [\{b, c\}, \{b, c\}]$, $\mathbf{A}_{14}(f) = [\{a\}, \{a\}]$, $\mathbf{A}_{15}(e) = \tilde{\mathcal{O}}$,
 $\mathbf{A}_{15}(f) = [\{a\}, \{a\}]$, $\mathbf{A}_{16}(e) = \tilde{X}$, $\mathbf{A}_{16}(f) = [\{b, c\}, \{b, c\}]$,
 $\mathbf{A}_{17}(e) = [\{a\}, \{a\}]$, $\mathbf{A}_{17}(f) = [\{b, c\}, \{b, c\}]$, $\mathbf{A}_{18}(e) = [\{b\}, \{b\}]$,
 $\mathbf{A}_{18}(f) = [\{b, c\}, \{b, c\}]$, $\mathbf{A}_{19}(e) = [\{c\}, \{c\}]$, $\mathbf{A}_{19}(f) = [\{b, c\}, \{b, c\}]$,
 $\mathbf{A}_{20}(e) = [\{a, b\}, \{a, b\}]$, $\mathbf{A}_{20}(f) = [\{b, c\}, \{b, c\}]$, $\mathbf{A}_{21}(e) = [\{a, c\}, \{a, c\}]$,
 $\mathbf{A}_{21}(f) = [\{b, c\}, \{b, c\}]$, $\mathbf{A}_{22}(e) = [\{b, c\}, \{b, c\}]$, $\mathbf{A}_{22}(f) = [\{b, c\}, \{b, c\}]$,
 $\mathbf{A}_{23}(e) = \tilde{\mathcal{O}}$, $\mathbf{A}_{23}(f) = [\{b, c\}, \{b, c\}]$, $\mathbf{A}_{24}(e) = [\{a\}, \{a\}]$, $\mathbf{A}_{24}(f) = \tilde{X}$,
 $\mathbf{A}_{25}(e) = [\{b\}, \{b\}]$, $\mathbf{A}_{25}(f) = \tilde{X}$, $\mathbf{A}_{26}(e) = [\{c\}, \{c\}]$, $\mathbf{A}_{26}(f) = \tilde{X}$,
 $\mathbf{A}_{27}(e) = [\{a, b\}, X]$, $\mathbf{A}_{27}(f) = \tilde{X}$, $\mathbf{A}_{28}(e) = [\{a, c\}, X]$, $\mathbf{A}_{28}(f) = \tilde{X}$,
 $\mathbf{A}_{29}(e) = [\{b, c\}, X]$, $\mathbf{A}_{29}(f) = \tilde{X}$, $\mathbf{A}_{30}(e) = \tilde{\mathcal{O}}$, $\mathbf{A}_{30}(f) = \tilde{X}$.
 $\mathbf{A}_{31}(e) = \tilde{X}$, $\mathbf{A}_{31}(f) = \tilde{\mathcal{O}}$, $\mathbf{A}_{32}(e) = \tilde{X}$, $\mathbf{A}_{32}(f) = [\{a\}, \{a\}]$.

Then observe that (X, τ, E) is an $\text{IVST}_3(\text{i})$ -space but not an $\text{IVST}_2(\text{i})$ -space.

On the other hand, by Result 3.4, we have

$$\tau_e = \{\tilde{\mathcal{O}}, [\{a\}, \{a\}], [\{b\}, \{b\}], [\{c\}, \{c\}], [\{a, b\}, \{a, b\}], [\{a, c\}, \{a, c\}], [\{b, c\}, \{b, c\}], \tilde{X}\}$$

and

$$\tau_f = \{\tilde{\mathcal{O}}, [\{a\}, \{a\}], [\{b, c\}, \{b, c\}], \tilde{X}\}.$$

Obviously, (X, τ_f) is not an interval-valued $T_3(\text{i})$ -space. Moreover, we have

$$\tau_e^- = \tau_e^+ = \{\emptyset, \{a\}, \{b\}, \{c\}, \{a, b\}, \{a, c\}, \{b, c\}, X\}$$

and

$$\tau_f^- = \tau_f^+ = \{\emptyset, \{a\}, \{b, c\}, X\}.$$

Thus we can see that (X, τ_f^-) is not a T_3 -space.

Definition 3.29. An IVSTS (X, τ, E) is called an:

- (i) *interval-valued soft normal space* (briefly, IVSNS), If for any IVSCSs $\mathbf{F}_1, \mathbf{F}_2$ in X with $\mathbf{F}_1 \cap \mathbf{F}_2 = \tilde{\mathcal{O}}_E$, there are $\mathbf{U}, \mathbf{V} \in \tau$ such that $\mathbf{F}_1 \subset \mathbf{U}$, $\mathbf{F}_2 \subset \mathbf{V}$ and $\mathbf{U} \cap \mathbf{V} = \tilde{\mathcal{O}}_E$,
- (ii) *interval-valued soft $T_4(\text{i})$ -space* (briefly, $\text{IVST}_4(\text{i})$ -space), if it is an $T_1(\text{i})$ -space and an IVSNS,
- (iii) *interval-valued soft $T_4(\text{ii})$ -space* (briefly, $\text{IVST}_4(\text{ii})$ -space), if it is an $T_1(\text{ii})$ -space and an IVSNS.

Remark 3.30. (1) An $IVST_4(i)$ [resp. $IVST_4(ii)$]-space need not an $IVST_3(i)$ [resp. $IVST_3(ii)$]-space (See Example 3.31).

(2) If (X, τ, E) is an $IVST_4(i)$ [resp. $IVST_4(ii)$]-space, then (X, τ_e) may not be an interval-valued T_4 -space for each $e \in E$ in the sense of Lee et al. [43] Furthermore, (X, τ_e^-) and (X, τ_e^+) may not be T_4 -spaces (See Example 3.31).

Example 3.31. Let $X = \{a, b, c, d\}$, let $E = \{e, f\}$ and consider the IVST τ on X given by:

$$\tau = \{\tilde{\varnothing}_E, \mathbf{A}_1, \mathbf{A}_2, \mathbf{A}_3, \mathbf{A}_4, \mathbf{A}_5, \mathbf{A}_6, \mathbf{A}_7, \mathbf{A}_8, \tilde{X}_E\},$$

$$\begin{aligned} \text{where } \mathbf{A}_1(e) &= [\{a, b, d\}, \{a, b, d\}], \mathbf{A}_1(f) = [\{a, b, c\}, \{a, b, c\}], \\ \mathbf{A}_2(e) &= [\{a, c, d\}, \{a, c, d\}], \mathbf{A}_2(f) = [\{a, b, c\}, \{a, b, c\}], \\ \mathbf{A}_3(e) &= [\{a, d\}, \{a, d\}], \mathbf{A}_3(f) = [\{a, b, c\}, \{a, b, c\}], \\ \mathbf{A}_4(e) &= [\{b, c\}, \{b, c\}], \mathbf{A}_4(f) = [\{a, b, c\}, \{a, b, c\}], \\ \mathbf{A}_5(e) &= [\{b\}, \{b\}], \mathbf{A}_5(f) = [\{a, b, c\}, \{a, b, c\}], \\ \mathbf{A}_6(e) &= [\{c\}, \{c\}], \mathbf{A}_6(f) = [\{a, b, c\}, \{a, b, c\}], \\ \mathbf{A}_7(e) &= \tilde{\varnothing}, \mathbf{A}_7(f) = [\{a, b, c\}, \{a, b, c\}], \\ \mathbf{A}_8(e) &= \tilde{X}, \mathbf{A}_8(f) = [\{a, b, c\}, \{a, b, c\}]. \end{aligned}$$

Then we can check that (X, τ, E) is an $IVST_4(i)$ -space but not an $IVST_3(i)$ -space.

On the other hand, by Result 3.4, we have

$$\begin{aligned} \tau_e &= \{\tilde{\varnothing}, [\{b\}, \{b\}], [\{c\}, \{c\}], [\{a, d\}, \{a, d\}], [\{b, c\}, \{b, c\}], \\ &\quad [\{a, c, d\}, \{a, c, d\}], [\{a, b, d\}, \{a, b, d\}], \tilde{X}\} \end{aligned}$$

and

$$\tau_f = \{\tilde{\varnothing}, [\{a, b, c\}, \{a, b, c\}], \tilde{X}\}.$$

Then we can easily see that (X, τ_e) and (X, τ_f) are not interval-valued T_4 -spaces. Furthermore, we get

$$\tau_e^- = \tau_e^+ = \{\varnothing, \{b\}, \{c\}, \{a, d\}, \{b, c\}, \{a, c, d\}, \{a, b, d\}, X\}$$

and

$$\tau_f^- = \tau_f^+ = \{\varnothing, \{a, b, c\}, X\}.$$

Thus we can check that (X, τ_e^-) and (X, τ_f^-) are not T_4 -spaces.

4. INTERVAL-VALUED SOFT SUBSPACES

In this section, we deal with some hereditary problems in an IVSTS. For this reason, we introduce the notion of interval-valued soft subspaces of an IVSTS.

Definition 4.1. Let Y be a nonempty subset of X and $\mathbf{A} \in IVSS_E(X)$. Then

(i) the *interval valued soft set* (Y, E) over X , denoted by $\tilde{Y}_E$, is defined as follows:

$$\tilde{Y}_E(e) = [Y, Y] \text{ for each } e \in E,$$

(ii) the *interval-valued soft subset of $\mathbf{A}$ over Y* , denoted by $\mathbf{A}_Y$, is defined as follows:

$$\mathbf{A}_Y = \tilde{Y}_E \cap \mathbf{A}, \text{ i.e., } \mathbf{A}_Y(e) = [Y \cap A^-(e), Y \cap A^+(e)] \text{ for each } e \in E.$$

Example 4.2. Let $X = \{a, b, c, d\}$, $Y = \{a, b, c\}$, $E = \{e, f\}$ and $\mathbf{A}$ be the IVSS over X given by:

$$\mathbf{A}(e) = [\{a, b\}, \{a, b, d\}], \mathbf{A}(f) = [\{a, d\}, \{a, c, d\}].$$

Then clearly, $\mathbf{A}_Y(e) = [\{a, b\}, \{a, b\}]$, $\mathbf{A}_Y(f) = [\{a\}, \{a, c\}]$.

Proposition 4.3. *Let (X, τ, E) be an IVSTS and Y a nonempty subset of X . Then $\tau_Y = \{\mathbf{A}_Y : \mathbf{A} \in \tau\}$ is an IVST on Y .*

In this case, τ_Y is called the *interval-valued soft relative topology* on Y and (Y, τ_Y, E) is called an *interval-valued soft subspace* (briefly, IVS-subspace) of (X, τ, E) . Each member of τ_Y is called an *IVSOS in Y* and an IVSS $\mathbf{A}$ over Y is called an *IVSCS in Y* , if $[Y, Y] \setminus \mathbf{A} = [Y \setminus A^+, Y \setminus A^-] \in \tau_Y$.

Proof. The proof is straightforward. $\square$

We can see that (Y, τ_Y, E) is a special interval-valued soft subspace of an IVSTS (X, τ, E) (See Proposition 4.15, [34]).

Remark 4.4. Every IVS-subspace of an interval-valued soft discrete [resp. indiscrete] space is an interval-valued soft discrete [resp. indiscrete] space.

Lemma 4.5. *Let (X, τ) be an IVTS and Y a nonempty subset of X . Then $\tau_{[Y, Y]} = \{[Y, Y] \cap \mathbf{A} : \mathbf{A} \in \tau\} = \{[Y \cap A^-, [Y \cap A^+] : \mathbf{A} \in \tau\}$ is an IVT on Y .*

In this case, $\tau_{[Y, Y]}$ is called the *interval-valued relative topology* on Y and $([Y, Y], \tau_{[Y, Y]})$ is called an *interval-valued subspace* (briefly, IV-subspace) of (X, τ) .

Proof. The proof is straightforward. $\square$

Proposition 4.6. *Let (X, τ, E) be an IVSTS and Y a nonempty subset of X . Then $(Y, (\tau_Y)_e) = ([Y, Y], \tau_{e, [Y, Y]})$, i.e., $(Y, (\tau_Y)_e)$ is an interval-valued subspace of IVTS (X, τ_e) for each $e \in E$, where $\tau_{e, [Y, Y]} = \{[Y, Y] \cap \mathbf{A}(e) : \mathbf{A}(e) \in \tau_e\}$.*

Proof. The proof follows from Proposition 4.3 and Lemma 4.5. $\square$

Corollary 4.7 (See Proposition 11, [16]). *Let (X, τ, E) be an IVSTS and Y a nonempty subset of X . Then $(Y, (\tau_Y)_e^-) = (Y, (\tau_e^-)_Y)$ [resp. $(Y, (\tau_Y)_e^+) = (Y, (\tau_e^+)_Y)$], i.e., $(Y, (\tau_Y)_e^-)$ [resp. $(Y, (\tau_Y)_e^+)$] is a subspace of (X, τ_e^-) [resp. (X, τ_e^+)] for each $e \in E$.*

Proposition 4.8. *Let (Y, τ_Y, E) be an IVS-subspace of an IVSTS (X, τ, E) and $\mathbf{A}$ an IVSOS in Y . If $\tilde{Y}_E \in \tau$, then $\mathbf{A} \in \tau$.*

Proof. The proof is obvious. $\square$

Theorem 4.9. *Let (Y, τ_Y, E) be an IVS-subspace of an IVSTS (X, τ, E) and $\mathbf{A} \in IVSS_E(X)$.*

(1) $\mathbf{A} \in \tau_Y$ if and only if there is $\mathbf{B} \in \tau$ such $\mathbf{A} = \tilde{Y}_E \cap \mathbf{B}$.

(2) $\mathbf{A} \in \tau_Y^c$ if and only if there is $\mathbf{B} \in \tau^c$ such $\mathbf{A} = \tilde{Y}_E \cap \mathbf{B}$.

Proof. (1) The proof follows from Proposition 4.3.

(2) Suppose $\mathbf{A} \in \tau_Y^c$. Then clearly, $[Y, Y] - \mathbf{A} = [Y - A^+, Y - A^-] \in \tau_Y$. Thus by (1), there is $\mathbf{B} \in \tau$ such that $[Y - A^+, Y - A^-] = \tilde{Y}_E \cap \mathbf{B}$. Now let $e \in E$. Then

$$[Y - A^+, Y - A^-](e) = \tilde{Y}_E \cap \mathbf{B}(e), \text{ i.e.,}$$

$$[Y - A^+(e), Y - A^-(e)] = [Y \cap B^-(e), Y \cap B^-(e)], \text{ i.e.,}$$

$$Y - A^+(e) = Y \cap B^-(e), Y - A^-(e) = Y \cap B^-(e).$$

Thus we have

$$A^+(e) = Y - (Y - A^+(e)) = Y - (Y \cap B^-(e)) = Y \cap (Y - B^-(e)) = Y \cap (X - B^-(e)),$$

$$A^-(e) = Y - (Y - A^-(e)) = Y - (Y \cap B^+(e)) = Y \cap (Y - B^+(e)) = Y \cap (X - B^+(e)).$$

$$\text{So } \mathbf{A}(e) = [A^-(e), A^+(e)] = [Y \cap (X - B^+(e)), Y \cap (X - B^-(e))] = (\tilde{Y}_E \cap \mathbf{B}^c)(e).$$

Hence $\mathbf{A} = \tilde{Y}_E \cap \mathbf{B}^c$. Since $\mathbf{B} \in \tau$, $\mathbf{B}^c \in \tau^c$. Therefore the necessary condition holds.

Conversely, suppose the necessary condition holds, i.e., there is $\mathbf{B} \in \tau^c$ such that $\mathbf{A} = \tilde{Y}_E \cap \mathbf{B}$. Then $\mathbf{B}^c = \tilde{X}_E - \mathbf{B} \in \tau$. We will show that $[Y, Y] - \mathbf{A} = \tilde{Y}_E \cap \mathbf{B}^c$. Let $e \in E$. Then we get

$$\begin{aligned} ([Y, Y] - \mathbf{A})(e) &= [Y - A^+(e), Y - A^-(e)] \\ &= [Y - Y \cap B^-(e), Y - Y \cap B^+(e)] \\ &= [Y \cap (Y - B^-(e)), Y \cap (Y - B^+(e))] \\ &= [Y \cap (X - B^+(e)), Y \cap (X - B^-(e))]. \end{aligned}$$

Thus $[Y, Y] - \mathbf{A} = \tilde{Y}_E \cap (\tilde{X}_E - \mathbf{B}) = \tilde{Y}_E \cap \mathbf{B}^c$. Since $\mathbf{B}^c \in \tau$, we have $[Y, Y] - \mathbf{A} \in \tau_Y$. So $\mathbf{A} \in \tau_Y^c$. This completes the proof. $\square$

Proposition 4.10. *Let (X, τ, E) be an IVSTS and Y a nonempty subset of X .*

(1) *If X is an $IVST_0$ (i) [resp. $IVST_0$ (ii)]-space, then (Y, τ_Y, E) is an $IVST_0$ (i) [resp. $IVST_0$ (ii)]-space.*

(2) *If X is an $IVST_1$ (i) [resp. $IVST_1$ (ii)]-space, then (Y, τ_Y, E) is an $IVST_1$ (i) [resp. $IVST_1$ (ii)]-space.*

Proof. (1) Suppose X is an $IVST_0$ (i)-space and let $x \neq y \in Y$. Then there are $\mathbf{U}, \mathbf{V} \in \tau$ such that either $x_1 \in \mathbf{U}, y_1 \notin \mathbf{U}$ or $y_1 \in \mathbf{U}, x_1 \notin \mathbf{U}$, say $x_1 \in \mathbf{U}, y_1 \notin \mathbf{U}$. Since $x \in Y, x_1 \in \tilde{Y}_E$. Since $x_1 \in \mathbf{U}$, we have $x_1 \in \tilde{Y}_E \cap \mathbf{U} = \mathbf{U}_Y$. Since $y_1 \notin \mathbf{U}$, there is $e \in E$ such that $y \notin \mathbf{U}(e)$, i.e., $y \notin U^-(e)$. Thus $y \notin Y \cap U^-(e) = (Y \cap U^-)(e)$. So $y_1 \notin \tilde{Y}_E \cap \mathbf{U} = \mathbf{U}_Y$. Since $\mathbf{U} \in \tau, \mathbf{U}_Y \in \tau_Y$. Hence (Y, τ_Y, E) is an $IVST_1$ (i)-space. The proof of the second part is similar.

(2) The proof is similar to (1). $\square$

Proposition 4.11. *Let (X, τ, E) be an IVSTS and Y a nonempty subset of X . If X is an $IVST_2$ (i) [resp. $IVST_2$ (ii)]-space, then (Y, τ_Y, E) is an $IVST_2$ (i) [resp. $IVST_2$ (ii)]-space.*

Proof. Suppose X is an $IVST_2$ (i)-space and let $x \neq y \in Y$. Then there are $\mathbf{U}, \mathbf{V} \in \tau$ such that either $x_1 \in \mathbf{U}, y_1 \in \mathbf{V}, \mathbf{U} \cap \mathbf{V} = \tilde{X}_E$. Then clearly, we have

$$x \in U^-(e), y \in V^-(e), U^+(e) \cap V^+(e) = \emptyset.$$

Thus $x \in Y \cap U^-(e), y \in Y \cap V^-(e), (Y \cap U^+(e)) \cap (Y \cap V^+(e)) = \emptyset$. So we get

$$x \in U_Y^-(e), y \in V_Y^-(e), U_Y^+(e) \cap V_Y^+(e) = \emptyset.$$

Hence $x_1 \in \mathbf{U}_Y, y_1 \in \mathbf{V}_Y, \mathbf{U}_Y \cap \mathbf{V}_Y = \tilde{\mathcal{Q}}_E$. Therefore (Y, τ_Y, E) is an $IVST_2$ (i)-space. The proof of the second part is similar. $\square$

Proposition 4.12. *Let (X, τ, E) be an IVSTS and Y a nonempty subset of X . If X is an $IVST_3(i)$ [resp. $IVST_3(ii)$]-space, then (Y, τ_Y, E) is an $IVST_3(i)$ [resp. $IVST_3(ii)$]-space.*

Proof. Suppose X is an $IVST_3(i)$ -space. Then by Proposition 4.10 (2), (X, τ_Y, E) is an $IVST_1(i)$ -space. Let $y \in Y$ and $\mathbf{A} \in \tau_Y^c$ such that $y_1 \notin \mathbf{A}$. Then by Theorem 4.9 (2), there is $\mathbf{B} \in \tau^c$ such that $\mathbf{A} = \tilde{Y}_E \cap \mathbf{B}$. Since $y_1 \notin \mathbf{A}$, $y_1 \notin \tilde{Y}_E \cap \mathbf{B}$. But $y_1 \in \tilde{Y}_E$. Thus $y_1 \notin \mathbf{B}$. By the hypothesis, there are $\mathbf{U}, \mathbf{V} \in \tau$ such that $y_1 \in \mathbf{U}$, $\mathbf{B} \subset \mathbf{V}$ and $\mathbf{U} \cap \mathbf{V} = \tilde{\mathcal{O}}_E$. So by Proposition 4.10 (1), $\tilde{Y}_E \cap \mathbf{U}, \tilde{Y}_E \cap \mathbf{V} \in \tau_Y$. Furthermore, $y_1 \notin \tilde{Y}_E \cap \mathbf{U}$, $\mathbf{A} \subset \tilde{Y}_E \cap \mathbf{V}$ and $(\tilde{Y}_E \cap \mathbf{U}) \cap (\tilde{Y}_E \cap \mathbf{V}) = \tilde{\mathcal{O}}_E$. Hence (Y, τ_Y, E) is an $IVST_3(i)$ -space. The proof of the second part is similar. $\square$

Remark 4.13. Let (X, τ, E) be an IVSTS and let Y be a nonempty subset of X . When (X, τ, E) is an $IVST_4(i)$ [resp. $IVST_4(ii)$]-space and Y is a nonempty subset of X , (Y, τ_Y, E) need not be an $IVST_3(i)$ [resp. $IVST_3(ii)$]-space (See Example 4.14).

Example 4.14. Consider the $IVST_4(i)$ -space (X, τ, E) given in Example 3.31 and let $Y = \{a, b, c\}$. Then clearly, we have

$$\tau_Y = \{\tilde{\mathcal{O}}_E, \mathbf{A}_{1_Y}, \mathbf{A}_{2_Y}, \mathbf{A}_{3_Y}, \mathbf{A}_{4_Y}, \mathbf{A}_{5_Y}, \mathbf{A}_{6_Y}, \mathbf{A}_{7_Y}, \mathbf{A}_{7_Y}, \tilde{Y}_E\},$$

where $\mathbf{A}_{1_Y}(e) = [\{a, b\}, \{a, b\}]$, $\mathbf{A}_{1_Y}(f) = [Y, Y]$,
 $\mathbf{A}_{2_Y}(e) = [\{a, c\}, \{a, c\}]$, $\mathbf{A}_{2_Y}(f) = [Y, Y]$,
 $\mathbf{A}_{3_Y}(e) = [\{a\}, \{a\}]$, $\mathbf{A}_{3_Y}(f) = [Y, Y]$,
 $\mathbf{A}_{4_Y}(e) = [\{b, c\}, \{b, c\}]$, $\mathbf{A}_{4_Y}(f) = [Y, Y]$,
 $\mathbf{A}_{5_Y}(e) = [\{b\}, \{b\}]$, $\mathbf{A}_{5_Y}(f) = [Y, Y]$,
 $\mathbf{A}_{6_Y}(e) = [\{c\}, \{c\}]$, $\mathbf{A}_{6_Y}(f) = [Y, Y]$,
 $\mathbf{A}_{7_Y}(e) = \tilde{\mathcal{O}}$, $\mathbf{A}_{7_Y}(f) = [Y, Y]$,
 $\mathbf{A}_{8_Y}(e) = \tilde{X}$, $\mathbf{A}_{7_Y}(f) = [Y, Y]$.

Thus we can check that (Y, τ_Y, E) is not an $IVST_3(i)$ -space.

5. INTERVAL-VALUED SOFT α -CLOSURES [RESP. INTERIORS]

In this section, we define an interval-valued α -closure [resp. interior] of an interval-valued soft set and discuss some of their properties, and give some examples.

Definition 5.1. Let (X, τ, E) be a soft topological space over X and $A, B \in SS_E(X)$.

(i) The *soft closure* of A [16], denoted by $cl_\tau(A)$ or $cl(A)$ or $\bar{A}$, is a soft set over X defined as follows:

$$cl_\tau(A) = \bigcap \{F \in \tau^c : A \subset F\}.$$

(ii) The *soft interior* of A [20], denoted by $int_\tau(A)$ or $int(A)$ or A° , is a soft set over X defined as follows:

$$int_\tau(A) = \bigcup \{U \in \tau : U \subset A\}.$$

It is obvious that $cl(A)$ is the smallest soft closed set in X which contains A and $int(A)$ is the largest soft open set in X which is contained in A .

Definition 5.2 ([34]). Let (X, τ, E) be an IVSTS and $\mathbf{A} \in IVSS_E(X)$.

(i) The *interval-valued soft closure* of $\mathbf{A}$ w.r.t. τ , denoted by $IVScl(\mathbf{A})$, is an IVSS over X defined as:

$$IVScl(\mathbf{A}) = \bigcap \{\mathbf{K} \in \tau^c : \mathbf{A} \subset \mathbf{K}\}.$$

(ii) The *interval-valued soft interior* of $\mathbf{A}$ w.r.t. τ , denoted by $IVSint(\mathbf{A})$, is an IVSS over X defined as:

$$IVSint(\mathbf{A}) = \bigcup \{\mathbf{U} : \mathbf{U} \in \tau \text{ and } \mathbf{U} \subset \mathbf{A}\}.$$

We can see that $IVScl(\mathbf{A})$ is the smallest IVSCS in X containing $\mathbf{A}$ and $IVSint(\mathbf{A})$ is the largest IVSOS in X contained $\mathbf{A}$.

Remark 5.3. $IVScl(\mathbf{A}) = [cl_{\tau^-}(A^-), cl_{\tau^+}(A^+)]$, $IVSint(\mathbf{A}) = [int_{\tau^-}(A^-), int_{\tau^+}(A^+)]$.

Example 5.4. Let $X = \{a, b, c\}$, $E = \{e, f\}$ and consider τ the IVST on X given by:

$$\tau = \{\tilde{\varnothing}_E, \mathbf{U}, \tilde{X}_E\},$$

where $\mathbf{U}(e) = [\{a\}, \{a, b\}]$, $\mathbf{U}(f) = [\{b\}, \{b, c\}]$. Then we have

$$\tau^- = \{\varnothing_E, U^-, X_E\}, \quad \tau^+ = \{\varnothing_E, U^+, X_E\},$$

where $U^-(e) = \{a\}$, $U^-(f) = \{b\}$, $U^+(e) = \{a, b\}$, $U^+(f) = \{b, c\}$. Moreover,

$$\tau^c = \{\tilde{\varnothing}_E, \mathbf{U}^c, \tilde{X}_E\},$$

where $\mathbf{U}^c(e) = [\{c\}, \{b, c\}]$, $\mathbf{U}^c(f) = [\{a\}, \{a, c\}]$.

Consider the IVSS $\mathbf{A}$ over X defined by $\mathbf{A}(e) = [\{c\}, \{c\}]$, $\mathbf{A}(f) = [\{a\}, \{a\}]$. Then clearly, $\mathbf{A} \subset \mathbf{U}^c$ and $\mathbf{A} \subset \tilde{X}_E$. Thus $cl(\mathbf{A}) = \mathbf{U}^c \cap \tilde{X}_E = \mathbf{U}^c$. On the other hand, we get

$$cl_{\tau^-}(A^-) = U^{c,-}, \quad cl_{\tau^+}(A^+) = U^{c,+},$$

where $U^{c,-}(e) = \{c\}$, $U^{c,-}(f) = \{a\}$, $U^{c,+}(e) = \{b, c\}$, $U^{c,+}(f) = \{a, c\}$. So $cl(\mathbf{A}) = [cl_{\tau^-}(A^-), cl_{\tau^+}(A^+)]$.

Definition 5.5 ([36]). Let (X, τ, E) be a soft topological space and $A \in SS_E(X)$. A is called a *soft α -open set* in X , if $A \subset int(cl(int(A)))$. The complement of a soft α -open set is called a *soft α -closed set* in X .

The set of all soft α -open [resp. closed] sets in a soft topological space (X, τ, E) is denoted by $S\alpha OS(X)$ [resp. $S\alpha CS(X)$].

Definition 5.6. (i) Let (X, τ, E) be an IVSTS and $\mathbf{A} \in IVSS_E(X)$. Then $\mathbf{A}$ is called an *interval-valued soft α -open set* (briefly, $IVS\alpha OS$) in X , if it satisfies the following conditions:

$$\mathbf{A} \subset IVSint(IVScl(IVSint(\mathbf{A}))).$$

The complement of an $IVS\alpha OS$ is called an *interval-valued soft α -closed set* (briefly, $IVS\alpha CS$) in X .

(ii) Let (X, τ) be an IVTS and let $A \in IVS(X)$. Then A is called an *interval-valued α -open set* (briefly, $IV\alpha OS$) in X , if $A \subset IVint(IVcl(IVint(A)))$, where $IVint(A)$ and $IVcl(A)$ denote the interval-valued interior and the interval-valued closure of A (See [37]). The complement of an $IV\alpha OS$ is called an *interval-valued α -closed set* (briefly, $IV\alpha CS$) in X .

We will denote the set of all IVS α OSs [resp. IVS α CS] by IVS α OS(X) [resp. IVS α CS(X)]. Also, We will denote the set of all IV α OSs [resp. IV α CS] by IV α OS(X) [resp. IV α CS(X)].

From Remark 5.3, it is obvious that $\mathbf{A} \in IVS\alpha OS(X)$, then $A^- \in S\alpha OS(X, \tau^-, E)$ and $A^+ \in S\alpha OS(X, \tau^+, E)$.

Example 5.7. Let (X, τ, E) be the IVSTS given in Example 5.4. Observe that $\mathbf{U} \in IVS\alpha OS(X)$. Furthermore we can confirm that $U^- \in S\alpha OS(X, \tau^-, E)$ and $U^+ \in S\alpha OS(X, \tau^+, E)$.

Proposition 5.8. Let (X, τ, E) be an IVSTS. Then

- (1) $\bigcup_{j \in J} \mathbf{A}_j \in IVS\alpha OS(X)$ for each $(\mathbf{A}_j)_{j \in J} \subset IVS\alpha OS(X)$,
- (2) $\bigcap_{j \in J} \mathbf{A}_j \in IVS\alpha CS(X)$ for each $(\mathbf{A}_j)_{j \in J} \subset IVS\alpha CS(X)$.

Proof. The proofs are straightforward. $\square$

Remark 5.9. In an IVSTS (X, τ, E) , $\tau \subset IVS\alpha OS(X)$ and $\tau^c \subset IVS\alpha CS(X)$. But the converse need not be true (See Example 5.10)

Example 5.10. Let $X = \{a, b, c, d\}$, $E = \{e, f, g\}$ and consider the IVST τ given by:

$$\tau = \{\tilde{\mathcal{O}}_E, \mathbf{A}_1, \mathbf{A}_2, \dots, \mathbf{A}_{17}, \tilde{X}_E\},$$

where $\mathbf{A}_1(e) = [\{a\}, \{a\}]$, $\mathbf{A}_1(f) = [\{b, c\}, \{b, c\}]$, $\mathbf{A}_1(g) = [\{a, d\}, \{a, d\}]$,
 $\mathbf{A}_2(e) = [\{b, d\}, \{b, d\}]$, $\mathbf{A}_2(f) = [\{a, c, d\}, \{a, c, d\}]$, $\mathbf{A}_2(g) = [\{a, b, d\}, \{a, b, d\}]$,
 $\mathbf{A}_3(e) = [\emptyset, \emptyset]$, $\mathbf{A}_3(f) = [\{c\}, \{c\}]$, $\mathbf{A}_3(g) = [\{a\}, \{a\}]$,
 $\mathbf{A}_4(e) = [\{a, b, d\}, \{a, b, d\}]$, $\mathbf{A}_4(f) = [X, X]$, $\mathbf{A}_4(g) = [X, X]$,
 $\mathbf{A}_5(e) = [\{a, c\}, \{a, c\}]$, $\mathbf{A}_5(f) = [\{b, d\}, \{b, d\}]$, $\mathbf{A}_5(g) = [\{b\}, \{b\}]$,
 $\mathbf{A}_6(e) = [\{a\}, \{a\}]$, $\mathbf{A}_6(f) = [\{b\}, \{b\}]$, $\mathbf{A}_6(g) = [\emptyset, \emptyset]$,
 $\mathbf{A}_7(e) = [\{a, c\}, \{a, c\}]$, $\mathbf{A}_7(f) = [\{b, d, d\}, \{b, c, d\}]$, $\mathbf{A}_7(g) = [\{a, b, d\}, \{a, b, d\}]$,
 $\mathbf{A}_8(e) = [\emptyset, \emptyset]$, $\mathbf{A}_8(f) = [\{d\}, \{d\}]$, $\mathbf{A}_8(g) = [\{b\}, \{b\}]$,
 $\mathbf{A}_9(e) = [X, X]$, $\mathbf{A}_9(f) = [X, X]$, $\mathbf{A}_9(g) = [\{a, b, c\}, \{a, b, c\}]$,
 $\mathbf{A}_{10}(e) = [\{a, c\}, \{a, c\}]$, $\mathbf{A}_{10}(f) = [\{b, c, d\}, \{b, c, d\}]$, $\mathbf{A}_{10}(g) = [\{a, b\}, \{a, b\}]$,
 $\mathbf{A}_{11}(e) = [\{b, c, d\}, \{b, c, d\}]$, $\mathbf{A}_{11}(f) = [X, X]$, $\mathbf{A}_{11}(g) = [\{a, b, c\}, \{a, b, c\}]$,
 $\mathbf{A}_{12}(e) = [\{a\}, \{a\}]$, $\mathbf{A}_{12}(f) = [\{b, c, d\}, \{b, c, d\}]$, $\mathbf{A}_{12}(g) = [\{a, b, d\}, \{a, b, d\}]$,
 $\mathbf{A}_{13}(e) = [\{a\}, \{a\}]$, $\mathbf{A}_{13}(f) = [\{b, d\}, \{b, d\}]$, $\mathbf{A}_{13}(g) = [\{b\}, \{b\}]$,
 $\mathbf{A}_{14}(e) = [\{c, d\}, \{c, d\}]$, $\mathbf{A}_{14}(f) = [\{a, b\}, \{a, b\}]$, $\mathbf{A}_{14}(g) = [\emptyset, \emptyset]$,
 $\mathbf{A}_{15}(e) = [\{a\}, \{a\}]$, $\mathbf{A}_{15}(f) = [\{b, c\}, \{b, c\}]$, $\mathbf{A}_{15}(g) = [\{a\}, \{a\}]$,
 $\mathbf{A}_{16}(e) = [\emptyset, \emptyset]$, $\mathbf{A}_{16}(f) = [\{c\}, \{c\}]$, $\mathbf{A}_{16}(g) = [\{a, d\}, \{a, d\}]$,
 $\mathbf{A}_{17}(e) = [\{a, b, d\}, \{a, b, d\}]$, $\mathbf{A}_{17}(f) = [X, X]$, $\mathbf{A}_{17}(g) = [\{a, b, d\}, \{a, b, d\}]$.

Then we have

$$\tau^c = \{\tilde{\mathcal{O}}_E, \mathbf{A}_1^c, \mathbf{A}_2^c, \dots, \mathbf{A}_{15}^c, \tilde{X}_E\}.$$

Let $\mathbf{A}$ be the IVSS over X defined by:

$$\mathbf{A}(e) = [\{a, b, c\}, \{a, b, c\}], \mathbf{A}(f) = [\{b, c, d\}, \{b, c, d\}], \mathbf{A}(g) = [\{a, b\}, \{a, b\}].$$

Then clearly, $IVSint(\mathbf{A}) = \mathbf{A}_{10}$. Thus $IVScl(IVSint(\mathbf{A})) = \tilde{X}_E$. So we have

$$IVSint(IVScl(IVSint(\mathbf{A}))) = \tilde{X}_E, \text{ i.e., } \mathbf{A} \subset IVSint(IVScl(IVSint(\mathbf{A}))).$$

Hence $\mathbf{A} \in IVS\alpha OS(X)$ but $\mathbf{A} \notin \tau$. Therefore $IVS\alpha OS(X) \not\subset \tau$.

Definition 5.11 ([36]). Let (X, τ, E) be a soft topological space and $A \in SS_E(X)$. Then

(i) the *soft α -closure* of A , denoted by $cl_{S\alpha}(A)$ or $cl_{S\alpha, \tau}(A)$, is a soft set over X defined as follows:

$$cl_{S\alpha}(A) = \bigcap \{F \in S\alpha CS(X) : A \subset F\},$$

(ii) the *soft α -interior* of A , denoted by $int_{S\alpha}(A)$ or $int_{S\alpha, \tau}(A)$, is a soft set over X defined as follows:

$$cl_{S\alpha}(A) = \bigcup \{U \in S\alpha OS(X) : U \subset A\}.$$

It is clear that $cl_{S\alpha}(A)$ is the smallest soft α -closed set over X which contains A and $int_{S\alpha}(A)$ is the largest soft α -open set over X which is contained in A .

Definition 5.12. Let (X, τ, E) be an IVSTS and $\mathbf{A} \in IVSS_E(X)$. Then

(i) the *interval-valued soft α -closure* of $\mathbf{A}$, denoted by $cl_{IVS\alpha}(\mathbf{A})$, is an IVSS over X defined as follows:

$$cl_{IVS\alpha}(\mathbf{A}) = \bigcap \{\mathbf{F} \in IVS\alpha CS(X) : \mathbf{A} \subset \mathbf{F}\},$$

(ii) the *interval-valued soft α -interior* of $\mathbf{A}$, denoted by $int_{IVS\alpha}(\mathbf{A})$, is an IVSS over X defined as follows:

$$int_{IVS\alpha}(\mathbf{A}) = \bigcup \{\mathbf{U} \in IVS\alpha OS(X) : \mathbf{U} \subset \mathbf{A}\}.$$

It is obvious that $cl_{IVS\alpha}(\mathbf{A})$ is the smallest interval-valued soft α -closed set over X containing $\mathbf{A}$ and $int_{IVS\alpha}(\mathbf{A})$ is the largest interval-valued soft α -open set over X contained in $\mathbf{A}$.

Remark 5.13. Let (X, τ, E) be an IVSTS and $\mathbf{A} \in IVSS_E(X)$. Then

- (1) $cl_{IVS\alpha}(\mathbf{A}) = [cl_{S\alpha, \tau-}(A^-), cl_{S\alpha, \tau+}(A^+)]$,
- (2) $int_{IVS\alpha}(\mathbf{A}) = [int_{S\alpha, \tau-}(A^-), int_{S\alpha, \tau+}(A^+)]$.

Example 5.14. Let (X, τ, E) be the IVSTS given in Example 5.4. To find IVS α OSs in X , consider the following IVSSs containing $\mathbf{U}$:

$$\mathbf{U}, \mathbf{A}_1, \mathbf{A}_2, \dots, \mathbf{A}_{20},$$

where $\mathbf{A}_1(e) = [\{a\}, \{a, b\}]$, $\mathbf{A}_1(f) = [\{b\}, X]$, $\mathbf{A}_2(e) = [\{a\}, \{a, b\}]$,
 $\mathbf{A}_2(f) = [\{b, c\}, \{b, c\}]$, $\mathbf{A}_3(e) = [\{a\}, \{a, b\}]$, $\mathbf{A}_3(f) = [\{b, c\}, X]$,
 $\mathbf{A}_4(e) = [\{a\}, \{a, b\}]$, $\mathbf{A}_4(f) = [\{b, c\}, X]$, $\mathbf{A}_5(e) = [\{a\}, \{a, b\}]$,
 $\mathbf{A}_5(f) = [X, X]$, $\mathbf{A}_6(e) = [\{a, b\}, \{a, b\}]$, $\mathbf{A}_6(f) = [\{b\}, \{b, c\}]$,
 $\mathbf{A}_6(e) = [\{a, b\}, \{a, b\}]$, $\mathbf{A}_6(f) = [\{b\}, \{b, c\}]$, $\mathbf{A}_7(e) = [\{a, b\}, \{a, b\}]$,
 $\mathbf{A}_6(f) = [\{b\}, X]$, $\mathbf{A}_8(e) = [\{a, b\}, \{a, b\}]$, $\mathbf{A}_8(f) = [\{b, c\}, \{b, c\}]$,
 $\mathbf{A}_9(e) = [\{a, b\}, \{a, b\}]$, $\mathbf{A}_9(f) = [\{b, c\}, X]$, $\mathbf{A}_{10}(e) = [\{a, b\}, \{a, b\}]$,
 $\mathbf{A}_{10}(f) = [X, X]$, $\mathbf{A}_{11}(e) = [\{a, b\}, X]$, $\mathbf{A}_{11}(f) = [\{b\}, \{b, c\}]$,
 $\mathbf{A}_{12}(e) = [\{a, b\}, X]$, $\mathbf{A}_{12}(f) = [\{b\}, X]$, $\mathbf{A}_{13}(e) = [\{a, b\}, X]$,
 $\mathbf{A}_{13}(f) = [\{b, c\}, \{b, c\}]$, $\mathbf{A}_{14}(e) = [\{a, b\}, X]$, $\mathbf{A}_{14}(f) = [\{b, c\}, X]$,
 $\mathbf{A}_{15}(e) = [\{a, b\}, X]$, $\mathbf{A}_{15}(f) = [X, X]$, $\mathbf{A}_{16}(e) = [X, X]$,
 $\mathbf{A}_{16}(f) = [\{b\}, \{b, c\}]$, $\mathbf{A}_{17}(e) = [X, X]$, $\mathbf{A}_{17}(f) = [\{b\}, X]$,
 $\mathbf{A}_{18}(e) = [X, X]$, $\mathbf{A}_{18}(f) = [\{b, c\}, \{b, c\}]$, $\mathbf{A}_{19}(e) = [X, X]$,

$$\mathbf{A}_{19}(f) = [\{b, c\}, X], \mathbf{A}_{20}(e) = [X, X], \mathbf{A}_{20}(f) = [X, X].$$

Then clearly, $IVSint(\mathbf{A}_i) = \mathbf{U}$ for each $i \in \{1, 2, \dots, 19\}$. Since $\tilde{X}_E$ is the only IVSCS in X containing $\mathbf{U}$, $IVScl(IVSint(\mathbf{A}_i)) = \tilde{X}_E$ for each $i \in \{1, 2, \dots, 20\}$. Thus by Proposition 5.26 in [34], $IVSint(IVScl(IVSint(\mathbf{A}_i))) = \tilde{X}_E$ for each $i \in \{1, 2, \dots, 20\}$. Since $\mathbf{A}_i \subset \tilde{X}_E$, $\mathbf{A}_i \in IVS\alpha OS(X)$ for each $i \in \{1, 2, \dots, 20\}$. So $cl_{IVS\alpha}(\mathbf{A}_i) = \tilde{X}_E$ for each $i \in \{1, 2, \dots, 20\}$.

Theorem 5.15. *Let (X, τ, E) be an IVSTS and $\mathbf{A} \in IVSS_E(X)$. Then*

- (1) $\mathbf{A} \in IVS\alpha CS(X)$ if and only if $A = cl_{IVS\alpha}(\mathbf{A})$,
- (2) $\mathbf{A} \in IVS\alpha OS(X)$ if and only if $A = int_{IVS\alpha}(\mathbf{A})$.

Proof. The proofs are straightforward. $\square$

Proposition 5.16. *Let (X, τ, E) be an IVSTS and $\mathbf{A} \in IVSS_E(X)$. Then*

- (1) $[cl_{IVS\alpha}(\mathbf{A})]^c = int_{IVS\alpha}(\mathbf{A}^c)$,
- (2) $[int_{IVS\alpha}(\mathbf{A})]^c = cl_{IVS\alpha}(\mathbf{A}^c)$.

Proof. The proofs follow from Definition 5.12. $\square$

Proposition 5.17. *Let (X, τ, E) be an IVSTS and $\mathbf{A}, \mathbf{B} \in IVSS_E(X)$. Then*

- (1) $cl_{IVS\alpha}(\tilde{\emptyset}_E) = \tilde{\emptyset}_E$, $cl_{IVS\alpha}(\tilde{X}_E) = \tilde{X}_E$,
- (2) $cl_{IVS\alpha}(\mathbf{A}) \in IVS\alpha CS(X)$,
- (3) if $\mathbf{A} \subset \mathbf{B}$, then $cl_{IVS\alpha}(\mathbf{A}) \subset cl_{IVS\alpha}(\mathbf{B})$,
- (4) $cl_{IVS\alpha}(cl_{IVS\alpha}(\mathbf{A})) = cl_{IVS\alpha}(\mathbf{A})$,
- (5) $cl_{IVS\alpha}(\mathbf{A} \cup \mathbf{B}) = cl_{IVS\alpha}(\mathbf{A}) \cup cl_{IVS\alpha}(\mathbf{B})$,
- (6) $cl_{IVS\alpha}(\mathbf{A} \cap \mathbf{B}) \subset cl_{IVS\alpha}(\mathbf{A}) \cap cl_{IVS\alpha}(\mathbf{B})$.

Proof. The proofs are straightforward. $\square$

Proposition 5.18. *Let (X, τ, E) be an IVSTS and $\mathbf{A}, \mathbf{B} \in IVSS_E(X)$. Then*

- (1) $int_{IVS\alpha}(\tilde{\emptyset}_E) = \tilde{\emptyset}_E$, $int_{IVS\alpha}(\tilde{X}_E) = \tilde{X}_E$,
- (2) $int_{IVS\alpha}(\mathbf{A}) \in IVS\alpha OS(X)$,
- (3) if $\mathbf{A} \subset \mathbf{B}$, then $int_{IVS\alpha}(\mathbf{A}) \subset int_{IVS\alpha}(\mathbf{B})$,
- (4) $int_{IVS\alpha}(int_{IVS\alpha}(\mathbf{A})) = int_{IVS\alpha}(\mathbf{A})$,
- (5) $int_{IVS\alpha}(\mathbf{A} \cup \mathbf{B}) \subset int_{IVS\alpha}(\mathbf{A}) \cup int_{IVS\alpha}(\mathbf{B})$,
- (6) $int_{IVS\alpha}(\mathbf{A} \cap \mathbf{B}) = int_{IVS\alpha}(\mathbf{A}) \cap int_{IVS\alpha}(\mathbf{B})$.

Proof. The proofs are straightforward. $\square$

6. INTERVAL-VALUED SOFT α -SEPARATION AXIOMS

In this section, we propose some of new separation axioms such as the $IVS\alpha T_0(j)$, $IVS\alpha T_1(j)$, $IVS\alpha T_2(j)$, $IVS\alpha T_3(j)$ and $IVS\alpha T_4(j)$ axioms for $j=i, ii$ as a generalization of separation axioms discussed in Section 3. Furthermore, we study some of their properties and the relations between them in the general framework of IVSTSs.

Definition 6.1 ([35]). A soft topological space (X, τ, E) is called a *soft αT_0 -space*, if for any $x \neq y \in X$, there are $U, V \in S\alpha OS(X)$ such that either $x \in U, y \notin U$ or $y \in V, x \notin V$.

Definition 6.2. An IVSTS (X, τ, E) is called an:

- (i) *interval-valued soft αT_0 (i)-space* (briefly, $\text{IVS}\alpha T_0$ (i)-space), if for any $x \neq y \in X$, there are $\mathbf{U}, \mathbf{V} \in \text{IVS}\alpha\text{OS}(X)$ such that either $x_1 \in \mathbf{U}, y_1 \notin \mathbf{U}$ or $y_1 \in \mathbf{V}, x_1 \notin \mathbf{V}$,
- (ii) *interval-valued soft αT_0 (ii)-space* (briefly, $\text{IVS}\alpha T_0$ (ii)-space), if for any $x \neq y \in X$, there are $\mathbf{U}, \mathbf{V} \in \text{IVS}\alpha\text{OS}(X)$ such that either $x_0 \in \mathbf{U}, y_0 \notin \mathbf{U}$ or $y_0 \in \mathbf{V}, x_0 \notin \mathbf{V}$.

Remark 6.3. (1) Every IVST_0 (i) [resp. IVST_0 (ii)]-space is an $\text{IVS}\alpha T_0$ (i) [resp. $\text{IVS}\alpha T_0$ (ii)]-space.

(2) Every $\text{IVS}\alpha T_0$ (i)-space is an $\text{IVS}\alpha T_0$ (ii)-space. The converse is not true in general (See Example 6.4).

(3) If an IVSTS (X, τ, E) is an $\text{IVS}\alpha T_0$ (i) [resp. $\text{IVS}\alpha T_0$ (ii)]-space, then (X, τ^-, E) and (X, τ^+, E) are soft αT_0 -spaces [resp. (X, τ^+, E) is a soft αT_0 -space].

Example 6.4. Let $X = \{a, b\}$, $E = \{e, f\}$ and consider the IVST τ_1 given by:

$$\tau_1 = \{\tilde{\varnothing}_E, \mathbf{A}, \tilde{X}_E\},$$

where $\mathbf{A}(e) = \mathbf{A}(f) = [\{a\}, \{a\}]$. Then we can easily check that (X, τ_1, E) is an $\text{IVS}\alpha T_0$ (i)-space. Moreover, we can confirm that $\tau_1^- = \tau_1^+$ and (X, τ_1^-, E) is a soft αT_0 -space.

Now consider the IVST τ_2 given by:

$$\tau_2 = \{\tilde{\varnothing}_E, \mathbf{A}, \tilde{X}_E\},$$

where $\mathbf{A}(e) = \mathbf{A}(f) = \{\varnothing, \{a\}\}$. Then clearly, (X, τ_2, E) is an $\text{IVS}\alpha T_0$ (ii)-space and (X, τ_2^+, E) is a soft αT_0 -space. But (X, τ_2, E) is not an $\text{IVS}\alpha T_0$ (i)-space.

The following is a similar consequence of Proposition 3.13.

Proposition 6.5. Let (X, τ, E) be an IVSTS and let $x \neq y \in X$.

- (1) If there are $\mathbf{U}, \mathbf{U} \in \text{IVS}\alpha\text{OS}(X)$ such that either $x_1 \in \mathbf{U}, y_1 \in \mathbf{U}^c$ or $y_1 \in \mathbf{V}, x_1 \in \mathbf{V}^c$, then X is an $\text{IVS}\alpha T_0$ (i)-space.
- (2) If there are $\mathbf{U}, \mathbf{U} \in \text{IVS}\alpha\text{OS}(X)$ such that either $x_0 \in \mathbf{U}, y_0 \in \mathbf{U}^c$ or $y_0 \in \mathbf{V}, x_0 \in \mathbf{V}^c$, then X is an $\text{IVS}\alpha T_0$ (ii)-space.

Proof. The proofs are similar to Proposition 3.13. □

Also, we obtain a similar consequence of Proposition 4.10 (1).

Proposition 6.6. Let (X, τ, E) be an IVSTS and let Y be a nonempty subset of X . If X is an $\text{IVS}\alpha T_0$ (i) [resp. $\text{IVS}\alpha T_0$ (ii)]-space, then (Y, τ_Y, E) is an $\text{IVS}\alpha T_0$ (i) [resp. $\text{IVS}\alpha T_0$ (ii)]-space.

Proof. The proofs are similar to Proposition 4.10 (1). □

Definition 6.7 ([35]). A soft topological space (X, τ, E) is called a *soft αT_1 -space*, if for any $x \neq y \in X$, there are $U, V \in \text{S}\alpha\text{OS}(X)$ such that $x \in U, y \notin U$ and $y \in V, x \notin V$.

Definition 6.8. An IVSTS (X, τ, E) is called an:

- (i) *interval-valued soft αT_1 (i)-space* (briefly, $\text{IVS}\alpha T_1$ (i)-space), if for any $x \neq y \in X$, there are $\mathbf{U}, \mathbf{V} \in \text{IVS}\alpha\text{OS}(X)$ such that $x_1 \in \mathbf{U}, y_1 \notin \mathbf{U}$ and $y_1 \in \mathbf{V}, x_1 \notin \mathbf{V}$,

(ii) *interval-valued soft αT_1 (ii)-space* (briefly, $\text{IVS}\alpha T_1$ (ii)-space), if for any $x \neq y \in X$, there are $\mathbf{U}, \mathbf{V} \in \text{IVS}\alpha\text{OS}(X)$ such that $x_0 \in \mathbf{U}$, $y_0 \notin \mathbf{U}$ and $y_0 \in \mathbf{V}$, $x_0 \notin \mathbf{V}$.

Remark 6.9. (1) Every $\text{IVS}\alpha T_1$ (i)-space is an $\text{IVS}\alpha T_1$ (ii)-space.

(2) If an $\text{IVSTS} (X, \tau, E)$ is an $\text{IVS}\alpha T_1$ (i) [resp. $\text{IVS}\alpha T_1$ (ii)]-space, then (X, τ^-, E) and (X, τ^+, E) are soft αT_1 -spaces [resp. (X, τ^+, E) is a soft αT_1 -space].

Example 6.10. Let $X = \{a, b\}$, $E = \{e, f\}$ and consider the $\text{IVST} \tau_1$ given by:

$$\tau_1 = \{\tilde{\varnothing}_E, \mathbf{A}_1, \mathbf{A}_2, \mathbf{A}_3, \tilde{X}_E\},$$

where $\mathbf{A}_1(e) = [X, X]$, $\mathbf{A}_1(f) = [\{b\}, \{b\}]$,
 $\mathbf{A}_2(e) = [\{a\}, \{a\}]$, $\mathbf{A}_2(f) = [X, X]$,
 $\mathbf{A}_3(e) = [\{a\}, \{a\}]$, $\mathbf{A}_3(f) = [\{b\}, \{b\}]$.

Then we can easily see that (X, τ_1, E) is an $\text{IVS}\alpha T_1$ (i)-space. Moreover, (X, τ_1^-, E) and (X, τ_1^+, E) are soft αT_1 -spaces.

Now consider the $\text{IVST} \tau_2$ given by:

$$\tau_2 = \{\tilde{\varnothing}_E, \mathbf{A}_1, \mathbf{A}_2, \mathbf{A}_3, \tilde{X}_E\},$$

where $\mathbf{A}_1(e) = [\varnothing, X]$, $\mathbf{A}_1(f) = [\varnothing, \{b\}]$,
 $\mathbf{A}_2(e) = [\varnothing, \{a\}]$, $\mathbf{A}_2(f) = [\varnothing, X]$,
 $\mathbf{A}_3(e) = [\varnothing, \{a\}]$, $\mathbf{A}_3(f) = [\varnothing, \{b\}]$.

Then clearly (X, τ_2, E) is an $\text{IVS}\alpha T_1$ (ii)-space but not an $\text{IVS}\alpha T_1$ (i)-space. Furthermore, (X, τ_2^+, E) is a soft αT_1 -spaces.

The following is an immediate consequence of Definitions 6.5 and 6.8.

Proposition 6.11. *Every IVST_1 (i) [resp. IVST_1 (ii)]-space is an $\text{IVS}\alpha T_1$ (i) [resp. $\text{IVS}\alpha T_1$ (ii)]-space. But the converse is not true in general (See Example 6.12).*

Proof. The proof is obvious. $\square$

Example 6.12. Let $X = \{a, b, c, d\}$, $E = \{e, f\}$ and consider the $\text{IVST} \tau_1$ given by:

$$\tau_1 = \{\tilde{\varnothing}_E, \mathbf{A}_1, \mathbf{A}_2, \mathbf{A}_3, \tilde{X}_E\},$$

where $\mathbf{A}_1(e) = \mathbf{A}_1(f) = [\{a, b, c\}, \{a, b, c\}]$,
 $\mathbf{A}_2(e) = \mathbf{A}_2(f) = [\{b, c\}, \{b, c\}]$,
 $\mathbf{A}_3(e) = \mathbf{A}_3(f) = [\{b\}, \{b\}]$.

Then we can see that (X, τ_1, E) is an $\text{IVS}\alpha T_0$ (i)-space but not an $\text{IVS}\alpha T_1$ (i)-space.

Now consider $\text{IVST} \tau_2$ given by:

$$\tau_2 = \{\tilde{\varnothing}_E, \mathbf{A}_1, \mathbf{A}_2, \mathbf{A}_3, \tilde{X}_E\},$$

where $\mathbf{A}_1(e)\mathbf{A}_1(f) = [\varnothing, \{a, b, c\}]$,
 $\mathbf{A}_2(e) = \mathbf{A}_2(f) = [\varnothing, \{b, c\}]$,
 $\mathbf{A}_3(e) = \mathbf{A}_3(f) = [\varnothing, \{b\}]$.

Then we can easily check that (X, τ_2, E) is an $\text{IVS}\alpha T_0$ (i)-space but not an $\text{IVS}\alpha T_1$ (ii)-space.

We have a similar consequence of Proposition 3.10.

Proposition 6.13. *Let (X, τ, E) be an IVSTS .*

(1) *If $\mathbf{x}_1$ is an $\text{IVS}\alpha\text{CS}$ in X for each $x \in X$, then X is an $\text{IVS}\alpha T_1$ (i)-space.*

(2) *If $\mathbf{x}_0$ is an $\text{IVS}\alpha\text{CS}$ in X for each $x \in X$, then X is an $\text{IVS}\alpha T_1$ (ii)-space.*

Proof. The proofs are similar to Proposition 3.10. $\square$

Theorem 6.14. Let (X, τ, E) be an IVSTS.

- (1) X is an $IVS\alpha T_1(i)$ -space if and only if $x_1 \in IVS\alpha CS(X)$ for each $x \in X$.
- (2) X is an $IVS\alpha T_1(ii)$ -space if and only if $x_0 \in IVS\alpha CS(X)$ for each $x \in X$.

Proof. (1) Suppose X is an $IVS\alpha T_1(i)$ -space and let $x \neq y \in X$. Then there is $\mathbf{U}_{x_1} \in IVS\alpha OS(X)$ such that $x_1 \in \mathbf{U}_{x_1}$, $y_1 \notin \mathbf{U}_{x_1}$. Thus by Proposition 5.8 (1), $x_1 \in \bigcup_{x \in X} \mathbf{U}_{x_1} \in IVS\alpha OS(X)$. So $x_1 = [\bigcup_{x \in X} \mathbf{U}_{x_1}]^c \in IVS\alpha CS(X)$.

Conversely, suppose the necessary condition holds and let $x \neq y \in X$. Then clearly, $y_1 \in [X - \{x\}, X - \{x\}] \in IVS\alpha OS(X)$ and $x_1 \notin [X - \{x\}, X - \{x\}]$. Similarly, $x_1 \in [X - \{y\}, X - \{y\}] \in IVS\alpha OS(X)$ and $y_1 \notin [X - \{y\}, X - \{y\}]$. Thus X is an $IVS\alpha T_1(i)$ -space.

- (2) The proof is similar to (1). $\square$

The following is a similar consequence of Proposition 4.10 (2).

Proposition 6.15. Let (X, τ, E) be an IVSTS and let Y be a nonempty subset of X . If X is an $IVS\alpha T_1(i)$ [resp. $IVS\alpha T_1(ii)$]-space, then (Y, τ_Y, E) is an $IVS\alpha T_1(i)$ [resp. $IVS\alpha T_1(ii)$]-space

Proof. The proof is similar to Proposition 4.10 (2). $\square$

Definition 6.16 ([35]). A soft topological space (X, τ, E) is called a *soft αT_2 -space*, if for any $x \neq y \in X$, there are $U, V \in S\alpha OS(X)$ such that $x \in U$, $y \in V$ and $U \cap V = \emptyset_E$.

Definition 6.17. An IVSTS (X, τ, E) is called an:

- (i) *interval-valued soft $\alpha T_2(i)$ -space* (briefly, $IVS\alpha T_2(i)$ -space), if for any $x \neq y \in X$, there are $\mathbf{U}, \mathbf{V} \in IVS\alpha OS(X)$ such that $x_1 \in \mathbf{U}$, $y_1 \in \mathbf{V}$ and $\mathbf{U} \cap \mathbf{V} = \tilde{\emptyset}_E$,
- (ii) *interval-valued soft $\alpha T_2(ii)$ -space* (briefly, $IVS\alpha T_2(ii)$ -space), if for any $x \neq y \in X$, there are $\mathbf{U}, \mathbf{V} \in IVS\alpha OS(X)$ such that $x_0 \in \mathbf{U}$, $y_0 \in \mathbf{V}$ and $\mathbf{U} \cap \mathbf{V} = \tilde{\emptyset}_E$.

Remark 6.18. (1) Every $IVS\alpha T_2(i)$ -space is an $IVS\alpha T_2(ii)$ -space.

(2) If an IVSTS (X, τ, E) is an $IVS\alpha T_2(i)$ [resp. $IVS\alpha T_2(ii)$]-space, then (X, τ^-, E) and (X, τ^+, E) are soft αT_2 -spaces [resp. (X, τ^+, E) is a soft αT_2 -space].

Example 6.19. Let $X = \{a, b\}$, $E = \{e, f\}$ and consider the IVST τ_1 given by:

$$\tau_1 = \{\tilde{\emptyset}_E, \mathbf{A}_1, \mathbf{A}_2, \tilde{X}_E\},$$

where $\mathbf{A}_1(e) = \mathbf{A}_1(f) = [\{a\}, \{a\}]$, $\mathbf{A}_2(e) = \mathbf{A}_2(f) = [\{b\}, \{b\}]$.

Then clearly, (X, τ_1, E) is an $IVS\alpha T_2(i)$ -space, and, (X, τ_1^-, E) and (X, τ_1^+, E) are soft αT_2 -spaces.

Now consider the IVST τ_2 given by:

$$\tau_2 = \{\tilde{\emptyset}_E, \mathbf{A}_1, \mathbf{A}_2, \tilde{X}_E\},$$

where $\mathbf{A}_1(e) = \mathbf{A}_1(f) = [\emptyset, \{a\}]$, $\mathbf{A}_2(e) = \mathbf{A}_2(f) = [\emptyset, \{b\}]$.

Then it is conspicuous that (X, τ_2, E) is an $IVS\alpha T_2(ii)$ -space but not an $IVS\alpha T_2(i)$ -space. Moreover, (X, τ_2^+, E) is a soft αT_2 -spaces.

Proposition 6.20. Every $IVS\alpha T_2(i)$ [resp. $IVS\alpha T_2(ii)$]-space is an $IVS\alpha T_1(i)$ [resp. $IVS\alpha T_1(ii)$]-space. But the converse is not true in general (See Example 6.21).

Proof. The proof is clear. $\square$

Example 6.21. Consider the $\text{IVS}\alpha T_1(\text{i})$ -space (X, τ_1, E) [resp. $\text{IVS}\alpha T_1(\text{ii})$ -space (X, τ_2, E)] given in Example 6.10. Then we can see that (X, τ_1, E) is not an $\text{IVS}\alpha T_2(\text{i})$ [resp. $\text{IVS}\alpha T_2(\text{ii})$]-space.

We have a similar consequence of Proposition 4.11.

Proposition 6.22. Let (X, τ, E) be an IVSTS and let Y be a nonempty subset of X . If X is an $\text{IVS}\alpha T_2(\text{i})$ [resp. $\text{IVS}\alpha T_2(\text{ii})$]-space, then (X, τ_Y, E) is an $\text{IVS}\alpha T_2(\text{i})$ [resp. $\text{IVS}\alpha T_2(\text{ii})$]-space.

Proof. The proof is similar to Proposition 4.11. $\square$

Definition 6.23 ([35]). A soft topological space (X, τ, E) is called a *soft α -regular space*, if for each $A \in \text{S}\alpha\text{CS}(X)$ and each $x \in X$ with $x \notin A$, there are $U, V \in \text{S}\alpha\text{OS}(X)$ such that $x \in U$, $A \subset V$ and $U \cap V = \emptyset_E$.

Definition 6.24. An IVSTS (X, τ, E) is called an:

- (i) *interval-valued soft α -regular(i)-space* (briefly, $\text{IVS}\alpha\text{R}(\text{i})$ -space), if for each $\mathbf{A} \in \text{IVS}\alpha\text{CS}(X)$ and each $x \in X$ with $x_1 \notin \mathbf{A}$, there are $\mathbf{U}, \mathbf{V} \in \text{IVS}\alpha\text{OS}(X)$ such that $x_1 \in \mathbf{U}$, $\mathbf{A} \subset \mathbf{V}$ and $\mathbf{U} \cap \mathbf{V} = \tilde{\emptyset}_E$,
- (ii) *interval-valued soft α -regular(ii)-space* (briefly, $\text{IVS}\alpha\text{R}(\text{ii})$ -space), if for each $\mathbf{A} \in \text{IVS}\alpha\text{CS}(X)$ and each $x \in X$ with $x_0 \notin \mathbf{A}$, there are $\mathbf{U}, \mathbf{V} \in \text{IVS}\alpha\text{OS}(X)$ such that $x_0 \in \mathbf{U}$, $\mathbf{A} \subset \mathbf{V}$ and $\mathbf{U} \cap \mathbf{V} = \tilde{\emptyset}_E$.

It is obvious that if an IVSTS (X, τ, E) is an $\text{IVSR}(\text{i})$ [resp. $\text{IVSR}(\text{ii})$]-space, then it is an $\text{IVS}\alpha\text{R}(\text{i})$ [resp. $\text{IVS}\alpha\text{R}(\text{ii})$]-space.

Example 6.25. Let $X = \{a, b, c\}$, $E = \{e, f\}$ and consider the IVST τ_1 on X defined by:

$$\tau_1 = \{\tilde{\emptyset}_E, \mathbf{A}_1, \mathbf{A}_2, \tilde{X}_E\},$$

where $\mathbf{A}_1(e) = \mathbf{A}_1(f) = [\{a\}, \{a\}]$, $\mathbf{A}_2(e) = \mathbf{A}_2(f) = [\{b, c\}, \{b, c\}]$.

Then we can easily check that (X, τ_1, E) is an $\text{IVS}\alpha\text{R}(\text{i})$ -space but not an $\text{IVS}\alpha T_1(\text{i})$ -space. Now consider the IVST τ_2 on X defined by:

$$\tau_2 = \{\tilde{\emptyset}_E, \mathbf{A}_1, \mathbf{A}_2, \tilde{X}_E\},$$

where $\mathbf{A}_1(e) = \mathbf{A}_1(f) = [\emptyset, \{a\}]$, $\mathbf{A}_2(e) = \mathbf{A}_2(f) = [\emptyset, \{b, c\}]$.

Then clearly, (X, τ_1, E) is an $\text{IVS}\alpha\text{R}(\text{ii})$ -space but not an $\text{IVS}\alpha T_1(\text{ii})$ -space.

Definition 6.26 ([35]). A soft topological space (X, τ, E) is called a *soft αT_3 -space*, if it is a soft αT_1 -space and a soft α -regular space.

Definition 6.27. An IVSTS (X, τ, E) is called an:

- (i) *interval-valued soft $\alpha T_3(\text{i})$ -space* (briefly, $\text{IVS}\alpha T_3(\text{i})$ -space), if it is an $\text{IVS}\alpha T_1(\text{i})$ -space and an $\text{IVS}\alpha\text{R}(\text{i})$ -space,
- (ii) *interval-valued soft $\alpha T_3(\text{ii})$ -space* (briefly, $\text{IVS}\alpha T_3(\text{ii})$ -space), if it is an $\text{IVS}\alpha T_1(\text{ii})$ -space and an $\text{IVS}\alpha\text{R}(\text{ii})$ -space.

Remark 6.28. Every $\text{IVS}\alpha T_3(\text{i})$ -space is an $\text{IVS}\alpha T_3(\text{ii})$ -space.

Proposition 6.29. Every $\text{IVS}\alpha T_3(\text{i})$ [resp. $\text{IVS}\alpha T_3(\text{ii})$]-space is an $\text{IVS}\alpha T_2(\text{i})$ [resp. $\text{IVS}\alpha T_2(\text{ii})$]-space. But the converse is not true in general (See Example 6.30).

Proof. Let (X, τ, E) be an $\text{IVS}\alpha T_3(\text{i})$ -space and let $x \neq y \in X$. Since X is an $\text{IVS}\alpha T_1(\text{i})$ -space, by Theorem 6.14, $y_1 \in \text{IVS}\alpha CS(X)$ and $x_1 \notin y_1$. Since X is an $\text{IVS}\alpha R(\text{i})$ -space, there are $\mathbf{U}, \mathbf{V} \in \text{IVS}\alpha OS(X)$ such that $x_1 \in \mathbf{U}$, $y_1 \subset \mathbf{V}$ and $\mathbf{U} \cap \mathbf{V} = \tilde{\varnothing}_E$. Then (X, τ, E) is an $\text{IVS}\alpha T_2(\text{i})$ -space. The proof of the second part can be done by the same token. $\square$

We obtain a similar consequence of Proposition 4.12.

Proposition 6.30. *Let (X, τ, E) be an IVSTS and let Y be a nonempty subset of X . If X is an $\text{IVS}\alpha T_3(\text{i})$ [resp. $\text{IVS}\alpha T_3(\text{ii})$]-space, then (X, τ_Y, E) is an $\text{IVS}\alpha T_3(\text{i})$ [resp. $\text{IVS}\alpha T_3(\text{ii})$]-space.*

Proof. The proof is similar to Proposition 4.12. $\square$

Definition 6.31 ([35]). A soft topological space (X, τ, E) is called a *soft α -normal space*, if for each $A, B \in \text{S}\alpha CS(X)$ with $A \cap B = \varnothing_E$, there are $U, V \in \text{S}\alpha OS(X)$ such that $A \subset U$, $B \subset V$ and $U \cap V = \varnothing_E$.

Definition 6.32. An IVSTS (X, τ, E) is called an *interval-valued soft α -normal space* (briefly, $\text{IVS}\alpha N$ -space), if for each $\mathbf{A}, \mathbf{B} \in \text{IVS}\alpha CS(X)$ with $\mathbf{A} \cap \mathbf{B} = \tilde{\varnothing}_E$, there are $\mathbf{U}, \mathbf{V} \in \text{IVS}\alpha OS(X)$ such that $x_1 \in \mathbf{U}$, $\mathbf{A} \subset \mathbf{V}$ and $\mathbf{U} \cap \mathbf{V} = \tilde{\varnothing}_E$.

Remark 6.33. Every $\text{IVS}\alpha N$ -space may be neither an $\text{IVS}\alpha R(\text{i})$ [resp. $\text{IVS}\alpha R(\text{ii})$]-space nor an $\text{IVS}\alpha T_1(\text{i})$ [resp. $\text{IVS}\alpha T_1(\text{ii})$]-space (See Example 6.34).

Example 6.34. Let $X = \{a, b, c\}$, $E = \{e, f\}$ and consider the IVST τ on X given by:

$$\tau = \{\tilde{\varnothing}_E, \mathbf{A}_1, \mathbf{A}_2, \mathbf{A}_3, \tilde{X}_E\},$$

where $\mathbf{A}_1(e) = \mathbf{A}_1(f) = [\{a\}, \{a\}]$, $\mathbf{A}_2(e) = \mathbf{A}_2(f) = [\{b\}, \{b\}]$,
 $\mathbf{A}_3(e) = \mathbf{A}_3(f) = [\{a, b\}, \{a, b\}]$.

Then we can see that (X, τ, E) is an $\text{IVS}\alpha N$ -space but neither an $\text{IVS}\alpha R(\text{i})$ [resp. $\text{IVS}\alpha R(\text{ii})$]-space nor an $\text{IVS}\alpha T_1(\text{i})$ [resp. $\text{IVS}\alpha T_1(\text{ii})$]-space.

Definition 6.35 ([35]). A soft topological space (X, τ, E) is called a *soft αT_4 -space*, if it is a soft αT_1 -space and a soft α -normal space.

Definition 6.36. An IVSTS (X, τ, E) is called an:

- (i) *interval-valued soft $\alpha T_4(\text{i})$ -space* (briefly, $\text{IVS}\alpha T_4(\text{i})$ -space), if It is an $\text{IVS}\alpha T_1(\text{i})$ -space and an $\text{IVS}\alpha N$ -space,
- (ii) *interval-valued soft $\alpha T_4(\text{ii})$ -space* (briefly, $\text{IVS}\alpha T_4(\text{ii})$ -space), if It is an $\text{IVS}\alpha T_1(\text{ii})$ -space and an $\text{IVS}\alpha N$ -space.

Example 6.37. In Example 6.19, we can easily check that an IVSTS (X, τ_1, E) [resp. (X, τ_2, E)] is an $\text{IVS}\alpha T_4(\text{i})$ [resp. $\text{IVS}\alpha T_4(\text{ii})$]-space.

Proposition 6.38. *Every $\text{IVS}\alpha T_4(\text{i})$ [resp. $\text{IVS}\alpha T_4(\text{ii})$]-space is an $\text{IVS}\alpha T_3(\text{i})$ [resp. $\text{IVS}\alpha T_3(\text{ii})$]-space.*

Proof. Let X be an $\text{IVS}\alpha T_4(\text{i})$ -space. Since X is an $\text{IVS}\alpha T_1(\text{i})$ -space, it is enough to prove that X is an $\text{IVS}\alpha R(\text{i})$ -space. Let $\mathbf{A} \in \text{IVS}\alpha CS(X)$ with $x_1 \notin \mathbf{A}$. Since X is an $\text{IVS}\alpha T_1(\text{i})$ -space, by Theorem 6.14 (1), $x_1 \in \text{IVS}\alpha CS(X)$. Note that $\mathbf{A} \cap x_1 = \varnothing_E$. Since X is an $\text{IVS}\alpha N$ -space, there are $\mathbf{U}, \mathbf{V} \in \text{IVS}\alpha OS(X)$ such that $x_1 \in \mathbf{U}$,

$\mathbf{A} \subset \mathbf{V}$ and $\mathbf{U} \cap \mathbf{V} = \tilde{\mathcal{O}}_E$. Thus X is an $\text{IVS}\alpha T_3(\text{i})$ -space. The proof of the second part is analogous and therefore omitted. $\square$

Remark 6.39. From Propositions 6.11, 6.20, 6.29 and 6.38, we have relationships among $\text{IVS}\alpha T_0(\text{i})$ [resp. $\text{IVS}\alpha T_0(\text{ii})$], $\text{IVS}\alpha T_1(\text{i})$ [resp. $\text{IVS}\alpha T_1(\text{ii})$], $\text{IVS}\alpha T_2(\text{i})$ [resp. $\text{IVS}\alpha T_2(\text{ii})$], $\text{IVS}\alpha T_3(\text{i})$ [resp. $\text{IVS}\alpha T_3(\text{ii})$] and $\text{IVS}\alpha T_4(\text{i})$ [resp. $\text{IVS}\alpha T_4(\text{ii})$]:

$$\begin{aligned} \text{IVS}\alpha T_4(\text{i}) \text{ [resp. } \text{IVS}\alpha T_4(\text{ii})] &\implies \text{IVS}\alpha T_3(\text{i}) \text{ [resp. } \text{IVS}\alpha T_3(\text{ii})] \\ &\implies \text{IVS}\alpha T_2(\text{i}) \text{ [resp. } \text{IVS}\alpha T_2(\text{ii})] \\ &\implies \text{IVS}\alpha T_1(\text{i}) \text{ [resp. } \text{IVS}\alpha T_1(\text{ii})] \\ &\implies \text{IVS}\alpha T_0(\text{i}) \text{ [resp. } \text{IVS}\alpha T_0(\text{ii})]. \end{aligned}$$

7. CONCLUSIONS

First, we defined separation axioms, i.e., $\text{IVST}_0(\text{i})$ [resp. $\text{IVST}_0(\text{ii})$], $\text{IVST}_1(\text{i})$ [resp. $\text{IVST}_1(\text{ii})$], $\text{IVST}_2(\text{i})$ [resp. $\text{IVST}_2(\text{ii})$], $\text{IVST}_3(\text{i})$ [resp. $\text{IVST}_3(\text{ii})$] and $\text{IVST}_4(\text{i})$ [resp. $\text{IVST}_4(\text{ii})$], and studied some of their relationships. Second, by using interval-valued soft α -open sets, we introduced some of new separation axioms, i.e., $\text{IVS}\alpha T_0(\text{i})$ [resp. $\text{IVS}\alpha T_0(\text{ii})$], $\text{IVS}\alpha T_1(\text{i})$ [resp. $\text{IVS}\alpha T_1(\text{ii})$], $\text{IVS}\alpha T_2(\text{i})$ [resp. $\text{IVS}\alpha T_2(\text{ii})$], $\text{IVS}\alpha T_3(\text{i})$ [resp. $\text{IVS}\alpha T_3(\text{ii})$] and $\text{IVS}\alpha T_4(\text{i})$ [resp. $\text{IVS}\alpha T_4(\text{ii})$], and discussed relationships among them. Moreover, we dealt with hereditary properties of each separation axiom.

In the future, we would like to study new separation axioms as well as decision making problems that were not covered in our study.

Acknowledgements. We thank the four judges for their detailed and kind advice. This paper was supported by Wonkwang University in 2024.

REFERENCES

- [1] D. Molodtsov, Soft set theory—First results, *Comput. Math. Appl.* 37 (4-5) (1999) 19–31.
- [2] P. K. Maji, R. Biswas and A. R. Roy, Soft set theory, *Comput. Math. Appl.* 45 (4-5) (2003) 555–562.
- [3] M. I. Ali, F. Feng, X. Liu, W. K. Min and M. Shabir, On some new operations in soft set theory, *Comput. Math. Appl.* 57 (2009) 1547–1553.
- [4] K. V. Babitha and J. J. Sunil, Soft set relations and functions, *Comput. Math. Appl.* 60 (7) (2010) 1840–1849.
- [5] Athar Kharal and B. Ahmad, Mappings on soft classes, *New Math. Nat. Comput.* 7 (3) (2011) 471–481.
- [6] H. Aktaş and N. Çağman, Soft sets and soft groups, *Infor. Sci.* 1 (77) (2007) 2726–2735.
- [7] F. Feng, Y. B. Jun and X. Zhao, Soft semirings, *Fuzzy Sets and Systems: Theory and Applications* 56 (10) (2008) 2621–2628.
- [8] U. Acar, F. Koyuncu and B. Tanay, Soft sets and soft rings, *Comput. Math. Appl.* 59 (2010) 3458–3463.
- [9] Q. M. Sun, Z. L. Zhang and J. Liu, Soft sets and soft modules, *Proceedings of Rough Sets and Knowledge Technology, Third International Conference, RSKT 2008*, 17–19 May, Chengdu, China 403–409.
- [10] Y. B. Jun, Soft BCK/BCI -algebras, *Comput. Math. Appl.* 56 (2008) 1408–1413.
- [11] Y. B. Jun and C. H. Park, Applications of soft sets in ideal theory in BCK/BCI -algebras, *Inform. Sci.* 178 (2008) 2466–2475.
- [12] Y. B. Jun and C. H. Park, Applications of soft sets in Hilbert algebras, *Iranian Journal Fuzzy Systems* 6 (2) (2009) 55–86.

- [13] P. Majumdar and S. K. Samanta, Similarity measure of soft sets, *New Mathematics and Natural Computation* 4 (1) (2008) 1–12.
- [14] N. Çağman and S. Enginoglu, Soft set theory and uni-int decision making, *European Journal of Operational Research* 207 (2010) 848–855.
- [15] N. Çağman and S. Enginoglu, Soft matrix theory and its decision making, *Comput. Math. Appl.* 59 (2010) 3308–3314.
- [16] M. Shabir and M. Naz, On soft topological spaces, *Comput. Math. Appl.* 61 (2011) 1786–1799.
- [17] N. Çağman, S. Karataş and S. Enginoglu, Soft topology, *Comput. Math. Appl.* 62 (2011) 351–358.
- [18] W. K. Min, A note on soft topological spaces, *Comput. Math. Appl.* 62 (2011) 3524–3528.
- [19] S. Hussain and B. Ahmad, Some properties of soft topological spaces, *Comput. Math. Appl.* 62 (11) (2011) 4058–4067.
- [20] I. Zorlutuna, M. Akdag, W. K. Min and S. Atmaca, Remarks on soft topological spaces, *Ann. Fuzzy Math. Inform.* 3 (2) (2012) 171–185.
- [21] A. Aygunoglu and H. Aygun, Some notes on soft topological spaces, *Neural. Comput. Appl.* 21 (1) (2012) 113–119.
- [22] Sk. Nazmul and S. K. Samanta, Neighborhood properties of soft topological spaces, *Ann. Fuzzy Math. Inform.* 6 (1) (2013) 1–15.
- [23] D. N. Georgiou and A. C. Megaritis, Soft set theory and topology, *Applied General topology* 14 (2013) 93–100.
- [24] D. N. Georgiou, A. C. Megaritis and V. I. Petropoulos, On soft topological spaces, *Appl. Math. Inf. Sci.* 7 (5) (2013) 1889–1901.
- [25] Sadi Bayramov and Çiğdem Gündüz Aras, A new approach to separability and compactness in soft topological spaces, *TWMS J. Pure Appl. Math.* 9 (1) (2018) 82–93.
- [26] P. Debnath and B. C. Tripathy, On separation axioms in soft bitopological spaces, *Songklanakarin Journal of Science and Technology* 42 (4) (2020) 830–835.
- [27] Sk. Nazmul and S. K. Samanta, Soft topological soft groups, *Math. Sci. (Springer)*, 6:Art. 66, 10, 2012.
- [28] Sk. Nazmul and S. K. Samanta, Group soft topology, *The Journal of Fuzzy Mathematics* 22 (2) (2014) 435–450.
- [29] Takanori Hida, Soft topological group, *Ann. Fuzzy Math. Inform.* 8 (6) (2014) 1001–1025.
- [30] Mohammad K. Tahat, Fawzan Sidky and M. Abo-Elhamayel, Soft topological soft groups and soft rings, *Soft Computing* 22 (21) (2018) 7143–7156.
- [31] Mohammad K. Tahat, Fawzan Sidky and M. Abo-Elhamayel, Soft topological rings, *Journal of King Saud University-Science* 31 (4) (2019) 1127–1136.
- [32] Fawzan Sidky, M. E. El-Shafei and M. K. Tahat, Soft topological soft modules, *Ann. Fuzzy Math. Inform.* 20 (3) (2020) 257–272.
- [33] Sadi Bayramov, Çiğdem Gündüz Aras and Ljubiša D. R. Kočinac, Interval-valued topology on soft sets, *Axioms* 2023, 12, 692. <https://doi.org/10.3390/axioms12070692>.
- [34] J. G. Lee, G. Şenel, Y. B. Jun, Fadhil Abbas, K. Hur, Topological structures via interval-valued soft sets, *Ann. Fuzzy Math. Inform.* 22 (2) (2021) 133–169.
- [35] Metin Akdag and Alkan Ozkan, On soft α -separation axioms, *Journal of Advanced Studies in Topology* 5 (4) (2014) 16–24.
- [36] Metin Akdag and Alkan Ozkan, Soft α -pen sets and soft α -continuous functions, *Abstract and Applied Analysis* 2014 (2014) Article ID 891341 7 pages, <http://dx.doi.org/10.1155/2014/891341>.
- [37] J. Kim, Y. B. Jun, J. G. Lee, K. Hur, Topological structures based on interval-valued sets, *Ann. Fuzzy Math. Inform.* 20 (3) (2020) 273–295.
- [38] Y. Yao, Interval sets and interval set algebras, *Proc. 8th IEEE Int. Conf. on Cognitive Informatics (ICCI'09)* (2009) 307–314.
- [39] L. A. Zadeh, The concept of a linguistic variable and its application to approximate reasoning-I, *Inform. Sci.* 8 (1975) 199–249.
- [40] X. Yang, T. Y. Lin, J. Yang, Y. Li and D. Yu, Combination of interval-valued fuzzy set and soft set, *Computers & Mathematics with Applications* 58 (3) (2009) 521–527.

- [41] Mabruka Ali, A. Kiliçman and Azadeh Zahedi Khameneh, Separation axioms interval-valued fuzzy soft topology via quasi-neighborhood structure, Mathematics 2020,8,178;doi:10.3390/math8020178.
- [42] J. C. Kelly, Bitopological spaces, Proc. London Math. Soc. 13 (1963) 71–89.
- [43] J. G. Lee, G. Şenel, S. M. Mostafa, J. Kim and K. Hur, Continuities and separation axioms in interval-valued topological spaces, Ann. Fuzzy Math. Inform. 25 (1) (2023) 55–89.

J. I. BAEK (jibaek@wku.ac.kr)

School of Big Data and Financial Statistics, Wonkwang University, Korea

G. ŞENEL (g.senel@amasya.edu.tr)

Department of Mathematics, University of Amasya, Turkey

S. JAFARI (jafaripersia@gmail.com, saeidjafari@topositus.com)

Dr.rer.nat in Mathematics (Graz University of Technology-Graz, Austria)

Mathematical and Physical Science Foundation Sidevej 5 4200 Slagelse Denmark

S. H. HAN (shhan235@wku.ac.kr)

Department of Applied Mathematics, Wonkwang University, Korea

K. HUR (kulhur@wku.ac.kr)

Division of Applied Mathematics, Wonkwang University, 460, Iksan-daero, Iksan-Si, Jeonbuk 54538, Korea