

@FMII

# Soft set theory applied to residuated multilattices

Diagram illustrating the recursive construction of the Sierpinski triangle using the word "atomi". The root node is "atomi" (yellow box). It branches into "at" and "omi" (pink boxes). "at" branches into "a" and "t" (light blue boxes). "omi" branches into "o" and "mi" (light blue boxes). "a" branches into "a" and "t" (light blue boxes). "t" branches into "t" and "i" (light blue boxes). "o" branches into "o" and "m" (light blue boxes). "m" branches into "m" and "i" (light blue boxes). The final row shows the full expansion: "a", "t", "i", "m", "i" (light blue boxes).

Reprinted from the  
Annals of Fuzzy Mathematics and Informatics  
Vol. 29, No. 1, February 2025

## Soft set theory applied to residuated multilattices

KOUANKAM DJOUOHOU LÉA AUBIN, KOGUEP NJIONOU BLAISE BLÉRIOT, LÉLÉ  
CÉLESTIN

Received 22 May 2024; Revised 24 July 2024; Accepted 11 September 2024

**ABSTRACT.** Dealing with uncertainty, imprecision and fuzziness, *Soft set theory* is one of many mathematical tools; it is a classical generalization of fuzzy set theory. Many researchers study the case of a universe set endowed with an algebraic structure. This paper uses residuated multilattices as the universe set for soft set theory. The notions of *f*-soft residuated multilattice, *r*-soft residuated multilattice, soft filter, filteristic residuated multilattice are defined, and several related properties are investigated.

2020 AMS Classification: 03E72, 08A72

**Keywords:** Residuated multilattice, Soft set, *f*-soft residuated multilattice, *r*-soft residuated multilattice, Soft filter.

**Corresponding Author:** Kouankam Djouohou Léa Aubin ([akouankam@gmail.com](mailto:akouankam@gmail.com))

### 1. INTRODUCTION

Uncertainties occur in all areas of science and must therefore be managed rather than avoided. Theories such as probability, fuzzy sets [1], vague sets [2] and rough sets [3] are mathematical tools for handling uncertainties. As pointed out in [4], the major drawback of all these theories is the lack of consideration of parameters. That is why, soft set theory [4] was proposed by Molodtsov in 1999 to deal with uncertainty in a parametric manner. A soft set is defined as a parameterized family of sets which can be considered as an approximated description of an object, precisely consisting of two parts, namely, predicate and approximate value set. Because of its great potential, soft set theory has been rapidly developed with several applications. In [5, 6], a theoretical study is carried out on soft sets. A survey on some of the main developments of applications of soft sets theory in decision-making problems is given in [4, 7, 8]. Applications of soft set theory to some algebraic structures such as groups [9], rings [10, 11], lattices [12, 13], BCK/BCI-algebras [14], BL-algebras [15] and to general algebras in [16].

One of the most widely used algebraic theories is lattice theory. A lattice is a poset where every pair of elements has a least upper bound and a greatest lower bound. However, there are posets in which the set of upper (resp. lower) bounds of a pair of elements has no smallest (resp. greatest) element, but minimal (resp. maximal) elements. Benado formalized this in [17] by extending the notion of lattice to that of multilattice by weakening the uniqueness condition regarding upper and lower bounds to the existence of minimal upper bounds and maximal lower bounds. Residuation plays an important role in the algebraic study of logical systems; this lead to logical algebraic structures, i.e., algebraic structures modeling logical systems. The most widely used and important of these are residuated lattices. Focusing on the link between the theory of multilattices and residuation, Cabrera et al. introduce residuated multilattices [18]. Some authors have used multilattices and residuated multilattices in the framework of fuzzy sets theory [19, 20].

In [21], we applied soft set theory to multilattices. Now, we focus on residuated multilattices in the framework of soft sets. We defined the notions of *f-soft residuated multilattice*, *r-soft residuated multilattice*, *soft filter*, *filteristic soft residuated multilattice* and studied their related properties. The paper is organized as follows: in Section 2, we set some preliminaries on soft sets and residuated multilattices. In Section 3, we defined the notions of *f-soft residuated multilattice* and *r-soft residuated multilattice* and study their properties related to operations on soft sets. We focus on soft filter of an *f-soft residuated multilattice* in Section 4 while Section 5 deals with filteristic soft residuated multilattice. Finally, we draw some conclusions and prospects for future work in Section 6.

## 2. PRELIMINARIES

Given an initial universe set  $U$  and a set  $E$  of parameters, Molodtsov [4] defines the notion on *soft set* over  $U$  as a pair  $(f, E)$  such that  $f$  is a mapping from  $E$  to the power set  $P(U)$  of  $U$ . Thus, a soft set over a universe  $U$  is a parameterized family of subsets of  $U$ . To define a soft set over a universe  $U$  means, for each parameter, to point out the set of all elements of  $U$  that possess the given parameter. Of course, this set can be empty for some parameters. For a soft set  $(f, E)$ , the set of parameters with a non-empty image subset of  $U$  is called the *support* of  $(f, E)$  and denoted by  $\text{supp}(f, E)$ . Then  $\text{supp}(f, E) = \{p \in E : f(p) \neq \emptyset\}$ . It is often the practice to define a soft set with a part of the set of parameters rather than the whole parameter's set. In what follows, we denote a soft set  $(f, E)$  with support  $A$  by  $f_A$ . If the support  $A$  is empty, then  $f_\emptyset$  is the *null soft set* and will be denoted by  $\Phi$ .

Let us recall some basics operators on soft sets (one can refer to [6]). Let  $f_A$  and  $h_B$  be two soft sets over a universe  $U$  with the set of parameters  $E$ . Then  $f_A$  is called a *soft subset* of  $h_B$ , if  $A \subseteq B$  and  $f(p) \subseteq h(p)$  for all  $p \in A$ . In this case,  $h_B$  is a *soft supper set* of  $f_A$ .  $f_A$  and  $h_B$  are said to be *soft equal*, if  $f_A$  is a soft subset and a soft supper set of  $h_B$ . The *intersection* of  $f_A$  and  $h_B$  denoted by  $f_A \cap h_B$  is the soft set  $(t, E)$ , where  $t(p) = f(p) \cap h(p)$  for all  $p \in E$ . Clearly, if  $A \cap B = \emptyset$ , then  $f_A \cap h_B$  is the null soft set. If  $f_A \cap h_B$  is non-null, then its support is the set  $\{p \in A \cap B : f(p) \cap h(p) \neq \emptyset\}$ . The *union* of the soft sets  $f_A$  and  $h_B$  is the soft set  $(t, E)$ , where  $t(p) = f(p) \cup h(p)$  for all  $p \in E$ . As shown in [5], it is possible to restrict the parameter set of the union or extend that of the intersection of two

soft sets and obtain a *restricted union*  $\cup_r$  by considering the intersection of the sets of parameters or an *extended intersection*  $\cap_e$  by considering the union of the sets of parameters. Clearly, the restricted union of soft sets  $f_A$  and  $h_B$  over a common universe  $U$  denoted by  $f_A \cup_r h_B$  is the soft set  $t_{A \cap B}$  such that  $t(p) = f(p) \cup h(p)$  if  $p \in A \cap B$  and the extended intersection of  $f_A$  and  $h_B$  denoted by  $f_A \cap_e h_B$  is the

soft set  $t_{A \cup B}$  such that for all  $p \in A \cup B$ ,  $t(p) = \begin{cases} f(p) & \text{if } p \in A \setminus B \\ h(p) & \text{if } p \in B \setminus A \\ f(p) \cap h(p) & \text{else} \end{cases}$

and the restricted union of  $f_A$  and  $h_B$  with  $A \cap B \neq \emptyset$  is the soft set  $t_{A \cap B}$  such that for all  $p \in A \cap B$ ,  $t(p) = f(p) \cup h(p)$ .

Given two soft sets  $f_A$  and  $h_B$  over a common universe parameterized by a set  $E$ , one can define some soft sets with the set of parameters being a subset of  $E \times E$ . The soft set “ $f_A$  AND  $h_B$ ”, denoted by  $f_A \wedge h_B$  is defined by  $f_A \wedge h_B = (t, A \times B)$ , where  $t(p, q) = f(p) \cap h(q)$  for all  $(p, q) \in E \times E$ . While “ $f_A$  OR  $h_B$ ”, denoted by  $f_A \vee h_B$  is defined by  $f_A \vee h_B = (s, A \times B)$ , with  $s(p, q) = f(p) \cup h(q)$ , for all  $(p, q) \in A \times B$ .


Additional operations such as difference, restricted difference, symmetric difference and restricted symmetric difference of two soft sets can be found in [5, 21].

Now, let us give some preliminaries on residuated multilattices (See [17, 22]). Multilattices are natural generalization of lattices. Given a partially ordered set (poset)  $(P, \leq)$ ,  $x$  and  $y$  two elements of  $P$  such that  $x \leq y$ , we say that  $x$  is *below*  $y$  or  $y$  is *above*  $x$ . A poset  $(P, \leq)$  is called a *multilattice*, if for any finite subset  $X$  of  $P$ , each upper bound of  $X$  is above a minimal upper bound of  $X$  and each lower bound of  $X$  is below a maximal lower bound of  $X$ . We denote  $x \sqcap y$  (resp.  $x \sqcup y$ ) the set of minimal upper bounds (resp. maximal lower bounds) of  $\{x, y\}$ . We will sometimes write  $x \sqcap y = a$  instead of  $x \sqcap y = \{a\}$  and  $x \sqcup y = b$  instead of  $x \sqcup y = \{b\}$  when  $x \sqcap y$  or  $x \sqcup y$  is a singleton. A multilattice is said to be *full*, if  $x \sqcap y$  and  $x \sqcup y$  are non-empty for all  $x, y$ . A map  $\varphi : M \rightarrow N$  between two multilattices is said to be a *homomorphism* [17] if  $\varphi(x \sqcap y) \subseteq \varphi(x) \sqcap \varphi(y)$  and  $\varphi(x \sqcup y) \subseteq \varphi(x) \sqcup \varphi(y)$  for all  $x, y \in M$ . The authors in [18] showed that when the initial multilattice is full, the notion of homomorphism can be characterized in terms of equalities as follows: let  $\varphi : M \rightarrow N$  be a map between multilattices such that  $M$  is full. Then  $\varphi$  is a homomorphism iff  $\varphi(x \sqcap y) = (\varphi(x) \sqcap \varphi(y)) \cap \varphi(M)$  and  $\varphi(x \sqcup y) = (\varphi(x) \sqcup \varphi(y)) \cap \varphi(M)$  for all  $x, y \in M$ .

A structure  $\mathcal{P} := (P, \leq, \top, \odot, \rightarrow)$ , is said to be *pocrim* (partially ordered commutative residuated integral monoid), if  $(P, \leq, \top)$  is a poset with a greatest element  $\top$  and  $(P, \odot, \top)$  is a commutative monoid such that

$$(2.1) \quad x \odot y \leq z \Leftrightarrow x \leq y \rightarrow z \quad \text{for all } x, y, z \in P.$$

Then the pair  $(\odot, \rightarrow)$  is called the *residuation*. If a pocrim has a least element, usually denoted by  $\perp$ , it is said to be *bounded*. A *residuated multilattice* ( $\mathcal{RML}$  for short) is a multilattice endowed with a residuation. It combines a structure of pocrim and a structure of multilattice. Quite simply, it is a pocrim whose underlying poset is a multilattice [18]. When the underlying poset is a lattice, the pocrim is called a *residuated lattice*. Clearly, a residuated lattice is an  $\mathcal{RML}$ . When an  $\mathcal{RML}$  is not a residuated lattice, it is said to be *pure*. The authors in [23] show that every


 FIGURE 1. A residuated multilattice with seven elements: *RML-7*.

bounded pure  $\mathcal{RML}$  has at least seven elements. Figure 1 gives the  $\mathcal{RML}$  with seven elements *RML-7* from [23] that we will use throughout the paper.

Let  $\mathcal{P}$  be an  $\mathcal{RML}$  and  $X$  be a subset of  $\mathcal{P}$ .  $X$  is called a *full residuated submultilattice* (*f-sub- $\mathcal{RML}$* ), if  $X$  contains  $\top$ ,  $x \odot y, x \rightarrow y$ , and  $x \sqcap y, x \sqcup y \subseteq X$  for all  $x, y \in X$ . Similarly,  $X$  is called a *restricted residuated submultilattice* (*r-sub- $\mathcal{RML}$* ), if  $X$  contains  $\top$ ,  $x \odot y, x \rightarrow y$  and  $(x \sqcap y) \cap X \neq \emptyset, (x \sqcup y) \cap X \neq \emptyset$  for all  $x, y \in X$ . For example,  $\{a, c, e, \top\}$  is an *f-sub- $\mathcal{RML}$*  and an *r-sub- $\mathcal{RML}$*  of *RML-7*.  $\{a, c, d, e, \top\}$  is an *r-sub- $\mathcal{RML}$*  of *RML-7* which is not an *f-sub- $\mathcal{RML}$* , since  $c \sqcap d = \{a, b\}$  and  $b \notin \{a, c, d, e, \top\}$ . One can remark that, as for Multilattices, an *f-sub- $\mathcal{RML}$*  of an  $\mathcal{RML}$  is an  $\mathcal{RML}$  on its own. Moreover, any intersection of two *f-sub- $\mathcal{RML}$* s is also an *f-sub- $\mathcal{RML}$* , but it is not the case for *r-sub- $\mathcal{RML}$* s. Indeed,  $X = \{\perp, a, b, c, e, \top\}$  and  $Y = \{\perp, a, b, d, e, \top\}$  are *r-sub- $\mathcal{RML}$* s, but  $X \cap Y = \{\perp, a, b, e, \top\}$  is not an *r-sub- $\mathcal{RML}$* , since  $a \sqcup b = \{c, d\}$  and  $\{c, d\} \cap (X \cap Y) = \emptyset$ .

We end this section by recalling some useful properties that hold in a pocrim in general and in an  $\mathcal{RML}$  in particular. We consider a pocrim  $\mathcal{P}$  which is an  $\mathcal{RML}$ . Let  $x, y, z \in \mathcal{P}$ .

- P1**  $x \rightarrow x = \top$  and  $\top \rightarrow x = x$ ,
- P2**  $x \odot y \leq x$  and  $x \odot y \leq y$ , then  $x \leq y \rightarrow x$ ,
- P3**  $x \odot y \leq x \rightarrow y$ ,
- P4**  $x \leq y \Leftrightarrow x \rightarrow y = \top$ ,
- P5**  $x \rightarrow y = y \rightarrow x = \top \Leftrightarrow x = y$ .

### 3. *f*-SOFT RESIDUATED MULTILATTICE AND *r*-SOFT RESIDUATED MULTILATTICE

Throughout this section and the rest of the study, we will consider an  $\mathcal{RML}$   $\mathcal{M} := (M, \leq, \top, \odot, \rightarrow)$  as our universe set and  $E$  a set of parameters. To simplify, we will write  $M$  to denote the support or the whole residuated multilattice.

**Definition 3.1.** Let  $M$  be an  $\mathcal{RML}$  and  $f_A$  be a non-null soft set over  $M$ . Then  $f_A$  said to be:

- (i) *full soft residuated multilattice* (briefly, *f-soft- $\mathcal{RML}$* ) over  $M$ , if  $f(p)$  is an *f-sub- $\mathcal{RML}$*  of  $M$  for all  $p \in A$ ,

(ii) *restricted soft residuated multilattice* (briefly,  $r$ -soft- $\mathcal{RML}$ ) over  $M$ , if  $f(p)$  is an  $r$ -sub- $\mathcal{RML}$  of  $M$  for all  $p \in A$ .

**Remark 3.2.** Let  $M$  be an  $\mathcal{RML}$  and  $x \in M$ . The following statements hold:

- (1)  $\{\top\}$  is an  $f$ -sub- $\mathcal{RML}$  of  $M$ ,
- (2)  $\{x, \top\}$  is an  $f$ -sub- $\mathcal{RML}$  of  $M$  iff  $x \odot x = x$ .

**Example 3.3.** Consider  $RML-7$  in Figure 1 and  $E = \{p_1, p_2, p_3\}$ . Let  $f_A$  and  $h_B$  be the soft sets over  $RML-7$  defined by

$$\begin{aligned} f_A &= \{(p_1, \{\perp, \top\}), (p_2, \{a, \top\})\}, \\ h_B &= \{(p_1, \{\perp, a, b, c, e, \top\}), (p_2, \{\perp, b, d, e, \top\})\}, \\ t_C &= \{(p_1, \{\perp, \top\}), (p_2, \{\perp, a, \top\})\}. \end{aligned}$$

Then we have  $f(p_1) = \{\perp, \top\}$ ,  $f(p_2) = \{a, \top\}$ . By Remark 3.2,  $f(p_1)$  and  $f(p_2)$  are  $f$ -sub- $\mathcal{RML}$ s of  $RML-7$ . Thus  $f_A$  is an  $f$ -soft- $\mathcal{RML}$  over  $RML-7$ . However  $h(p_1) = \{\perp, a, b, c, e, \top\}$  is not an  $f$ -sub- $\mathcal{RML}$  of  $RML-7$ , as  $a \sqcup b = \{c, d\}$  and we have  $\{c, d\} \not\subseteq \{\perp, a, b, c, e, \top\}$ . So  $h_B$  is not an  $f$ -soft- $\mathcal{RML}$  over  $RML-7$ . Nevertheless,  $h_B$  is an  $r$ -soft- $\mathcal{RML}$  over  $RML-7$ .

$t_C$  is neither an  $f$ -soft- $\mathcal{RML}$  nor an  $r$ -soft- $\mathcal{RML}$  because  $t(p_2) = \{\perp, a, \top\}$  and  $a \rightarrow \perp = b \notin t(p_2)$ .

**Remark 3.4.** Let  $M$  be an  $\mathcal{RML}$  and  $f_A$  be an  $f$ -soft- $\mathcal{RML}$  (resp.  $r$ -soft- $\mathcal{RML}$ ) over  $M$ . If  $\emptyset \neq B \subseteq A$ , then  $f_B$  is an  $f$ -soft- $\mathcal{RML}$  (resp.  $r$ -soft- $\mathcal{RML}$ ) over  $M$ .

Knowing that the intersection of  $f$ -sub- $\mathcal{RML}$ s of an  $\mathcal{RML}$   $M$  is also an  $f$ -sub- $\mathcal{RML}$  of  $M$ , we have the following result concerning the intersection of two  $f$ -soft- $\mathcal{RML}$ s over an  $\mathcal{RML}$   $M$ , whose proof is straightforward.

**Proposition 3.5.** Let  $M$  be an  $\mathcal{RML}$ ,  $f_A$  and  $h_B$  be two non-null  $f$ -soft- $\mathcal{RML}$ s over  $M$ . Then  $f_A \cap h_B$  and  $f_A \wedge h_B$  are  $f$ -soft- $\mathcal{RML}$ s over  $M$  if they are non-null.

We have shown in Section 2 that the intersection of  $r$ -sub- $\mathcal{RML}$ s of an  $\mathcal{RML}$   $M$  is not always an  $r$ -sub- $\mathcal{RML}$ . This is also the case for the intersection of  $r$ -soft- $\mathcal{RML}$ s. Indeed, by considering the  $\mathcal{RML}$  in Figure 1,  $f_A = \{(p_1, \{\perp, a, b, c, e, \top\})\}$  and  $h_B = \{(p_1, \{\perp, a, b, d, e, \top\})\}$ , clearly,  $f_A$  and  $h_B$  are  $r$ -soft- $\mathcal{RML}$ s over  $RML-7$  but  $f_A \cap h_B$  is not since  $f(p_1) \cap h(p_1) = \{\perp, a, b, e, \top\}$  is not an  $r$ -sub- $\mathcal{RML}$  of  $RML-7$ . Moreover,  $f_A \wedge h_B = \{(p_1, p_1), \{\perp, a, b, e, \top\})\}$  is not an  $r$ -soft- $\mathcal{RML}$ .

**Proposition 3.6.** Let  $M$  be an  $\mathcal{RML}$ ,  $f_A$  and  $h_B$  be two non-null  $f$ -soft- $\mathcal{RML}$ s (resp.  $r$ -soft- $\mathcal{RML}$ s) over  $M$ . If  $A \cap B = \emptyset$ , then  $f_A \cup h_B$  is an  $f$ -soft- $\mathcal{RML}$  (resp.  $r$ -soft- $\mathcal{RML}$ ) over  $M$ .

*Proof.* We know that  $f_A \cup h_B$  is the soft set  $(t, E)$ , where  $t(p) = f(p) \cup h(p)$  for all  $p \in A \cup B$  and  $t(p) = \emptyset$  if  $p \notin A \cup B$ . Suppose that  $f_A$  and  $h_B$  are  $f$ -soft- $\mathcal{RML}$ s over  $M$  and  $A \cap B = \emptyset$ . Then for all  $p \in A \cup B$ , either  $t(p) = f(p)$  or  $t(p) = h(p)$ , as  $A \cup B = (A \setminus B) \cup (B \setminus A)$ . As  $f_A$  and  $h_B$  are  $f$ -soft- $\mathcal{RML}$ s over  $M$ , if  $p \in A$ , then  $f(p)$  is an  $f$ -sub- $\mathcal{RML}$  of  $M$  and if  $p \in B$ , then  $h(p)$  is an  $f$ -sub- $\mathcal{RML}$  of  $M$ . That is  $t(p)$  is an  $f$ -sub- $\mathcal{RML}$  of  $M$  for all  $p \in A \cup B$ . This implies that  $f_A \cup h_B$  is an  $f$ -soft- $\mathcal{RML}$  over  $M$  as required.

The proof for  $r$ -soft- $\mathcal{RML}$  can be obtained similarly.  $\square$

If  $A \cap B \neq \emptyset$ , then  $f_A \cup h_B$  may not be an  $f$ -soft- $\mathcal{RML}$ . Let us consider, for example  $f_A = \{(p_1, \{\perp, \top\})\}$  and  $h_B = \{(p_1, \{a, \top\})\}$  which are  $f$ -soft- $\mathcal{RML}$ s over RML-7. We have  $f_A \cup h_B = \{(p_1, \{\perp, a, \top\})\}$  which is not an  $f$ -soft- $\mathcal{RML}$  because  $\{\perp, a, \top\}$  is not an  $f$ -sub- $\mathcal{RML}$  of RML-7, since  $a \rightarrow \perp = b \notin \{\perp, a, \top\}$ . However,  $f_A \cup h_B$  can still be an  $f$ -soft- $\mathcal{RML}$  even if  $A \cap B \neq \emptyset$ , as explained in Proposition 3.7, whose proof is similar to Proposition 3.6.

**Proposition 3.7.** *Let  $M$  be an  $\mathcal{RML}$ ,  $f_A$  and  $h_B$  be two non-null  $f$ -soft- $\mathcal{RML}$ s (resp.  $r$ -soft- $\mathcal{RML}$ s) over  $M$  such that for all  $p \in E$ ,  $f(p) \subseteq h(p)$  or  $h(p) \subseteq f(p)$ . Then  $f_A \cup h_B$  is an  $f$ -soft- $\mathcal{RML}$  (resp.  $r$ -soft- $\mathcal{RML}$ ) over  $M$ .*

If  $f_A$  and  $h_B$  are two non-null  $f$ -soft- $\mathcal{RML}$ s over an  $\mathcal{RML}$   $M$ ,  $f_A \vee h_B$  may not be an  $f$ -soft- $\mathcal{RML}$  even if  $A \cap B = \emptyset$ . Indeed, from the  $\mathcal{RML}$  of Figure 1, let us consider  $f_A = \{(p_1, \{\perp, \top\})\}$  and  $h_B = \{(p_2, \{a, \top\})\}$ . By Remark 3.2,  $f_A$  and  $h_B$  are  $f$ -soft- $\mathcal{RML}$ s over RML-7. However,  $f_A \vee h_B = \{((p_1, p_2), \{\perp, a, \top\})\}$  is not an  $f$ -soft- $\mathcal{RML}$ . One can also observe that  $f_A$  and  $h_B$  are  $r$ -soft- $\mathcal{RML}$ s but  $f_A \vee h_B$  is not.

**Proposition 3.8.** *Let  $M$  be an  $\mathcal{RML}$ ,  $f_A$  and  $h_B$  be two non-null  $f$ -soft- $\mathcal{RML}$ s over  $M$ . Then  $f_A \cap_g h_B$  is an  $f$ -soft- $\mathcal{RML}$  over  $M$  if it is non-null.*

*Proof.* If  $f_A \cap_g h_B$  is non-null, then  $f_A \cap_g h_B$  is the soft set  $(t, E)$  such that for all

$$p \in A \cup B, t(p) = \begin{cases} f(p) & \text{if } p \in A \setminus B \\ h(p) & \text{if } p \in B \setminus A \\ f(p) \cap h(p) & \text{else} . \end{cases}$$

In each case,  $t(p)$  is an  $f$ -sub- $\mathcal{RML}$  of  $M$ . We conclude that  $f_A \cap_g h_B$  is an  $f$ -soft- $\mathcal{RML}$  over  $M$ .  $\square$

#### 4. SOFT FILTER OF AN $f$ -SOFT RESIDUATED MULTILATTICE

The notion of filter, originated in topology, is very useful in various algebraic structures endowed with a (partial) order. As stated above, the structure of residuated multilattice is a combination of two structures, namely the structure of pocrim and the structure of multilattice. In what follows, we will consider the notion of filter in the structure of pocrim that we will call p-filter, the notion of filter in a multilattice that we will call m-filter and we will call filter the one that combines the structures of pocrim and multilattice. In this section, we extend this notion to the framework of soft set theory.

Let recall from [18] the definition and some properties on filters in residuated multilattices.

**Definition 4.1.** Let  $M$  be an  $\mathcal{RML}$ . A non-empty subset  $F \subseteq M$  is said to be a  $p$ -filter of  $M$ , if the following conditions hold:

- (i) if  $x, y \in F$ , then  $x \odot y \in F$ ,
- (ii) if  $x \in F$  and  $x \leq y$ , then  $y \in F$ .

Note that if  $F$  is a p-filter of an  $\mathcal{RML}$   $M$ , then  $\top \in F$ . Moreover, if  $x, x \rightarrow y \in F$ , then  $y \in F$ . A subset  $D \subseteq M$  containing  $\top$  and satisfying  $(x, x \rightarrow y \in D) \implies y \in D$  is called a *deductive system*. Thus a p-filter is a deductive system. The converse is also true.

**Definition 4.2.** Let  $M$  be an  $\mathcal{RML}$ . A non-empty set  $F \subseteq M$  is said to be an *m-filter*, if the following conditions hold:

- (i) for all  $x, y \in F$ ,  $\emptyset \neq x \sqcap y \subseteq F$ ,
- (ii) for all  $x \in F$  and  $y \in M$ ,  $x \sqcup y \subseteq F$ ,
- (iii) for all  $x, y \in M$ , if  $(x \sqcup y) \cap F \neq \emptyset$ , then  $x \sqcup y \subseteq F$ .

Cabrera et al. [18] have proposed a notion of filter in the  $\mathcal{RML}$  structure which combines pocrim and multilattice structures.

**Definition 4.3.** Let  $M$  be an  $\mathcal{RML}$ . A non-empty subset  $F \subseteq M$  is said to be a *filter*, if it is a deductive system and the following condition holds:

$$x \rightarrow y \in F \text{ implies } (x \sqcup y) \rightarrow y \subseteq F \text{ and } x \rightarrow (x \sqcap y) \subseteq F.$$

It is known that a filter of an  $\mathcal{RML}$  is an m-filter and a p-filter but the converse is not true [18]. One can easily observe that:

**Remark 4.4.** Let  $M$  be an  $\mathcal{RML}$ . The following conditions hold:

- (1)  $\{\top\}$  and  $M$  are filters of  $M$ ,
- (2) if  $F$  is a filter of  $M$ , then  $F$  is an  $f$ -sub- $\mathcal{RML}$  of  $M$ .

**Proposition 4.5.** Let  $M$  be an  $\mathcal{RML}$ ,  $F$  and  $G$  be two filters of  $M$ . Then  $F \cap G$  is a filter of  $M$ .

One can easily observe that if  $X$  is an  $f$ -sub- $\mathcal{RML}$  of an  $\mathcal{RML}$   $M$ , then  $X$  is an  $\mathcal{RML}$  on its own right. Then we introduce the notion of soft filter of an  $f$ -soft- $\mathcal{RML}$  as follows:

**Definition 4.6.** Let  $h_F$  be a non-null soft set over an  $\mathcal{RML}$   $M$  and  $f_A$  be an  $f$ -soft- $\mathcal{RML}$  over  $M$ . Then  $h_F$  is called a *soft filter* of  $f_A$ , if the following conditions hold:

- (i)  $F \subseteq A$ ,
- (ii) for all  $p \in F$ ,  $h(p)$  is a filter of  $f(p)$ .

**Example 4.7.** Let  $M$  be the  $\mathcal{RML}$  described in Figure 2. The soft set  $f_A$  over  $M$  defined by  $f_A = \{(p_1, \{g, h, \top\}), (p_2, \{f, \top\}), (p_3, \{\perp, \top\})\}$  is clearly an  $f$ -soft- $\mathcal{RML}$  over  $M$ .  $h_F = \{(p_1, \{h, \top\}), (p_2, \{\top\})\}$  is a soft filter of  $f_A$ .


Using Remark 4.4, one can observe that any soft filter is an  $f$ -soft- $\mathcal{RML}$  over  $M$ . Let us now give some properties of soft filters.

**Proposition 4.8.** Let  $f_A$  be an  $f$ -soft- $\mathcal{RML}$  over an  $\mathcal{RML}$   $M$ . Let  $h_F$  and  $t_G$  be two soft filters of  $f_A$ . Then  $h_F \cap t_G$  is a soft filter of  $f_A$  if it is non-null.

*Proof.* Suppose  $h_F \cap t_G$  is non-null. Then  $h_F \cap t_G = u_C$  with  $C = \{p \in F \cap G : h(p) \cap t(p) \neq \emptyset\}$  and  $u(p) = h(p) \cap t(p)$  for all  $p \in C$ . As  $h_F$  and  $t_G$  are soft filters of  $f_A$ , we have  $F \subseteq A$  and  $G \subseteq A$ , that is,  $C \subseteq A$ . If  $p \in C$ , then  $p \in F$  and  $p \in G$ . This implies that  $h(p)$  and  $t(p)$  are filters of  $f(p)$ . By Proposition 4.5, we deduce that  $u(p) = h(p) \cap t(p)$  is a filter of  $f(p)$ . It follows that  $u_C$  is a soft filter of  $f_A$ .  $\square$

The union of two soft filters of an  $f$ -soft- $\mathcal{RML}$  is not always a soft filter. Indeed, let us consider  $M$  as in Example 4.7 and define  $f_A = \{(p_1, M)\}$  which is an  $f$ -soft- $\mathcal{RML}$  over  $M$ . The soft sets  $h_F = \{(p_1, \{g, h, \top\})\}$  and  $t_G = \{(p_1, \{f, h, \top\})\}$  are





$\odot$	$\perp$	$a$	$b$	$c$	$d$	$e$	$f$	$g$	$h$	$\top$
$\perp$	$\perp$	$\perp$	$\perp$	$\perp$	$\perp$	$\perp$	$\perp$	$\perp$	$\perp$	$\perp$
$a$	$\perp$	$\perp$	$\perp$	$\perp$	$\perp$	$\perp$	$\perp$	$\perp$	$\perp$	$a$
$b$	$\perp$	$\perp$	$\perp$	$\perp$	$\perp$	$\perp$	$\perp$	$\perp$	$\perp$	$b$
$c$	$\perp$	$\perp$	$\perp$	$\perp$	$\perp$	$\perp$	$\perp$	$\perp$	$\perp$	$c$
$d$	$\perp$	$\perp$	$\perp$	$\perp$	$\perp$	$\perp$	$\perp$	$\perp$	$\perp$	$d$
$e$	$\perp$	$\perp$	$\perp$	$\perp$	$\perp$	$\perp$	$\perp$	$\perp$	$\perp$	$e$
$f$	$\perp$	$\perp$	$\perp$	$\perp$	$\perp$	$\perp$	$f$	$\perp$	$f$	$f$
$g$	$\perp$	$\perp$	$\perp$	$\perp$	$\perp$	$\perp$	$\perp$	$g$	$g$	$g$
$h$	$\perp$	$\perp$	$\perp$	$\perp$	$\perp$	$\perp$	$f$	$g$	$h$	$h$
$\top$	$\perp$	$a$	$b$	$c$	$d$	$e$	$f$	$g$	$h$	$\top$

$\rightarrow$	$\perp$	$a$	$b$	$c$	$d$	$e$	$f$	$g$	$h$	$\top$
$\perp$	$\top$	$\top$	$\top$	$\top$	$\top$	$\top$	$\top$	$\top$	$\top$	$\top$
$a$	$h$	$\top$	$h$	$\top$	$\top$	$\top$	$\top$	$\top$	$\top$	$\top$
$b$	$h$	$h$	$\top$	$\top$	$\top$	$\top$	$\top$	$\top$	$\top$	$\top$
$c$	$h$	$h$	$h$	$\top$	$h$	$\top$	$\top$	$\top$	$\top$	$\top$
$d$	$h$	$h$	$h$	$h$	$\top$	$\top$	$\top$	$\top$	$\top$	$\top$
$e$	$h$	$h$	$h$	$h$	$h$	$\top$	$\top$	$\top$	$\top$	$\top$
$f$	$g$	$g$	$g$	$g$	$g$	$g$	$\top$	$g$	$\top$	$\top$
$g$	$f$	$f$	$f$	$f$	$f$	$f$	$f$	$\top$	$\top$	$\top$
$h$	$e$	$e$	$e$	$e$	$e$	$e$	$f$	$g$	$\top$	$\top$
$\top$	$\perp$	$a$	$b$	$c$	$d$	$e$	$f$	$g$	$h$	$\top$

FIGURE 2. Residuated multilattice with 10 elements

soft filters of  $f_A$  but  $h_F \cup t_G = \{(p_1, \{f, g, h, \top\})\}$  is not a soft filter of  $f_A$  because  $\{f, g, h, \top\}$  is not a filter of  $M$ , as  $f \sqcap g = e \notin \{f, g, h, \top\}$ . However, the following result gives a sufficient condition for the union of two soft filters of an  $f$ -soft- $\mathcal{RML}$  to be a soft filter.

**Proposition 4.9.** *Let  $f_A$  be an  $f$ -soft- $\mathcal{RML}$  over an  $\mathcal{RML}$   $M$ . Let  $h_F$  and  $t_G$  be two soft filters of  $f_A$ . If  $F \cap G = \emptyset$ , then  $h_F \cup t_G$  is a soft filter of  $f_A$ .*

*Proof.* We know that  $h_F \cup t_G$  is the soft set  $(u, E)$ , where  $u(p) = h(p) \cup t(p)$  for all  $p \in F \cup G$  and  $u(p) = \emptyset$  if  $p \notin F \cup G$ . As  $h_F$  and  $t_G$  are soft filters of  $f_A$ , we have  $F \subseteq A$  and  $G \subseteq A$ , which implies that  $F \cup G \subseteq A$ . If  $p \in F \cup G$ , then either  $p \in F \setminus G$  or  $p \in G \setminus F$ . That is,  $u(p) = h(p)$  or  $u(p) = t(p)$ . In both cases,  $t(p)$  is a filter of  $f(p)$ , as  $h_F$  and  $t_G$  are soft filters of  $f_A$ . We can conclude that  $h_F \cup t_G$  is a soft filter of  $f_A$ .  $\square$

**Proposition 4.10.** *Let  $f_A$  be an  $f$ -soft- $\mathcal{RML}$  over an  $\mathcal{RML}$   $M$ . Let  $h_F$  and  $t_G$  be two soft filters of  $f_A$ . Then  $h_F \sqcap_g t_G$  is a soft filter of  $f_A$  if it is non-null.*

*Proof.* Suppose  $h_F \cap_{\mathfrak{g}} t_G$  is non-null. Then  $h_F \cap_{\mathfrak{g}} t_G$  is the soft set  $(u, E)$ , where for all  $p \in F \cup G$ ,

$$u(p) = \begin{cases} h(p) & \text{if } p \in F \setminus G \\ t(p) & \text{if } p \in G \setminus F \\ h(p) \cap t(p) & \text{else.} \end{cases}$$

As  $h_F$  and  $t_G$  are soft filters of  $f_A$ ,  $F \subseteq A$  and  $G \subseteq A$ . That is,  $F \cup G \subseteq A$ . Knowing that the intersection of two filters of an  $\mathcal{RML}$  is also a filter (Proposition 4.5),  $u(p) = h(p) \cap t(p)$  is a filter of  $f(p)$ , if  $p \in F \cap G$ . For the others two cases,  $u(p)$  is evidently a filter of  $f(p)$ , as  $h_F$  and  $t_G$  are soft filters of  $f_A$ . It follows that  $h_F \cap_{\mathfrak{g}} t_G$  is a soft filter of  $f_A$ .  $\square$

## 5. FILTERISTIC SOFT RESIDUATED MULTILATTICE

In the previous section, we have presented the notion of soft filter, which is related to an  $f$ -soft- $\mathcal{RML}$ . In this section, we present the notion of filteristic soft residuated multilattice, which is related to the universe. We also look about the link between homomorphisms and filters.

**Definition 5.1.** A non-null soft set  $f_A$  over an  $\mathcal{RML}$   $M$  is called a *filteristic soft residuated multilattice* over  $M$ , if  $f(p)$  is a filter of  $M$  for all  $p \in A$ .

**Example 5.2.** Let  $M$  be the  $\mathcal{RML}$  of Figure 2.

$f_A = \{(p_1, \{f, h, \top\}), (p_2, \{g, h, \top\}), (p_3, \{h, \top\})\}$  is a filteristic soft residuated multilattice over  $M$ .

**Remark 5.3.** A soft filter of an  $f$ -soft- $\mathcal{RML}$  over an  $\mathcal{RML}$   $M$  is not always a filteristic soft residuated multilattice over  $M$ . Indeed, let  $M$  be the  $\mathcal{RML}$  of Figure 2. The soft set  $f_A = \{(p_1, M \setminus \{f, g\})\}$  is an  $f$ -soft- $\mathcal{RML}$  over  $M$  and  $h_F = \{(p_1, \{e, h, \top\})\}$  is a soft filter of  $f_A$  but it is not a filteristic soft residuated multilattice because  $h(p_1) = \{e, h, \top\}$  is not a filter of  $M$ .

**Proposition 5.4.** Let  $f_A$  and  $h_B$  be two filteristic soft residuated multilattices over an  $\mathcal{RML}$   $M$ . Then:

- (1)  $f_A \cap h_B$  is a filteristic soft residuated multilattice over  $M$  if it is non-null,
- (2) if  $A \cap B = \emptyset$ , then  $f_A \cup h_B$  is a filteristic soft residuated multilattice over  $M$ ,
- (3)  $f_A \wedge h_B$  is a filteristic soft residuated multilattice over  $M$  if it is non-null,
- (4)  $f_A \cap_{\mathfrak{g}} g_B$  is a filteristic soft residuated multilattice over  $M$  if it is non-null.

*Proof.* (1) The proof is similar to Proposition 4.8.

(2) The proof is similar to Proposition 4.9.

(3) The proof follows from Proposition 4.5.

(4) The proof is similar to Proposition 4.10.  $\square$

Given an  $\mathcal{RML}$   $M$ , we know that  $\{\top\}$  and  $M$  are filters of  $M$ . They are known as trivial filter and whole filter. We set a trivial and whole filteristic soft residuated multilattices to be as follows:

**Definition 5.5.** Let  $M$  be an  $\mathcal{RML}$ . Then a filteristic soft residuated multilattice  $f_A$  over  $M$  is said to be:

- (i) *trivial*, if  $f(p) = \{\top\}$  for all  $p \in A$ .
- (ii) *whole*, if  $f(p) = M$  for all  $p \in A$ .

In general, a homomorphism between residuated multilattices is a homomorphism of multilattice which preserves the multiplication ( $\odot$ ) and the implication ( $\rightarrow$ ). In what follows, we investigate the interaction between filters and homomorphisms.

According to the fact that an  $\mathcal{RML}$  is always full (as multilattice), we have the following definition:

**Definition 5.6** ([18]). Let  $M$  and  $N$  be two  $\mathcal{RML}$ s. A map  $\varphi : M \rightarrow N$  is said to be an *homomorphism*, if the following conditions hold: for all  $x, y \in M$ ,

- (i)  $\varphi(x \sqcap y) = (\varphi(x) \sqcap \varphi(y)) \sqcap \varphi(M)$  and  $\varphi(x \sqcup y) = (\varphi(x) \sqcup \varphi(y)) \sqcap \varphi(M)$ ,
- (ii)  $\varphi(x \odot y) = \varphi(x) \odot \varphi(y)$  and  $\varphi(x \rightarrow y) = \varphi(x) \rightarrow \varphi(y)$ .

One can easily observe that  $\varphi(\top) = \top$  for all homomorphism  $\varphi$  between residuated multilattices.

**Proposition 5.7.** [18] Let  $\varphi : M \rightarrow N$  be an homomorphism between residuated multilattices. Then:

- (1)  $\varphi^{-1}(\top)$  is a filter of  $M$ ,
- (2) if  $F$  is a filter of  $N$ , then  $\varphi^{-1}(F)$  is a filter of  $M$ .

**Proposition 5.8.** Let  $\varphi : M \rightarrow N$  be an homomorphism between residuated multilattices. If  $F$  is a filter of  $M$  and  $\varphi$  bijective, then  $\varphi(F)$  is a filter of  $N$ .

*Proof.* Suppose  $F$  is a filter of  $M$ . Note that as  $\varphi$  is surjective, we deduce from Definition 5.6 that  $\varphi(x \sqcap y) = \varphi(x) \sqcap \varphi(y)$  and  $\varphi(x \sqcup y) = \varphi(x) \sqcup \varphi(y)$  for all  $x, y \in M$ . Moreover, if  $x \leq y$ , then  $\varphi(x) = \varphi(x \sqcap y) = \varphi(x) \sqcap \varphi(y)$ , which implies that  $\varphi(x) \leq \varphi(y)$ . Then  $\varphi$  is an isotone function. Thus  $\varphi^{-1}$  is an isotone function.

- (i) Let  $z, t \in \varphi(F)$ . Then there exist  $x, y \in F$  such that  $z = \varphi(x)$  and  $t = \varphi(y)$ . Thus we have  $z \odot t = \varphi(x) \odot \varphi(y) = \varphi(x \odot y)$ . As  $F$  is a filter of  $M$ ,  $x \odot y \in F$ . That is,  $z \odot t = \varphi(x \odot y) \in \varphi(F)$ .
- (ii) Let  $z \in \varphi(F)$  and  $t \in N$  such that  $z \leq t$ . Then there exist  $x \in F$  such that  $\varphi(x) = z$  and by the surjectivity of  $\varphi$ , an element  $y \in M$  such that  $\varphi(y) = t$ . We have to show that  $t \in \varphi(F)$ . As  $\varphi^{-1}$  is isotone, we have from  $z \leq t$  that  $\varphi^{-1}(z) \leq \varphi^{-1}(t)$ . Thus  $\varphi^{-1}(\varphi(x)) \leq \varphi^{-1}(\varphi(y))$ , i.e.,  $x \leq y$ . As  $F$  is a filter of  $M$  and  $x \in F$ , we deduce that  $y \in F$ , which implies that  $t = \varphi(y) \in \varphi(F)$ .
- (iii) Let  $z, t \in N$  such that  $z \rightarrow t \in \varphi(F)$ . We will show that  $(z \sqcup t) \rightarrow t \subseteq \varphi(F)$  and  $z \rightarrow (z \sqcap t) \subseteq \varphi(F)$ . As  $\varphi$  is surjective, there exist  $x, y \in M$  such that  $z = \varphi(x)$  and  $t = \varphi(y)$ . From  $z \rightarrow t \in \varphi(F)$ , we have  $\varphi(x) \rightarrow \varphi(y) \in \varphi(F)$ , i.e.,  $\varphi(x \rightarrow y) \in \varphi(F)$ , which implies by the bijectivity of  $\varphi$  that  $x \rightarrow y \in F$ . Now, we have  $(z \sqcup t) \rightarrow t = (\varphi(x) \sqcup \varphi(y)) \rightarrow \varphi(y) = \varphi((x \sqcup y) \rightarrow y)$ . As  $x \rightarrow y \in F$ , we have  $(x \sqcup y) \rightarrow y \in F$ , which implies that  $(z \sqcup t) \rightarrow t = \varphi((x \sqcup y) \rightarrow y) \in \varphi(F)$ . Similarly, we obtain  $z \rightarrow (z \sqcap t) \subseteq \varphi(F)$ .

□

Let  $f_A$  be a soft set over an  $\mathcal{RML}$   $M$  and  $\varphi : M \rightarrow N$  be a map between  $\mathcal{RML}$ s. Let us consider the mapping  $\tilde{\varphi} : P(M) \rightarrow P(N)$  such that  $\tilde{\varphi}(X) = \varphi(X)$  (the direct image), for all  $X \in P(M)$ . We can see that  $(\tilde{\varphi} \circ f, E)$  is a soft set over  $N$ . If  $p \in A$ ,

then  $f(p) \neq \emptyset$ , i.e.,  $\varphi(f(p)) \neq \emptyset$ . If rather  $f(p) = \emptyset$  ( $p \notin A$ ), then  $\varphi(f(p)) = \emptyset$ . Thus  $f$  and  $(\tilde{\varphi} \circ f, E)$  have the same support  $A$ ; we will denote  $(\tilde{\varphi} \circ f, E)$  by  $\varphi f_A$  in the sequel.

**Proposition 5.9.** *Let  $M$  and  $N$  be two  $\mathcal{RML}$ s and  $\varphi : M \rightarrow N$  be a surjective homomorphism. If  $f_A$  is an  $f$ -soft- $\mathcal{RML}$  over  $M$ , then  $\varphi f_A$  is an  $f$ -soft- $\mathcal{RML}$  over  $N$ .*

*Proof.* Suppose  $f_A$  is an  $f$ -soft- $\mathcal{RML}$  over  $M$  and let  $p \in A$ . We have to show that  $\varphi f(p)$  is an  $f$ -sub- $\mathcal{RML}$  of  $N$ . As  $f_A$  is an  $f$ -soft- $\mathcal{RML}$ , we have  $\top \in f(p)$ , which implies  $\top = \varphi(\top) \in \varphi f(p)$ . Now, let  $z, t \in \varphi f(p)$ . Then there exists  $x, y \in f(p)$  such that  $z = \varphi(x)$  and  $t = \varphi(y)$ . We have to show that  $z \sqcup t \subseteq \varphi(f(p))$  and  $z \sqcap t \subseteq \varphi(f(p))$ . As  $f_A$  is an  $f$ -soft residuated multilattice over  $M$ ,  $f(p)$  is an  $f$ -sub- $\mathcal{RML}$  of  $M$ ,  $x \sqcup y \subseteq f(p)$  and  $x \sqcap y \subseteq f(p)$  which implies that

$$\varphi(x \sqcup y) \subseteq \varphi(f(p)) \text{ and } \varphi(x \sqcap y) \subseteq \varphi(f(p)).$$

As  $\varphi$  is surjective,  $\varphi(x \sqcup y) = \varphi(x) \sqcup \varphi(y)$  and  $\varphi(x \sqcap y) = \varphi(x) \sqcap \varphi(y)$ . Thus we have  $\varphi(x) \sqcup \varphi(y) \subseteq \varphi(f(p))$  and  $\varphi(x) \sqcap \varphi(y) \subseteq \varphi(f(p))$  which implies that  $z \sqcup t \subseteq \varphi(f(p))$  and  $z \sqcap t \subseteq \varphi(f(p))$  as required. It remains to show that  $z \odot t, z \rightarrow t \in \varphi f(p)$ . Since  $f(p)$  is an  $f$ -sub- $\mathcal{RML}$  of  $M$ ,  $x \odot y \in f(p)$  and  $x \rightarrow y \in f(p)$ , i.e.,  $\varphi(x) \odot \varphi(y) = \varphi(x \odot y) \in \varphi f(p)$  and  $\varphi(x) \rightarrow \varphi(y) = \varphi(x \rightarrow y) \in \varphi f(p)$ . It follows that  $z \odot t \in \varphi f(p)$  and  $z \rightarrow t \in \varphi f(p)$ .  $\square$

**Proposition 5.10.** *Let  $\varphi : M \rightarrow N$  be a bijective homomorphism of residuated multilattices and  $f_A$  be an  $f$ -soft- $\mathcal{RML}$  over  $M$ . If  $h_F$  is a soft filter of  $f_A$ , then  $\varphi h_F$  is a soft filter of  $\varphi f_A$ .*

*Proof.* Suppose  $h_F$  is a soft filter of  $f_A$ . Clearly,  $\varphi f_A$  is an  $f$ -soft- $\mathcal{RML}$  over  $N$  according to Proposition 5.9.  $F \subseteq A$  stems from the fact that  $h_F$  is a soft filter of  $f_A$ . It remains to show that  $\varphi(h(p))$  is a filter of  $\varphi(f(p))$  for all  $p \in F$ . In fact, if  $p \in F$ , then  $h(p)$  is a filter of  $f(p)$ . Since  $f(p)$  is an  $\mathcal{RML}$  on it own, by applying Proposition 5.8,  $\varphi(h(p))$  is a filter of  $\varphi(f(p))$  as required.  $\square$

**Proposition 5.11.** *Let  $\varphi : M \rightarrow N$  be a bijective homomorphism of residuated multilattices. If  $f_A$  is a filteristic soft residuated multilattice over  $M$ , then  $\varphi f_A$  is a filteristic soft residuated multilattice over  $N$ .*

*Proof.* Let  $\varphi$  be a bijective homomorphism and suppose  $f_A$  is a filteristic soft residuated multilattice. Then  $f_A$  is non-null, which implies that  $\varphi f_A$  is a non-null soft set over  $N$ . For all  $p \in A$ ,  $f(p)$  is a filter of  $M$ , as  $f_A$  is a filteristic soft residuated multilattice. As  $\varphi$  is bijective, we deduce from Proposition 5.8 that  $\varphi(f(p))$  is a filter of  $N$ . It follows that  $\varphi f_A$  is a filteristic soft residuated multilattice over  $N$ .  $\square$

**Proposition 5.12.** *Let  $f_A$  be a filteristic soft residuated multilattice over an  $\mathcal{RML}$   $M$  and  $\varphi : M \rightarrow N$  be a bijective homomorphism of residuated multilattices.*

- (1) *If  $f(p) = \varphi^{-1}(\top)$  for all  $p \in A$ , then  $\varphi f_A$  is a trivial filteristic soft residuated multilattice over  $N$ .*
- (2) *If  $f_A$  is whole, then  $\varphi f_A$  is a whole filteristic soft residuated multilattice over  $N$ .*

*Proof.* Let  $f_A$  be a filteristic soft residuated multilattice over  $M$  and  $\varphi$  be a bijective homomorphism.

(1) Suppose  $f(p) = \varphi^{-1}(\top)$  for all  $p \in A$ . Then  $\varphi(f(p)) = \varphi(\varphi^{-1}(\top)) = \top$ . Thus  $\varphi f_A$  is a trivial filteristic soft residuated multilattice.

(2) Suppose  $f_A$  is whole. Then for all  $p \in A$ ,  $f(p) = M$ , that is  $\varphi(f(p)) = \varphi(M) = N$ . Thus  $\varphi f_A$  is a whole filteristic soft residuated multilattice over  $M$ .  $\square$

## 6. CONCLUSION

We have used residuated multilattices as universe sets within the framework of soft set theory, studying the notions of f-soft residuated multilattices and r-soft residuated multilattices. We introduced some related notions such as soft filters and filteristic soft residuated multilattices with illustrative examples. Some properties of these concepts are proven. We plan to investigate some applications of this study to formal concept analysis.

## REFERENCES

- [1] L. A. Zadeh, Fuzzy sets, *Information and Control* 8 (1965) 338–353.
- [2] W. L. Gau and D. J. Buehrer, Vague sets, *IEEE Transactions on Systems, Man and Cybernetics* 23 (2) (1993) 610–614.
- [3] Z. Pawlak, Rough sets, *International Journal of Computer & Information Sciences* 11 (1982) 341–356.
- [4] D. Molodtsov, Soft set theory-first results, *Computers & Mathematics with Applications* 37 (4-5) (1999) 19–31.
- [5] M. I. Ali, F. Feng, X. Liu, W. K. Min and M. Shabir, On some new operations in soft set theory, *Computers & Mathematics with Applications* 57 (9) (2009) 1547–1553.
- [6] P. K. Maji, R. Biswas and A. R. Roy, Soft set theory, *Computers & Mathematics with Applications* 45 (4-5) (2003) 555–562.
- [7] D. Chen, E. Tsang, D. S. Yeung and X. Wang, The parameterization reduction of soft sets and its applications, *Computers & Mathematics with Applications* 49 (5-6) (2005) 757–763.
- [8] B. Sai, Applications of soft set in decision making problems, *Global Journal of Pure and Applied Mathematics* 16 (2) (2020) 305–324.
- [9] H. Aktaş and N. Çağman, Soft sets and soft groups, *Inform. Sci.* 177 (13) (2007) 2726–2735.
- [10] U. Acar, F. Koyuncu and B. Tanay, Soft sets and soft rings, *Computers & Mathematics with Applications* 59 (11) (2010) 3458–3463.
- [11] F. Feng, Y. B. Jun, X. Zhao, Soft semirings. *Computers & Mathematics with Applications* 56(10) (2008) 2621–2628.
- [12] L. Fu, Soft lattices, *Global Journal of Science Frontier Research* 10 (4) (2010) 57–59.
- [13] F. Karaaslan, N. Çağman and S. Enginoglu, Soft lattices, *Journal of New Results in Science* 1 (1) (2012) 5–17.
- [14] Y. B. Jun, Soft *BCK/BCI*-algebras, *Computers & Mathematics with Applications* 56 (5) (2008) 1408–1413.
- [15] J. Zhan and Y. B. Jun, Soft *BL*-algebras based on fuzzy sets, *Computers & Mathematics with Applications* 59 (6) (2010) 2037–2046.
- [16] G. Dymek and A. Walendziak, Soft set theory applied to general algebras, *Demonstratio Mathematica* 48 (1) (2015) 13–20.
- [17] M. Benado, Les ensembles partiellement ordonnés et le théorème de raffinement de schreier ii théorie des multistructures, *Czechoslovak Mathematical Journal* 5 (3) (1955) 308–344.
- [18] I. P. Cabrera, P. Cordero, G. Gutiérrez, J. Martínez and M. Ojeda-Aciego, On residuation in multilattices: filters, congruences and homomorphisms, *Fuzzy Sets and Systems* 234 (2014) 1–21.
- [19] P. C. Kengne, D. C. Awouafack, B. B. Njionou and C. Lélé,  $\mathcal{L}$ -fuzzy ideals and  $\mathcal{L}$ -fuzzy congruences of multilattices, *Advances in Fuzzy Sets and Systems* 26 (2021) 145–174.

- [20] L. M. Nzoda, B. Koguep, C. Lélé and L. Kwuida, Fuzzy setting of residuated multilattices, Ann. Fuzzy Math. inform. 10 (6) (2015) 929–948.
- [21] D. L. A. Kouankam, N. B. B. Koguep and C. Lélé, Application of soft sets theory to multilattices, Ann. Fuzzy Math. Inform. 26 (3) (2023) 245–259.
- [22] D. J. Hansen, An axiomatic characterization of multilattices, Discrete Mathematics 33 (1) (1981) 99–101.
- [23] N. B. B. Koguep, L. Kwuida and C. Lélé, Formal concepts and residuation on multilattices, Fundamenta Informaticae 188 (4) (2023) 217–237.

KOUANKAM DJOUOHOU LÉA AUBIN ([akouankam@gmail.com](mailto:akouankam@gmail.com))

Department of Mathematics and Computer Science, Faculty of Science, University of Dschang, Cameroon

KOGUEP NJIONOU BLAISE BLÉRIOT ([knbbkoguep2011@gmail.com](mailto:knbbkoguep2011@gmail.com))

Department of Mathematics and Computer Science, Faculty of Science, University of Dschang, Cameroon

LÉLÉ CÉLESTIN ([prcelestinlele@gmail.com](mailto:prcelestinlele@gmail.com))

Department of Mathematics and Computer Science, Faculty of Science, University of Dschang, Cameroon